

# THE $K$ -GROUPS AND THE INDEX THEORY OF CERTAIN COMPARISON $C^*$ -ALGEBRAS

BERTRAND MONTHUBERT AND VICTOR NISTOR

ABSTRACT. We compute the  $K$ -theory of comparison  $C^*$ -algebra associated to a manifold with corners. These comparison algebras are an example of the abstract pseudodifferential algebras introduced by Connes and Moscovici [12]. Our calculation is obtained by showing that the comparison algebras are a homomorphic image of a groupoid  $C^*$ -algebra. We then prove an index theorem with values in the  $K$ -theory groups of the comparison algebra.

## CONTENTS

Introduction	1
1. Groupoids and comparison algebras	4
2. The analytic index	7
3. Properties of the full $C^*$ -analytic index	9
4. A topological index theorem	11
5. $K$ -theory of comparison algebras	12
References	12

## INTRODUCTION

The work of Henri Moscovici encompasses many areas of mathematics, most notably Non-commutative Geometry, Group Representations, Geometry, and Abstract Analysis. His work on Non-commutative geometry, mostly joint works with Alain Connes, has lead to many breakthroughs in Index Theory and Operator Algebras, as well as to applications to other areas. We are happy to dedicate this paper to Henri Moscovici on the occasion of his 65th birthday.

---

*Date:* August 22, 2011.

Monthubert was partially supported by a ACI Jeunes Chercheurs. Manuscripts available from <http://bertrand.monthubert.net>. Nistor was partially supported by the NSF Grant DMS 0555831. Manuscripts available from <http://www.math.psu.edu/nistor/>.

The problem studied in this paper pertains to the general program of understanding index theory on singular and non-compact spaces. On such spaces, the Fredholm property depends on more than the principal symbol, so cyclic cocycles are needed in order to obtain explicit index formulas. Moscovici has obtained many results in this direction, including [4, 6, 11, 12, 33, 34]. See also [5, 20, 8, 9, 38].

One of the central concepts in a recent paper by Connes and Moscovici, is that of an abstract algebra of pseudodifferential operators [12]. These algebras generalize similar algebras introduced earlier. In this paper, we would like to study certain natural  $C^*$ -algebras associated to non-compact Riemannian manifolds, applying in particular the point of view of the work of Connes and Moscovici mentioned above.

Let us now explain the framework of this paper. Let  $M_0$  be a complete Riemannian manifold and let  $\Delta = d^*d$  be the *positive* Laplace operator on  $M_0$  associated to the metric. It is well known that  $\Delta$  is essentially self-adjoint [13, 44] and the references therein, and hence we can define  $\Lambda = (1 + \Delta)^{-1/2}$  using functional calculus. Let us also assume that a certain algebra  $\mathcal{D} = \cup \mathcal{D}_n$  of differential operators is given on  $M_0$ , where  $\mathcal{D}_n$  denotes the space of differential operators in  $\mathcal{D}$  of degree at most  $n$ . Let us assume that  $\Delta \in \mathcal{D}_2$  and that  $L_n \Lambda^n$  defines a bounded operator on  $L^2(M_0)$  for any  $L_n \in \mathcal{D}_n$ . Then the *comparison algebra* of  $M_0$  (and  $\mathcal{D}$ ) to be the  $C^*$ -algebra generated by the operators of the form  $L_n \Lambda^n$  for any  $L_n \in \mathcal{D}_n$ . This definition is almost the same as the one in [14, 15], where the comparison algebra was defined as the  $C^*$ -algebra generated by all operators of the form  $L_1 \Lambda$  for any  $L_1 \in \mathcal{D}_1$  and all compact operators. One of our results, Theorem 4, implies that the two definitions are the same for suitable  $M_0$ . Let us denote the comparison  $C^*$ -algebra of  $M_0$  by  $\mathfrak{A}(M_0)$  (the dependence on the algebra  $\mathcal{D}$  will be implicit). The comparison algebra  $\mathfrak{A}(M_0)$  is a convenient tool to study many analytic properties of differential operators on  $M_0$ , such as invertibility between Sobolev spaces, spectrum, compactness, the Fredholm property, and the index [10, 14, 19, 23, 25, 45]. For instance, the principal symbol of order zero pseudodifferential operators extends to a continuous map  $\sigma_0 : \mathfrak{A}(M_0) \rightarrow C(S^*A)$  with kernel denoted  $\mathfrak{A}_{-1}(M_0)$ , where  $S^*A$  is a suitable compactification of the co-sphere bundle of  $T^*M_0$ .

In this paper, we concentrate on the index properties of elliptic operators on a certain class of non-compact manifolds, called “manifolds with poly-cylindrical ends.” Recall that a *manifold with poly-cylindrical ends* is, locally, a product of manifolds with cylindrical ends. Our index depends only on the principal symbol, so it takes values in the  $K$ -theory

of the  $C^*$ -algebra  $\mathfrak{A}_{-1}(M_0)$ . Index calculations are sometimes necessary in applications, for instance in the study of Hartree's equation [21] and in the study of boundary value problems on polyhedral domain [26].

More precisely, let us first assume that the given manifold  $M_0$  is the interior of the space of units of a groupoid Lie  $\mathcal{G}$ . Then the Lie groupoid structure of  $\mathcal{G}$  gives rise to a natural algebra of differential operators  $\mathcal{D}$  on  $M_0$ , such as in the case of singular foliations [10, 2]. In general, there will be no natural metric on  $M_0$ , and even if a metric is chosen on  $M_0$ , the associated Laplace operator  $\Delta \notin \mathcal{D}$ . However, if  $M_0$  is a Lie manifold [1], then a natural class of metrics exists on  $M_0$  and  $\Delta \in \mathcal{D}$  for any metric in this class. Recall that  $M_0$  is a *Lie manifold* if the tangent bundle  $TM_0$  extends to a bundle  $A \rightarrow M$  on a compactification  $M$  of  $M_0$  to a manifold with corners such that the space of smooth section  $\mathcal{V} := \Gamma(A)$  of  $A$  has a natural Lie algebra structure induced by the Lie bracket of vector fields and such that that the diffeomorphisms generated by vector fields in  $\mathcal{V}$  preserve the faces of  $M$  [1].

We shall show that the comparison algebra of a Lie manifold  $M_0$  identifies with a subalgebra of a homomorphic image of a groupoid (pseudodifferential) algebra. For any manifold with corners, we shall denote by  $\mathcal{V}_M$  the Lie algebra of all vector fields tangent to all faces of  $M$ . Then  $\mathcal{V}_M = \Gamma(A_M)$  for a unique (up to isomorphism) vector bundle  $A_M \rightarrow M$ . If the vector bundle  $A \rightarrow M$  defining a Lie manifold  $M$  satisfies  $A = A_M$ , then we shall say that  $M_0$ , the interior of  $M$ , is a *manifold with poly-cylindrical ends*. In that case, we prove that  $\mathfrak{A}(M_0)$  is (isomorphic to) the norm closure of  $\Psi^0(\mathcal{G})$ . We then use this result to compute the  $K$ -theory of the algebra  $\mathfrak{A}_{-1}(M_0)$  and the index in  $K_0(\mathfrak{A}_{-1}(M_0))$  of elliptic operators in the comparison algebra of a manifold with poly-cylindrical ends.

Let us explain in a little more detail our results. Let us assume that our algebra  $\mathcal{D}$  of differential operators is generated by  $\mathcal{C}^\infty(M)$  and  $\mathcal{V}$ . Let  $D \in \mathcal{D}$ , then the principal symbol of  $D$  extends to a symbol defined on  $A^*$ . Assume that  $D$  is elliptic, in the sense that its principal symbol is invertible on  $A^*$  outside the zero section. Then the  $K$ -theory six-term exact sequence applied to the tangent (or adiabatic) groupoid of  $\mathcal{G}$  defines a map

$$(1) \quad \text{ind}_a = \text{ind}_a^M : K^0(A^*) \rightarrow K_0(\mathfrak{A}_{-1}(M_0)).$$

One of our main results is a computation of the groups  $K_0(\mathfrak{A}_{-1}(M_0))$  and of the map  $\text{ind}_a$  in case  $M_0$  is a manifold with poly-cylindrical ends.

Our calculation of the group  $K_0(\mathfrak{A}_{-1}(M_0))$  is as follows. Consider an embedding  $\iota : M \rightarrow X$  of manifolds with corners and let  $\iota_!$  be the push-forward map in  $K$ -theory. Morita equivalence then gives rise to a morphism  $\iota_* : K_0(C^*(M)) \rightarrow K_0(C^*(X))$ . Then Theorem 7 states that the following diagram commutes

$$(2) \quad \begin{array}{ccc} K_0(C^*(M)) & \xrightarrow{\iota_*} & K_0(C^*(X)) \\ \text{ind}_a^M \uparrow & & \uparrow \text{ind}_a^X \\ K^0(A_M^*) & \xrightarrow{\iota_!} & K^0(A_X^*). \end{array}$$

If the manifold with corners  $X$  is such that the natural morphisms  $\iota_* : K_0(C^*(M)) \rightarrow K_0(C^*(X))$  and  $\text{ind}_a^X : K^0(A_X^*) \rightarrow K_0(C^*(X))$  are isomorphisms, we are going to say that  $X$  is a *classifying space* for  $M$ . In that case, we can interpret the above diagram as a topological index theorem in the usual sense. We also obtain an identification of the groups  $K_0(C^*(M))$  and of the map  $\text{ind}_a^M$ .

Let us now very briefly summarize the contents of the paper. In Section 1, we introduce comparison algebras and we show that they are closely related to groupoid algebras. We show that the groupoid  $C^*$ -algebra  $\overline{\Psi^0(\mathcal{G}_M)}$  and the comparison  $C^*$ -algebra  $\mathfrak{A}(M_0)$  are in fact isomorphic for  $M_0$  a manifold with poly-cylindrical ends with compactification  $M$ . In Section 2, we recall the definition of the full  $C^*$ -analytic index using the tangent groupoid. In the process, we establish several technical results on tangent groupoids. Section 3 contains the main properties of the full  $C^*$ -analytic index. In this section, we also introduce the morphism  $j_*$  associated to an embedding of manifolds with corners  $j$  and we provide conditions for  $j_*$  and  $\text{ind}_a^M$  to be isomorphisms. We also discuss the compatibility of the full  $C^*$ -analytic index and of the shriek maps. This is then used to establish the equality of the full  $C^*$ -analytic and principal symbol topological index. Some of the proofs in this paper are only sketched. See [31] for full details.

We thank Bernd Ammann, Catarina Carvalho, Sergiu Moroianu, and Georges Skandalis for useful discussions. The second named author would like to thank the Max Planck Institute for Mathematics Bonn, where part of this work was completed, for hospitality and support.

## 1. GROUPOIDS AND COMPARISON ALGEBRAS

We shall need to consider pseudodifferential operators on groupoids [32, 40]. For simplicity, *we shall assume from now on that all our manifolds with corners have embedded faces.*

**1.1. Pseudodifferential operators on groupoids.** Throughout this paper, we shall fix a Lie groupoid  $\mathcal{G}$  with units  $M$  and Lie algebroid  $A = A(\mathcal{G})$ . Here  $M$  is allowed to have corners. To  $\mathcal{G}$ , there is associated the pseudodifferential calculus  $\Psi^\infty(\mathcal{G})$ , whose operators of order  $m$  form a linear space denoted  $\Psi^m(\mathcal{G})$ ,  $m \in \mathbb{R}$ , see [32, 40]. For short, this calculus is defined as follows. Let  $s : \mathcal{G} \rightarrow M$  be the source map and  $\mathcal{G}_x = s^{-1}(x)$ . Then  $\Psi^m(\mathcal{G})$ ,  $m \in \mathbb{Z}$ , consists of smooth families of classical, order  $m$  pseudodifferential operators  $(P_x \in \Psi^m(\mathcal{G}_x))$ ,  $x \in M$ , that are right invariant with respect to multiplication by elements of  $\mathcal{G}$  and are “uniformly supported.” To define what uniformly supported means, let us observe that the right invariance of the operators  $P_x$  implies that their distribution kernels  $K_{P_x}$  descend to a distribution  $k_P \in I^m(\mathcal{G}, M)$  [29, 40]. Then the family  $P = (P_x)$  is called *uniformly supported* if, by definition,  $k_P$  has compact support in  $\mathcal{G}$ .

We then have the following result [23, 30, 40].

**Theorem 1.** *The space  $\Psi^\infty(\mathcal{G})$  is a filtered algebra, closed under adjoints, so that the usual principal symbol of pseudodifferential operators defines a surjective*

$$\sigma_{\mathcal{G}}^{(m)} : \Psi^m(\mathcal{G}) \rightarrow S_{cl}^m(A^*)/S_{cl}^{m-1}(A^*),$$

with kernel  $\Psi^{m-1}(\mathcal{G})$ , for any  $m \in \mathbb{Z}$ .

**1.2. Comparison algebras.** We shall denote by  $\pi$  the natural action of  $\Psi^\infty(\mathcal{G})$  on  $\mathcal{C}_c^\infty(M_0)$  or on its completions and by  $\mathfrak{a}_{-\infty}$  the completion of  $\pi(\Psi^{-\infty}(\mathcal{G}))$  acting on all Sobolev spaces  $H^{-m}(M_0) \rightarrow H^m(M_0)$ . Let us denote  $\mathfrak{a}_n := \pi(\Psi^n(\mathcal{G})) + \mathfrak{a}_{-\infty}$ . The following result was proved in [24] (see also [1, 25]).

**Theorem 2.** *The space  $\mathfrak{a} := \pi(\Psi(\mathcal{G})) + \mathfrak{a}_{-\infty}$  is a filtered algebra by the pseudodifferential degree such that  $\mathfrak{a}_0 := \pi(\Psi^0(\mathcal{G})) + \mathfrak{a}_{-\infty}$  consists of bounded operators, is closed under the adjoint, and is spectrally invariant. Moreover,  $\mathfrak{a}_{-\infty}$  is a two-sided ideal of  $\mathfrak{a}$  and*

$$\Lambda := (1 + \Delta)^{-1/2} \in \mathfrak{a}_{-1} := \pi(\Psi^{-1}(\mathcal{G})) + \mathfrak{a}_{-\infty}.$$

From the above theorem we obtain that the comparison algebra is a subalgebra of the norm closure of  $\mathfrak{a}_0$ .

**Theorem 3.** *Let  $M_0$  be a Lie manifold. Then we have that*

$$\mathfrak{A}(M_0) \subset \overline{\mathfrak{a}_0} = \overline{\pi(\Psi^0(\mathcal{G}))}.$$

Moreover,  $\mathfrak{A}(M_0)$  contains all compact operators.

*Proof.* Let  $L_n \in \mathcal{D}_n$ . Since  $\Lambda \in \mathfrak{a}_{-1}$  and  $\mathcal{D}_n \subset \mathfrak{a}_n := \pi(\Psi^{-1}(\mathcal{G})) + \mathfrak{a}_{-\infty}$ . It follows that  $L_n \Lambda^n \in \mathfrak{a}_0$ , and hence the result.

To show that  $\mathfrak{A}(M_0)$  contains the subalgebra of compact operators, let us notice first that  $e^{-t\Delta} \in \mathfrak{A}(M_0)$ , since it can be written as a function of  $\Lambda := (1 + \Delta)^{-1/2}$ . Since  $C_c^\infty(M_0) \subset C^\infty(M)$ , we have that  $\phi e^{-t\Delta} \psi \in \mathfrak{A}(M_0)$ , and hence the latter contains compact operators. To show that all compact operators are in  $\mathfrak{A}(M_0)$ , it is enough to show that  $\mathfrak{A}(M_0)$  has no invariant subspaces. That is, it is enough to show that if  $f, g \in L_2(M_0)$  are such that the inner product  $(f, \phi e^{-t\Delta} \psi g)$  is zero for all  $\phi, \psi \in C_c^\infty(M_0)$ , then either  $f = 0$  or  $g = 0$ . Indeed, let us choose  $\phi_n, \psi_n \in C_c^\infty(M_0)$  such that  $\phi_n f \rightarrow |f|$  and  $\psi_n g \rightarrow |g|$  in  $L^2(M_0)$ . Then

$$(|f|, e^{-t\Delta} |g|) = \lim_{n \rightarrow \infty} (f, \phi_n e^{-t\Delta} \psi_n g) = 0,$$

which implies that either  $f = 0$  or  $g = 0$ , since the heat kernel  $e^{-t\Delta}$  has positive distribution kernel.  $\square$

Let  $M_0$  be a manifold with poly-cylindrical ends and  $\mathcal{G}$  be a groupoid such that  $A(\mathcal{G}) = A_M$ , where, we recall,  $A_M \rightarrow M$  is a vector bundle such that  $\mathcal{V}_M := \Gamma(A_M)$  consists of all smooth vector fields on  $M$  that are tangent to all the faces of  $M$ . A groupoid  $\mathcal{G}$  with this property is said *to integrate*  $A_M$ , and is not unique. However, if the fibers of the source map  $s : \mathcal{G} \rightarrow M$  are all connected and simply-connected, then  $\mathcal{G}$  is unique (up to isomorphism) and will be denoted  $\mathcal{G}_M$ . For  $\mathcal{G}_M$  the vector representation  $\pi$  is injective [25, 30]. We shall also denote by  $C^*(M) = C^*(\mathcal{G}_M)$ . Recall that  $C^*(\mathcal{G}_M) = \overline{\Psi^{-1}(\mathcal{G})}$ , [25, 30].

The algebra  $\Psi(\mathcal{G}_M)$  was considered before by many authors, including [27, 42, 43]. For this algebra, we actually have equality in the above theorem.

**Theorem 4.** *Let  $M_0$  be a manifold with poly-cylindrical ends. Then*

$$\mathfrak{A}(M_0) = \overline{\mathfrak{a}_0} \simeq \overline{\Psi^0(\mathcal{G}_M)}.$$

*Proof.* Let us recall that for manifolds with poly-cylindrical ends the vector representation  $\pi$  is injective on the norm closure  $\overline{\Psi^0(\mathcal{G}_M)}$ . We shall thus identify  $\Psi^0(\mathcal{G}_M)$  with  $\pi(\overline{\Psi^0(\mathcal{G}_M)})$ . Since the principal symbol map acting on both  $\mathfrak{A}(M_0)$  and on  $\overline{\Psi^0(\mathcal{G}_M)}$  has the same range, namely  $C(S^*A_M)$ , it is enough to show that  $\mathfrak{A}_{-1}(M_0) = \overline{\Psi^{-1}(\mathcal{G})} = C^*(M)$ .

Let us notice that we can consider families, so proving  $\mathfrak{A}_{-1}(M_0) = C^*(M)$  is equivalent to proving  $\mathfrak{A}_{-1}(M_0) \otimes C_0(X) = C^*(M) \otimes C_0(X)$ . Moreover, the inclusion  $\mathfrak{A}_{-1}(M_0) \subset C^*(M)$  is compatible with the natural representations of  $C^*(M)$  associated to the faces of  $M$ , as seen from their construction in [28]. It is enough then to prove that we have isomorphisms on subquotients defined by these representations,

which are all of the form  $\mathfrak{A}_{-1}(X_0) \otimes C_0(X)$ , for some lower dimensional manifolds. The proof finally reduces to show that  $\mathcal{K} \subset \mathfrak{A}_{-1}(M_0)$ , where  $\mathcal{K}$  is the algebra of compact operators. For this we use Theorem 3  $\square$

For manifolds with cylindrical ends (that is when  $M$  has no corners of codimension two or higher), this theorem was proved before in [16].

## 2. THE ANALYTIC INDEX

**2.1. The adiabatic and tangent groupoids.** For the definition and study of the full  $C^*$ -analytic index, we shall need the adiabatic and tangent groupoids associated to a differentiable groupoid  $\mathcal{G}$ . We now recall their definition.

Let  $\mathcal{G}$  be a Lie groupoid with space of units  $M$ . We construct both the *adiabatic groupoid*  ${}^{ad}\mathcal{G}$  and the *tangent groupoid*  ${}^T\mathcal{G}$  [10, 22, 23, 32, 41]. The space of units of  ${}^{ad}\mathcal{G}$  is  $M \times [0, \infty)$  and the tangent groupoid  ${}^T\mathcal{G}$  will be defined as the restriction of  ${}^{ad}\mathcal{G}$  to  $M \times [0, 1]$ . The underlying set of the groupoid  ${}^{ad}\mathcal{G}$  is the disjoint union:

$${}^{ad}\mathcal{G} = A(\mathcal{G}) \times \{0\} \cup \mathcal{G} \times (0, \infty).$$

We endow  $A(\mathcal{G}) \times \{0\}$  with the structure of commutative bundle of Lie groups induced by its vector bundle structure. We endow  $\mathcal{G} \times (0, \infty)$  with the product groupoid structure. Then the groupoid operations of  ${}^{ad}\mathcal{G}$  are such that  $A(\mathcal{G}) \times \{0\}$  and  $\mathcal{G} \times (0, \infty)$  are subgroupoids with the induced structure. Now let us endow  ${}^{ad}\mathcal{G}$  with a differentiable structure. To do so, it is enough to specify  $A({}^{ad}\mathcal{G})$ , since its knowledge completely determines the differentiable structure of  ${}^{ad}\mathcal{G}$  [39]. Then

$$(3) \quad \Gamma(A({}^{ad}\mathcal{G})) = t\Gamma(A(\mathcal{G} \times [0, \infty))).$$

More precisely, consider the product groupoid  $\mathcal{G} \times [0, \infty)$  with pointwise operations. Then a section  $X \in \Gamma(A(\mathcal{G} \times [0, \infty)))$  can be identified with a smooth function  $[0, \infty) \ni t \rightarrow X(t) \in \Gamma(A(\mathcal{G}))$ . We then require  $\Gamma(A({}^{ad}\mathcal{G})) = \{tX(t)\}$ , with  $X \in \Gamma(A(\mathcal{G} \times [0, \infty)))$ .

It is easy to show that

**Lemma 1.** *Let  $\mathcal{H} = \mathcal{G} \times \mathbb{R}^n$ , as above. We have that  $C^*(\mathcal{H}_{ad}) \simeq C^*(\mathcal{G}_{ad}) \otimes \mathcal{C}_0(\mathbb{R}^n)$  and that  $C^*({}^T\mathcal{H}) \simeq C^*({}^T\mathcal{G}) \otimes \mathcal{C}_0(\mathbb{R}^n)$ , the tensor product being the (complete, maximal)  $C^*$ -tensor product.*

**2.2. The full  $C^*$ -analytic index.** For each  $t \in [0, 1]$ ,  $M \times \{t\}$  is a closed invariant subset of  $M \times [0, \infty)$  for the adiabatic and tangent groupoids, and hence we obtain an *evaluation morphism*

$$e_t : C^*({}^T\mathcal{G}) \rightarrow C^*({}^T\mathcal{G}_{M \times \{t\}}),$$

and, in particular, an exact sequence

$$(4) \quad 0 \rightarrow C^*(T\mathcal{G}_{M \times (0,1]}) \rightarrow C^*(T\mathcal{G}) \xrightarrow{e_0} C^*(A(\mathcal{G})) \rightarrow 0.$$

Since  $K_*(C^*(T\mathcal{G}_{M \times (0,1]})) = K_*(C^*(\mathcal{G}) \otimes \mathcal{C}_0((0,1])) = 0$ , the evaluation map  $e_0$  induces an isomorphism in  $K$ -theory.

The  $C^*$ -algebra  $C^*(A(\mathcal{G}))$  is commutative and we have  $C^*(A(\mathcal{G})) \simeq \mathcal{C}_0(A^*(\mathcal{G}))$ . Therefore  $K_*(C^*(A(\mathcal{G}))) = K^*(A^*(\mathcal{G}))$ . In turn, this isomorphism allows us to define the *full  $C^*$ -analytic index*  $\text{ind}_a$  as the composition map

$$(5) \quad \text{ind}_a^{\mathcal{G}} = e_1 \circ e_0^{-1} : K^*(A^*(\mathcal{G})) \rightarrow K_*(C^*(\mathcal{G})),$$

where  $e_1 : C^*(T\mathcal{G}) \rightarrow C^*(T\mathcal{G}_{M \times \{1\}}) = C^*(\mathcal{G})$  is defined by the restriction map to  $M \times \{1\}$ . The definition of the full  $C^*$ -analytic index gives the following.

**Proposition 1.** *Let  $\mathcal{G}$  be a Lie groupoid with Lie algebroid  $\pi : A(\mathcal{G}) \rightarrow M$ . Also, let  $N \subset F \subset M$  be a closed, invariant subset which is an embedded submanifold of a face  $F$  of  $M$ . Then the full  $C^*$ -analytic index defines a morphism of the six-term exact sequences associated to the pair  $(A^*(\mathcal{G}), \pi^{-1}(N))$  and to the ideal  $C^*(\mathcal{G}_{N^c}) \subset C^*(\mathcal{G})$ ,  $N^c := M \setminus N$*

$$\begin{array}{ccccccc} K^0(\pi^{-1}(N^c)) & \longrightarrow & K^0(A^*(\mathcal{G})) & \longrightarrow & K^0(\pi^{-1}(N)) & \longrightarrow & K^1(\pi^{-1}(N^c)) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ K^0(C^*(\mathcal{G}_{N^c})) & \longrightarrow & K^0(C^*(\mathcal{G})) & \longrightarrow & K^0(C^*(\mathcal{G}_N)) & \longrightarrow & K^1(C^*(\mathcal{G}_{N^c})) \end{array}$$

*Proof.* The six-term, periodic long exact sequence in  $K$ -theory associated to the pair  $(A^*(\mathcal{G}), \pi^{-1}(N))$  is naturally isomorphic to the six-term exact sequence in  $K$ -theory associated to the pair  $\mathcal{C}_0(A^*_{M \setminus N}) \subset \mathcal{C}_0(A^*(\mathcal{G}))$ . The result follows from the naturality of the six-term exact sequence in  $K$ -theory and the definition of the full  $C^*$ -analytic index (6).  $\square$

Recall that for  $M$  a smooth manifold with corners with embedded faces, we have denoted  $A(\mathcal{G}_M) = A_M$  and  $C^*(M) := C^*(\mathcal{G}_M)$ . Then the full  $C^*$ -analytic index becomes the desired map

$$(6) \quad \text{ind}_a^M : K^*(A_M^*) \rightarrow K_*(C^*(M)).$$

See [17, 18] for more properties of the analytic index.

*Remark 5.* Assume  $M$  has no corners (or boundary). Then  $\mathcal{G}_M = M \times M$  is the product groupoid and hence  $\Psi^\infty(\mathcal{G}_M) = \Psi^\infty(M)$ . In particular,  $C^*(M) := C^*(\mathcal{G}_M) \simeq \mathcal{K}$ , the algebra of compact operators on  $M$ . In this case  $K_0(C^*(M)) = \mathbb{Z}$ , and  $\text{ind}_a$  is precisely the analytic



index as introduced by [3]. This construction holds also for the case when  $M$  is not compact, if one uses pseudodifferential operators of order zero that are “multiplication at infinity,” as in [7].

### 3. PROPERTIES OF THE FULL $C^*$ -ANALYTIC INDEX

The following proposition is an important step in the proof of our index theorem, Theorem 8.

**Proposition 2.** *Let  $X$  be a manifold with embedded faces such that each open face of  $X$  is diffeomorphic to a Euclidean space. Then the full  $C^*$ -analytic index*

$$\mathrm{ind}_a^X : K^*(A_X^*) \rightarrow K_*(C^*(X)),$$

*defined in Equation (6), is an isomorphism.*

*Proof.* The proof is by induction on the number of faces of  $X$  using Proposition 1, the six-term exact sequence in  $K$ -theory, and the Five Lemma in homological algebra.  $\square$

*Remark 6.* The above proposition can be regarded as a Baum–Connes isomorphism for manifolds with corners.

**Proposition 3.** *Let  $\iota : M \rightarrow X$  be a closed embedding of manifold with corners. Assume that, for each open face  $F$  of  $X$ , the intersection  $F \cap M$  is a non-empty open face of  $M$  and that every open face of  $M$  is obtained in this way. Then  $K_*(C^*(M)) \rightarrow K_*(C^*(X))$  is an isomorphism denoted  $\iota_*$ .*

*Proof.* Recall from [35] that two locally compact groupoids  $G$  and  $H$  are *equivalent* provided there exists a topological space  $\Omega$  and two continuous, surjective open maps  $r : \Omega \rightarrow \mathcal{G}^{(0)}$  and  $d : \Omega \rightarrow H^{(0)}$  together with a left (respectively right) action of  $G$  (respectively  $H$ ) on  $\Omega$  with respect to  $r$  (respectively  $d$ ), such that  $r$  (respectively  $d$ ) is a principal fibration of structural groupoid  $H$  (respectively  $G$ ). An important theorem of Muhly–Renault–Williams states that if  $G$  and  $H$  are equivalent, then  $K_*(C^*(G)) \simeq K_*(C^*(H))$  [35].

Our result then follows from the fact that  $\Omega := r^{-1}(M)$  establishes the desired equivalence between  $\mathcal{G}_M$  and  $\mathcal{G}_X$ .  $\square$

We can now prove a part of our principal symbol topological index theorem, Theorem 8, involving an embedding  $\iota : M \rightarrow X$  of our manifold with corners  $M$  into another manifold with corners  $X$ . This theorem amounts to the fact that the diagram (2) is commutative. In order to prove this, we shall first consider a tubular neighborhood

$$(7) \quad M \xrightarrow{k} U \xrightarrow{j} X$$

of  $M$  in  $X$ , so that  $\iota = j \circ k$ . The diagram (2) is then decomposed into the two diagrams below, and hence the proof of the commutativity of the diagram (2) reduces to the proof of the commutativity of the two diagrams below, whose morphisms are defined as follows: the morphism  $k_*$  is defined by Proposition 3 and the morphism  $j_*$  is defined by the inclusion of algebras. The morphism  $\iota_*$  is defined by  $\iota_* = j_* \circ k_*$ . Finally, the morphism  $k_!$  is the push-forward morphism.

Let us now turn our attention to the following diagram:

$$(8) \quad \begin{array}{ccccc} K_*(C^*(M)) & \xrightarrow{k_*} & K_*(C^*(U)) & \xrightarrow{j_*} & K_*(C^*(X)) \\ \text{ind}_a^M \uparrow & & \text{ind}_a^U \uparrow & & \uparrow \text{ind}_a^X \\ K^*(A_M^*) & \xrightarrow{k_!} & K^*(A_U^*) & \xrightarrow{j_!} & K^*(A_X^*). \end{array}$$

The commutativity of the left diagram is part of the following proposition, which is the most technical part of the proof. Its proof is obtained by integrating a Lie algebroid obtained as a double deformation of a tangent space.

**Proposition 4.** *Let  $\pi : U \rightarrow M$  be a vector bundle over a manifold with corners  $M$  and let  $k : M \rightarrow U$  be the “zero section” embedding. Then the following diagram commutes:*

$$(9) \quad \begin{array}{ccc} K_*(C^*(M)) & \xrightarrow[k_*]{\simeq} & K_*(C^*(U)) \\ \text{ind}_a^M \uparrow & & \uparrow \text{ind}_a^U \\ K^*(A_M^*) & \xrightarrow[k_!]{\simeq} & K^*(A_U^*) \end{array}$$

The commutativity of the second square in the Diagram 8 follows from the naturality of the tangent groupoid construction.

**Proposition 5.** *Let  $j : U \rightarrow X$  be the inclusion of the open subset  $U$ . Then the diagram below commutes:*

$$\begin{array}{ccc} K_*(C^*(U)) & \xrightarrow{j_*} & K_*(C^*(X)) \\ \text{ind}_a^U \uparrow & & \uparrow \text{ind}_a^X \\ K^*(A_U^*) & \xrightarrow[j_*]{\simeq} & K^*(A_X^*). \end{array}$$

As explained above, the previous two propositions give

**Theorem 7.** *Let  $M \xrightarrow{\iota} X$  be a closed embedding of manifolds with corners. Then the diagram*

$$(10) \quad \begin{array}{ccc} K_*(C^*(M)) & \xrightarrow{\iota_*} & K_*(C^*(X)) \\ \uparrow \text{ind}_a^M & & \text{ind}_a^X \uparrow \\ K^*(A_M^*) & \xrightarrow{\iota_!} & K^*(A_X^*) \end{array}$$

is commutative.

#### 4. A TOPOLOGICAL INDEX THEOREM

Motivated by Theorem 7 and by the results of Section 3 (see Propositions 2 and 3) we introduce the following definition.

**Definition 1.** *A strong classifying manifold  $X_M$  of  $M$  is a compact manifold with corners  $X_M$ , together with a closed embedding  $\iota : M \rightarrow X_M$  with the following properties:*

- (i) *each open face of  $X_M$  is diffeomorphic to a Euclidean space,*
- (ii)  *$F \rightarrow F \cap M$  induces a bijection between the open faces of  $X_M$  and  $M$ .*

Note that if  $M \subset X_M$  are as in the above definition, then each face of  $M$  is the transverse intersection of  $M$  with a face of  $X_M$ .

**Proposition 6.** *Let  $M$  be a manifold with embedded faces with embedded faces, and  $\iota : M \hookrightarrow X_M$  be a strong classifying space of  $M$ . Then the maps  $\iota_*$  and  $\text{ind}_a^X$  of Theorem 7 are isomorphisms. That is, a strong classifying space for  $M$  is a classifying space for  $M$ .*

*Proof.* This was proved in Propositions 2 and 3. □

Let  $\iota : M \rightarrow X_M$  be a classifying space for  $M$ . The above proposition then allows us to define (see diagram 10)

$$\text{ind}_t^M := \iota_*^{-1} \circ \text{ind}_a^X \circ \iota_! : K^*(A_M^*) \rightarrow K_*(C^*(M)).$$

If  $M$  is a smooth compact manifold (so, in particular,  $\partial M = \emptyset$ ), then  $C^*(M) = \mathcal{K}$ , the algebra of compact operators on  $L^2(M)$  and hence  $K_0(C^*(M)) = \mathbb{Z}$ . Any embedding  $\iota : M \hookrightarrow \mathbb{R}^N$  will then be a classifying space for  $M$ . Moreover, for  $X = \mathbb{R}^n$ , the map  $\iota_*^{-1} \circ \text{ind}_a^X : K^*(TX) \rightarrow \mathbb{Z}$  is the inverse of  $j_! : K^0(pt) \rightarrow K^0(T\mathbb{R}^N)$  and hence  $\text{ind}_t^{\mathbb{R}^N} = (j_!)^{-1} \iota_!$ , which is the definition of the topological index from [3]. In view of this fact, we shall also call the map  $\text{ind}_t^M$  the *topological index* associated to  $M$ . Theorem 7 then gives the following result:

**Theorem 8.** *Let  $M$  be a manifold with corners and  $A_M$  and  $C^*(M)$  be the Lie algebroid and the  $C^*$ -algebra associated to  $M$ . Then the principal topological index map  $\text{ind}_t^M$  depends only on  $M$ , that is, it is independent of the classifying space  $X_M$ , and we have*

$$\text{ind}_t^M = \text{ind}_a^M : K^*(A_M^*) \rightarrow K_*(C^*(M)).$$

If  $M$  is a smooth compact manifold (*no boundary*), this recovers the Atiyah-Singer index theorem on the equality of the full  $C^*$ -analytic and principal symbol topological index [3].

### 5. $K$ -THEORY OF COMPARISON ALGEBRAS

The isomorphism  $K_*(C^*(M)) \simeq K^*(X_M)$  provides us with a way of determining the groups  $K_*(C^*(M))$ . In particular, we have completed the determination of the  $K$ -theory groups of the comparison algebra  $\mathfrak{A}_{-1}(M_0) = C^*(M)$ , if the interior of  $M$  is endowed with a metric making it a manifold with poly-cylindrical ends.

**Theorem 9.** *Let  $M$  be a manifold with corners and embedded faces. We have  $K_*(C^*(M)) \simeq K^*(X_M)$ . Moreover  $K_j(C^*(M)) \otimes \mathbb{Q} \simeq \mathbb{Q}^{p_j}$ , where  $p_j$  is the number of faces of  $M$  of dimension  $\equiv j$  modulo 2.*

The last part of the above theorem is proved by showing that the Atiyah-Hirzebruch spectral sequence of  $X_M$  collapses at  $E^2$ . This is part of a joint work with Etienne Fieux.

It is not difficult to construct a classifying manifold  $X_M$  of  $M$  [31], that is, a manifold such that  $\iota_* : K_0(C^*(M)) \rightarrow K_0(C^*(X))$  and  $\text{ind}_a^X : K^0(A_X^*) \rightarrow K_0(C^*(X))$  are isomorphisms. Let us assume  $M$  is compact with embedded faces. The space  $X_M$  is obtained from an embedding  $X \rightarrow [0, \infty)^N$  for some large  $N$ , and then by removing suitable hyperplanes from the boundary of  $[0, \infty)^N$  such that each face of  $M$  is the transverse intersection of  $M$  and of a face of  $X_M$ . See also [36, 37].

### REFERENCES

- [1] B. Ammann, R. Lauter, and V. Nistor. On the geometry of Riemannian manifolds with a Lie structure at infinity. *Int. J. Math. Math. Sci.*, (1-4):161–193, 2004.
- [2] I. Androulidakis and G. Skandalis. The holonomy groupoid of a singular foliation. *J. Reine Angew. Math.*, 626:1–37, 2009.
- [3] M. F. Atiyah and I. M. Singer. The index of elliptic operators, i. *Annals of Math.*, 87:484–530, 1968.
- [4] D. Barbasch and H. Moscovici.  $L^2$ -index and the Selberg trace formula. *J. Funct. Anal.*, 53(2):151–201, 1983.

- [5] M.-T. Benamèur and J. Heitsch. Index theory and non-commutative geometry. I. Higher families index theory. *K-Theory*, 33(2):151–183, 2004.
- [6] J. Brüning and H. Moscovici.  $L^2$ -index for certain Dirac-Schrödinger operators. *Duke Math. J.*, 66(2):311–336, 1992.
- [7] C. Carvalho. A topological approach to the cobordism invariance of the index. *K-Theory*, 2006.
- [8] A. Connes. Sur la théorie non commutative de l'intégration. In *Algèbres d'opérateurs (Sém., Les Plans-sur-Bex, 1978)*, volume 725 of *Lecture Notes in Math.*, pages 19–143. Springer, Berlin, 1979.
- [9] A. Connes. A survey of foliations and operator algebras. In *Operator algebras and applications, Part I (Kingston, Ont., 1980)*, volume 38 of *Proc. Sympos. Pure Math.*, pages 521–628. Amer. Math. Soc., Providence, R.I., 1982.
- [10] A. Connes. *Noncommutative Geometry*. Academic Press, Inc., 1994.
- [11] A. Connes and H. Moscovici. Cyclic cohomology, the Novikov conjecture and hyperbolic groups. *Topology*, 29(3):345–388, 1990.
- [12] A. Connes and H. Moscovici. The local index formula in noncommutative geometry. *Geom. Funct. Anal.*, 5(2):174–243, 1995.
- [13] H. Cordes. On essential selfadjointness of powers and comparison algebras. *J. Anal. Math.*, 58:61–97, 1992. Festschrift on the occasion of the 70th birthday of Shmuel Agmon.
- [14] H. Cordes and R. McOwen. The  $C^*$ -algebra of a singular elliptic problem on a noncompact Riemannian manifold. *Math. Z.*, 153(2):101–116, 1977.
- [15] H. O. Cordes. *Spectral theory of linear differential operators and comparison algebras*, volume 76 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1987.
- [16] H. O. Cordes and S. H. Doong. The Laplace comparison algebra of spaces with conical and cylindrical ends. In *Pseudodifferential operators (Oberwolfach, 1986)*, volume 1256 of *Lecture Notes in Math.*, pages 55–90. Springer, Berlin, 1987.
- [17] C. Debord and J.-M. Lescure.  $K$ -duality for stratified pseudomanifolds. *Geom. Topol.*, 13(1):49–86, 2009.
- [18] C. Debord, J.-M. Lescure, and V. Nistor. Groupoids and an index theorem for conical pseudo-manifolds. *J. Reine Angew. Math.*, 628:1–35, 2009.
- [19] V. Georgescu. On the spectral analysis of quantum field Hamiltonians. *J. Funct. Anal.*, 245(1):89–143, 2007.
- [20] A. Gorokhovsky and J. Lott. Local index theory over foliation groupoids. *Adv. Math.*, 204(2):413–447, 2006.
- [21] E. Hunsicker, V. Nistor, and J. Sofo. Analysis of periodic Schrödinger operators: regularity and approximation of eigenfunctions. *J. Math. Phys.*, 49(8):083501, 21, 2008.
- [22] N. Landsman. Quantization and the tangent groupoid. In *Operator algebras and mathematical physics (Constanța, 2001)*, pages 251–265. Theta, Bucharest, 2003.
- [23] R. Lauter, B. Monthubert, and V. Nistor. Pseudodifferential analysis on continuous family groupoids. *Doc. Math.*, 5:625–655 (electronic), 2000.
- [24] R. Lauter, B. Monthubert, and V. Nistor. Spectral invariance for certain algebras of pseudodifferential operators. *J. Inst. Math. Jussieu*, 4(3):405–442, 2005.

- [25] R. Lauter and V. Nistor. Analysis of geometric operators on open manifolds: a groupoid approach. In *Quantization of singular symplectic quotients*, volume 198 of *Progr. Math.*, pages 181–229. Birkhäuser, Basel, 2001.
- [26] H. Li, A. Mazzucato, and V. Nistor. Analysis of the finite element method for transmission/mixed boundary value problems on general polygonal domains. *Electronic Transactions Numerical Analysis*, 37:41–69, 2010.
- [27] R. Melrose. *Geometric scattering theory*. Stanford Lectures. Cambridge University Press, Cambridge, 1995.
- [28] R. Melrose and V. Nistor.  $K$ -theory of  $C^*$ -algebras of  $b$ -pseudodifferential operators. *Geom. Funct. Anal.*, 8(1):88–122, 1998.
- [29] B. Monthubert. *Groupoïdes et calcul pseudo-différentiel sur les variétés à coins*. PhD thesis, Université Paris 7, 1998.
- [30] B. Monthubert. Pseudodifferential calculus on manifolds with corners and groupoids. *Proc. Amer. Math. Soc.*, 127(10):2871–2881, 1999.
- [31] B. Monthubert and V. Nistor. A principal symbol topological index theorem for manifolds with corners. Preprint.
- [32] B. Monthubert and F. Pierrot. Indice analytique et groupoïdes de Lie. *C. R. Acad. Sci. Paris Sér. I Math.*, 325(2):193–198, 1997.
- [33] H. Moscovici and R. Stanton.  $R$ -torsion and zeta functions for locally symmetric manifolds. *Invent. Math.*, 105(1):185–216, 1991.
- [34] H. Moscovici and A. Verona. Harmonically induced representations of nilpotent Lie groups. *Invent. Math.*, 48(1):61–73, 1978.
- [35] P. Muhly, J. Renault, and D. Williams. Equivalence and isomorphism for groupoid  $C^*$ -algebras. *J. Operator Theory*, 17(1):3–22, 1987.
- [36] V. Nazaïkinskiï, A. Savin, and B. Sternin. Elliptic theory on manifolds with corners. I. Dual manifolds and pseudodifferential operators. In  *$C^*$ -algebras and elliptic theory II*, Trends Math., pages 183–206. Birkhäuser, Basel, 2008.
- [37] V. Nazaïkinskiï, A. Savin, and B. Sternin. Noncommutative geometry and the classification of elliptic operators. *Sovrem. Mat. Fundam. Napravl.*, 29:131–164, 2008.
- [38] R. Nest and B. Tsygan. Algebraic index theorem. *Comm. Math. Phys.*, 172(2):223–262, 1995.
- [39] V. Nistor. Groupoids and the integration of Lie algebroids. *J. Math. Soc. Japan*, 52:847–868, 2000.
- [40] V. Nistor, A. Weinstein, and P. Xu. Pseudodifferential operators on groupoids. *Pacific J. Math.*, 189:117–152, 1999.
- [41] B. Ramazan. Limite classique de  $C^*$ -algèbres de groupoïdes de Lie. *C. R. Acad. Sci. Paris Sér. I Math.*, 329(7):603–606, 1999.
- [42] E. Schrohe and B.-W. Schulze. Boundary value problems in Boutet de Monvel’s algebra for manifolds with conical singularities. I. In *Pseudo-differential calculus and mathematical physics*, volume 5 of *Math. Top.*, pages 97–209. Akademie Verlag, Berlin, 1994.
- [43] B. Schulze. *Boundary value problems and singular pseudo-differential operators*. Wiley-Interscience Series in Pure and Applied Mathematics. Chichester: John Wiley & Sons., 1998.
- [44] R. Strichartz. Harmonic analysis as spectral theory of Laplacians. *J. Funct. Anal.*, 87(1):51–148, 1989.

- [45] M. Taylor. Gelfand theory of pseudo differential operators and hypoelliptic operators. *Trans. Amer. Math. Soc.*, 153:495–510, 1971.

UNIVERSITÉ PAUL SABATIER (UFR MIG), INSTITUT DE MATHÉMATIQUES DE  
TOULOUSE, F-31062 TOULOUSE CEDEX 4

*E-mail address:* `bertrand.monthubert@math.univ-toulouse.fr`

PENNSYLVANIA STATE UNIVERSITY, MATH. DEPT., UNIVERSITY PARK, PA  
16802

*E-mail address:* `nistor@math.psu.edu`