

# Propagation and quenching in a reactive Burgers-Boussinesq system

Peter Constantin\*      Jean-Michel Roquejoffre†      Lenya Ryzhik ‡  
Natalia Vladimirova §

July 26, 2007

## Abstract

We investigate the qualitative behavior of solutions of a Burgers-Boussinesq system – a reaction-diffusion equation coupled via gravity to a Burgers equation – by a combination of numerical, asymptotic and mathematical techniques. Numerical simulations suggest that when the gravity  $\rho$  is small the solutions decompose into a traveling wave and an accelerated shock wave moving in opposite directions. There exists  $\rho_{cr1}$  so that, when  $\rho > \rho_{cr1}$ , this structure changes drastically, and the solutions become more complicated. The solutions are composed of three elementary pieces: a wave fan, a combustion traveling wave, and an accelerating shock, the whole structure traveling in the same direction. There exists  $\rho_{cr2}$  so that when  $\rho > \rho_{cr2}$ , the wave fan catches up with the accelerating shock wave and the solution is quenched, no matter how large was the support of the initial temperature. We prove that the three building blocks (wave fans, combustion traveling waves and shocks) exist and we construct asymptotic solutions made up of these three elementary pieces. We finally prove, in a mathematically rigorous way, a quenching result irrespective of the size of the region where the temperature was above ignition – a major difference with what happens in advection-reaction-diffusion equations where an incompressible flow is imposed.

## 1 Introduction

In his pioneering papers [15, 16], Ya. I. Kanel made the following discovery: consider an initial value problem

$$T_t = T_{xx} + f(T),$$

on the real line,  $x \in \mathbb{R}$ , with the initial data  $T(0, x) = \chi_{[-L, L]}(x)$  which is the characteristic function of an interval  $[-L, L]$ . The nonlinearity  $f(T)$  is Lipschitz and of the ignition type: there exists  $\theta \in (0, 1)$  such that

$$\begin{aligned} f(T) &\equiv 0 && \text{on } [0, \theta] \cup \{1\} \\ f(T) &> 0 && \text{on } (\theta, 1), \end{aligned} \tag{1.1}$$

the range of  $T$  being the interval  $(0, 1)$ . Kanel has shown that there exists  $L_0$  so that if the initial "hot spot" size  $L$  satisfies  $L < L_0$  then there exists a time  $t_0 > 0$  so that  $0 < T(t_0, x) \leq \theta$  for all

---

\*Department of Mathematics, University of Chicago, Chicago, IL 60637; const@math.uchicago.edu

†Institut de Mathématiques de Toulouse (UMR CNRS 5219) and Institut Universitaire de France, Université Paul Sabatier, F-31062 Toulouse Cedex, France; roque@mip.ups-tlse.fr

‡Department of Mathematics, University of Chicago, Chicago, IL 60637; ryzhik@math.uchicago.edu

§ASC/Flash Centre, Department of Astronomy & Astrophysics, University of Chicago, Chicago, IL 60637; nata@flash.uchicago.edu

$x \in \mathbb{R}$  – hence, the reaction ceases at this time and solution decays to zero as  $t \rightarrow +\infty$ . We say that the solution *quenches* in that case. On the other hand, there exists  $L_1$  so that if  $L > L_1$  then the solution develops two traveling fronts, one going to the left, another to the right and  $T(t, x) \rightarrow 1$  as  $t \rightarrow +\infty$ , uniformly on compact sets. Very recently A. Zlatos has shown that  $L_0 = L_1$  [24].

The mathematical modeling of issues concerning flame propagation or quenching have received renewed attention recently, mainly regarding the effect that fluid flow has on the behavior discovered by Kanel: see, for instance, the direct simulations and formal asymptotic analysis in [12, 13, 14, 21]. A number of mathematically rigorous results generalizing Kanel’s results to ignition type reactions in the presence of a fluid flow are also available. Here is a typical example of a result of this kind: suppose that  $T(t, x, y)$  solves an advection-reaction-diffusion equation

$$\begin{aligned} T_t + u(x, y) \cdot \nabla T - \Delta T &= f(T), \\ \partial_\nu T &= 0, \text{ for } (x, y) \in \mathbb{R} \times \partial\Omega, \\ T(0, x, y) &= T_0(x, y), \end{aligned} \tag{1.2}$$

in a cylinder  $\Sigma = \{x, y \in \mathbb{R} \times \Omega\}$  where  $\Omega \subset \mathbb{R}^n$  is bounded and where  $f$  is a smooth ignition-type source term as in (1.1). Assume for simplicity that the initial datum  $T_0(x, y) = \chi_{[-L, L]}(x)$  depends only on the variable  $x$ , as in Kanel’s problem. Then there exists a constant  $L_0(u, f) > 0$  such that, if  $L < L_0(u, f)$ , then  $T(t, x, y)$  becomes uniformly smaller than  $\theta$  in finite time. This is an example of a finite time quenching. There also exists a constant  $L_1(u, f) \geq L_0(u, f)$  such that, if  $L > L_1$ , then  $T(t, x, y) \rightarrow 1$  as  $t \rightarrow +\infty$ , uniformly on compact sets in  $(x, y) \in \Sigma$ . It is not known whether  $L_0 = L_1$  when  $u \not\equiv 0$ . The main interest in these problems is in estimating the dependence of the quenching length  $L_0$  on the amplitude and geometry of the flow  $u(x, y)$ . In particular, precise results are known in advection-reaction-diffusion equations when a strong incompressible flow is imposed: see, for instance, [5, 17] for quenching by a strong shear flow and [11] for quenching by a strong cellular flow. In both cases the critical size  $L_0$  of an initially ”hot” region that can be quenched by the flow grows with the flow amplitude  $A$ , albeit at a rate depending on the flow geometry –  $L_0 \sim CA$  for generic shear flows and  $L_0 \sim CA^{1/4}$  in cellular flows. The increase in  $L_0$  is due to improved mixing by the incompressible flow.

The goal of the present paper is to investigate what happens when the fluid flow is no longer imposed, but rather obeys a hydrodynamic equation. What we wish to understand in this study is the following: what are the quenching rules when the reaction-diffusion is coupled to hydrodynamics? In particular, is quenching still a matter of the size of the zone where the temperature exceeds the ignition temperature? Such an investigation was initiated for reaction-diffusion equations coupled to incompressible hydrodynamics in the Boussinesq approximation in the spirit of [1, 2, 4, 6, 7, 9, 10, 18, 22, 23]. This was done in [8] for a bounded domain with Dirichlet boundary conditions. The cases of Neumann boundary conditions and of unbounded domains are still under investigation.

In this paper we are interested in the effects of a compressible flow on the quenching phenomenon, a subject that seems to not have been yet addressed in the mathematical literature. Studying this problem for the full compressible reactive Navier-Stokes problem is beyond our reach at the moment. We investigate therefore a drastically simplified model of a Boussinesq system, where the Navier-Stokes equation is replaced by the one dimensional Burgers equation coupled to a temperature equation by a gravity force:

$$\begin{aligned} T_t - T_{xx} + uT_x &= f(T) \\ u_t - \nu u_{xx} + uu_x &= \rho T. \end{aligned} \tag{1.3}$$

Here  $T(t, x)$  is the temperature and  $u(t, x)$  the velocity of the fluid. The reaction term  $f(T)$  satisfies

the assumptions (1.1), and moreover we require that

$$f'(1) < 0. \tag{1.4}$$

The coefficient  $\rho > 0$  represents the gravity, and the coefficient  $\nu \geq 0$  the kinematic viscosity.

The system (1.3) has another physical interpretation for a one-dimensional system of discrete excitable particles. The particles are mobile and inertial, can mix by diffusion, and can exchange momentum. A particle converts to an excited state if there is a high enough concentration of excited particles in its vicinity. Excited (and only excited) particles feel the presence of a force; all other properties of excited and non-excited particles are identical. Under some conditions the initially small excited region grows with time, as the excited particles spread around by diffusion and excite their neighbors. In addition, the driving force accelerates the excited particles and speeds up the process. However, if the force is too strong, the particles quickly spread around over a large area, their concentration drops below the threshold limit, and transition of new particles to the excited state terminates. Even though the particles excited earlier are still present in the system, we call this event extinction or quenching.

The continuum representation of the problem is the system (1.3);  $T(x)$  is the fraction of excited particles ( $0 \leq T \leq 1$ ), and  $u(x)$  is the locally averaged velocity. The system of Burgers and advection-reaction-diffusion equations describes the transport of momentum and the transport of excited species,  $\rho$  is the driving force, and  $f(T)$  is the reaction term which accounts for the transition of particles from non-excited to excited state.

The outcome of the present study is that the qualitative behavior of the reactive system under investigation is markedly different from that of a reactive system in a passive incompressible flow. In particular, if the parameter  $\rho$  is sufficiently large, then quenching may occur irrespective of the size of the set where  $T(0, x) \geq \theta$ . We note that, as in the case of an imposed flow, the temperature goes to zero as  $t \rightarrow +\infty$  as soon as it drops below  $\theta$  everywhere, provided that the flow is decaying at infinity.

The paper is organized as follows. In Section 2, we carry out a numerical investigation of the system (1.3). The subsequent mathematical analysis is based on these numerical computations: it would be very difficult for us to find the correct qualitative behavior of the solutions without them. A feature of the numerical simulations is that they are not very sensitive to the viscosity  $\nu$ ; therefore we set  $\nu = 0$  in the rest of the paper. Namely, we concentrate on the system

$$\begin{aligned} T_t - T_{xx} + uT_x &= f(T) \\ u_t + uu_x &= \rho T \end{aligned} \tag{1.5}$$

Briefly, the numerical simulations show the following picture for solutions with a sufficiently large set where initially  $T(0, x) = 1$ : there exists a critical value  $\rho_{cr1} > 0$  so that for  $\rho \in (0, \rho_{cr1})$  such solutions develop a left going traveling wave which moves with a constant speed. On the right boundary they have a shock wave accelerating in time to the right. When  $\rho > \rho_{cr1}$  the gravity does not permit a left-going traveling wave to develop. Instead, the solution is made up of three elementary building blocks pieced together: a wave fan in the back, followed by a traveling wave, and finally an accelerated shock. This whole structure propagates to the right. Finally, there exists a second critical threshold  $\rho_{cr2}$  so that for  $\rho > \rho_{cr2}$  the wave fan catches up with the shock, no matter how large was the support of  $T(0, x)$  and the reaction stops from this time onward: it is quenched. This seems to be the main difference between active compressible and passive incompressible flows – when a compressible flow is sufficiently strong, all solutions are quenched, regardless of their initial size. It would be very interesting to investigate this phenomenon in more realistic reactive flow models. The numerical results are presented in Section 2.

The next three sections are devoted to the mathematical study of the elementary solutions of the system (1.5). In Section 3 we establish the existence of wave fans, that is, self-similar solutions of (1.5) with  $f(T) \equiv 0$ . Suppressing the diffusivity in the temperature equation yields explicit wave fans; these can be seen in Section 2. Establishing their existence when the temperature diffusivity is positive turns out to be a surprisingly difficult task: the whole program is carried out in full details in Section 3.

In Section 4 we prove existence and study qualitative properties of the combustion traveling waves. We expect similar results to hold for  $\nu \neq 0$  as well, although the equations are different.

In Section 5, we construct asymptotic solutions to the full system (1.5). We point out that what we prove here is that the 'solutions' that we have constructed only satisfy the system up to an error that is  $O(t^{-1})$ , and not that they are true solutions to the system. However, they are constructed by matched asymptotic expansions, and we believe that it is possible to construct true solutions to (1.5) on the basis of these asymptotic solutions. This latter investigation is not, however, in the scope of this paper, and will be carried out elsewhere. The constructed solutions fully account for what we saw in the numerics of Section 2.

In Section 6 we prove a quenching result. For  $\rho$  sufficiently large there is numerical evidence from Section 2 that the formal solution constructed in Section 5 will quench. We prove rigorously that taking the asymptotic solution as the initial data, the temperature will drop below the ignition temperature in a finite time that we are able to estimate.

**Acknowledgment.** This work has been supported by ASC Flash Center at the University of Chicago. PC was supported by NSF grant DMS-0504213, and LR by NSF grant DMS-0604687. This work was completed during visits by PC and LR to Université Paul Sabatier and by JMR to the University of Chicago. We thank these institutions for their hospitality.

## 2 Numerical simulations

In this section we investigate numerically the system (1.3) with  $\nu = 1$ ; in other words, we consider

$$\begin{aligned} T_t - T_{xx} + uT_x &= f(T) \\ u_t - u_{xx} + uu_x &= \rho T. \end{aligned} \tag{2.1}$$

To be specific, we choose the following piecewise linear reaction rate:

$$\begin{aligned} f(T) &= \frac{\theta(1-T)}{(1-\theta)^2}, & \theta < T < 1; \\ f(T) &= 0, & \text{otherwise,} \end{aligned} \tag{2.2}$$

where  $\theta$  is the ignition temperature. We set  $\theta = 1/2$  in most simulations below. Our analysis does not depend on this particular choice of reaction – it is important only that the ignition cutoff exists and the reaction vanishes for  $T > 1$  and  $T < 0$ . In the absence of advection ( $\rho = 0$ ,  $u = 0$ ) the reaction-diffusion equation for temperature with reaction rate (2.2) has a traveling wave solution moving with unit speed,

$$\begin{aligned} T^*(x, t) &= 1 - (1 - \theta)e^{\frac{1-\theta}{\theta}(x-t)}, & x < t; \\ T^*(x, t) &= \theta e^{-(x-t)}, & x > t. \end{aligned} \tag{2.3}$$

Here the location  $x = t$  corresponds to the point where temperature equals to the ignition threshold value  $\theta$ .




Figure 1: Quenching of small hot spot by diffusion alone. Thick solid lines represent the initial profiles with  $h = h^* = 1.50$  and  $k = 0.75$  (left), and with  $h = h^* = 1.00$  and  $k = 1.50$  (right). Thin dashed lines correspond to the solutions at time  $t = 2$  with the initial widths just below and just above critical ( $\Delta h = \pm 0.01$ ). The dotted line is the threshold value  $\theta = 0.5$ .

As an initial condition for  $T$  we use a smooth localized distribution of width  $h$  and steepness  $k$ ,

$$T(x, 0) = \frac{1}{2} \tanh[k(x + h/2)] - \frac{1}{2} \tanh[k(x - h/2)]. \quad (2.4)$$

We use mostly  $k = 0.75$ , which corresponds to a profile with a more gradual slope compared to the traveling wave (2.3), but that overall better matches the theoretical solution. In a few cases we use a steeper interface with  $k = 1.5$ . The initial velocity is zero in the majority of our simulations with the exception of the case where we study quenching by a prescribed initial velocity.

We solve the system (2.1) with the reaction rate (2.2) numerically, using fourth-order central differences discretization and third-order Adams-Bashforth integration in time. The simulations are done at resolution  $\Delta x_0 = 1/16$  with an adaptive patch of highly resolved mesh ( $\Delta x_1 = \Delta x_0/64$ ) in the vicinity of the shock.

## 2.1 Quenching without gravity

We first consider the simple case  $\rho = 0$ . If initially  $u = 0$ , then the problem is reduced to a single scalar reaction-diffusion equation. If  $u(0, x) \neq 0$  then the equation for  $T(t, x)$  is driven by the flow  $u(t, x)$  that satisfies the viscous Burgers equation. Much of the material of this section is well-known, and has received an extensive mathematical treatment. It is, however, instructive to put it here: it will serve as a comparison basis for more sophisticated effects appearing when the gravity is present.

### 2.1.1 Zero velocity - Kanel's length

In the absence of gravity and initial velocity, initial data (2.4) evolve in agreement with the theoretical predictions [16, 24]. Namely, initially sufficiently large hot regions,  $h > h^*$ , develop two outward propagating traveling waves of the form (2.3), while initially small hot regions,  $h < h^*$ , are extinguished.

The critical length of the initially hot region  $h^*$  – Kanel's length – can be determined numerically. For the given  $\theta$ , the critical width depends on the steepness of the interface. We show in Figure 1 the initial profiles corresponding to the Kanel's lengths  $h^* = 1.50$ , for  $k = 0.75$ , and  $h^* = 1.00$ , for  $k = 1.50$  (in both cases  $\theta = 0.5$ ). Note that for  $k = 0.75$ , the maximum in the profile only slightly

exceeds the ignition temperature,  $T_{\max} = 0.51$ , while for steeper  $k = 1.50$  the maximum is higher,  $T_{\max} = 0.64$ . The solution with critical width  $h^*$  is unstable. The solution with  $h$  just below critical decays with time, while the one just above critical grows and eventually develops into a pair of outward propagating fronts.

### 2.1.2 Stationary compression

When  $\rho = 0$ , the Burgers equation for  $u(t, x)$  is uncoupled from the advection-reaction-diffusion equation for  $T(t, x)$  and has a stationary solution,

$$u^*(x) = -U \tanh \frac{Ux}{2}, \quad (2.5)$$

where  $U$  is the absolute velocity at  $x = \pm\infty$ . The stationary solution represents compression, in the sense that  $u^* < 0$  for  $x > 0$  and  $u^* > 0$  for  $x < 0$ . Although the velocity remains unaffected by temperature, it changes the temperature distribution and can facilitate quenching.

In this exercise, we study the quenching of initial data with different  $h$  by the stationary velocity (2.5) with different intensities,  $U$ . Both  $u(x)$  and  $T(x)$  are aligned at  $x = 0$ , so that the compression is symmetric with respect to the center of the hot spot.

We find that there exists a critical velocity,  $U_{cr}$ , that quenches any initial distribution of temperature, no matter how wide it is. The independence of initial size is not surprising: if the initial distribution is wide, both fronts are located in the region where the velocity is nearly constant,  $u(x) \approx \pm U$ . The fronts are advected toward the center with the speed  $V \approx U - 1$ , and eventually reach the center. Near the center the absolute velocity is lower, and the decrease of temperature due to compression might or might not be balanced by reaction. We found that if  $U > U_{cr} = 1.40$  the maximum of temperature drops below the ignition threshold, that is, the hot spot completely extinguishes. For  $1 < U < U_{cr}$ , the solution converges to a stationary profile  $\tilde{T}(x)$ . The shape of the stationary profile depends on the compression velocity; the profile is wider for lower  $U$  (see Figure 2, left panel). When  $U < 1$ , the hot spot grows outwards.

The above discussion applies to initially wide profiles,  $h \gg 1$ , or more specifically, to profiles wider than  $\tilde{T}(x)$ . Narrower profiles converge to a narrower stationary solution. We performed a test where we kept the same  $U = 1.3$  and  $k = 0.75$  and varied  $h$ . For  $h > 2.8$ , all solutions converge to  $\tilde{T}(x)$ . For  $1.66 > h > 2.7$ , the solutions converge to different profiles (see Figure 2, right panel). If  $h < 1.65$ , the solutions become extinct; recall that the Kanel size for this steepness at  $U = 0$  is  $h = 1.50$ .

### 2.1.3 Non-stationary stretching

If compression facilitates quenching, stretching facilitates burning: it increases the area where  $T$  is above the reaction threshold. We consider the no-gravity case,  $\rho = 0$ , with the initial velocity profile  $u(x, 0) = -u^*(x)$ , where  $u^*(x)$  is given by (2.5). We found that the critical size of the hot spot decreases with stretching velocity  $U$  (see table below). Note that the stretching solution evolves even in the absence of gravity.




Figure 2: Stationary profiles of temperature in prescribed, compressing velocity. On the left panel, the profiles are shown for compression  $U = 1.1, 1.2, 1.3,$  and  $1.39$ ; initial width of hot spot  $h \gg 1$ . On the right panel, the profiles are shown for different initial widths,  $h > 2.8$  (solid) and  $h = 2.5, 2.5, 2.0,$  and  $1.66$  (dashed), for  $U = 1.3$ .




Figure 3: Left: the profiles of temperature and velocity at time  $t = 2$  for stretching  $U = 0.2$  and  $U = 0.3$ ; the initial profile has the width  $h = 1.00$  with  $k = 1.50$ . Right: the profiles of  $T$  and velocity at time  $t = 1$  for stretching  $U = 4.2$  and  $U = 4.3$ ; initial profile has width  $h = 0.80$  with  $k = 1.50$ . The initial profiles are shown with dashed lines.




Figure 4: Profiles of velocity and concentration at times  $t = 32, 64, 96, 128$  for  $g = 1/4$  and  $\theta = 0.5$ . Profiles of velocity (solid lines) are compared to velocity as suggested by Eq. (2.11) with  $c = -0.75$  (dashed lines).

$U$	$h_{cr}$ ( $k = 0.75$ )	$h_{cr}$ ( $k = 1.50$ )
0	1.50	1.00
1	1.48	0.94
2	1.48	0.87
4	1.47	0.80

Finally, we point out that the above statement, “compression facilitates quenching while rarefaction (stretching) facilitates burning”, sounds counter-intuitive from the point of view of gas thermodynamics. We remind that in our model  $T$  has a physical meaning of fraction of “hot particles” rather than temperature, and the reaction rate is the function of  $T$  only. Compression and stretching affect the density of both hot and cold particles but preserve their fractions. For instance, squeezing the region with  $T = 0.5$  does not result in the increase of the temperature and the reaction rate. A better model would include a reaction rate that depends on the density of the hot particles rather than their fraction. However implementing such model involves introducing the concepts of density and pressure and an equation of state.






Figure 5: Profiles of velocity and concentration at times  $t = 8, 16, 24, 32$  for  $g = 4$  and  $\theta = 0.5$ . The dashed lines on the top panel show evolution  $u_f(x_f)$  as suggested by Eq. (2.7) and velocity given by Eq. (2.10) at the times matching simulation data. On the bottom panel, dashed line shows  $\phi_0 = 0.195$ .

## 2.2 Quenching by the gravity force

Here we study the growth of an initially small hot spot with constant non-zero gravity in an initially quiescent fluid,  $u(x, 0) = 0$ . In the tested range,  $1 \leq \rho \leq 8$ , the gravity has no influence on the critical size of the initial hot spot. All solutions with  $h$  higher than the Kanel's length grow, at least initially. For hot spots with initial size above Kanel's length, the effect of the initial size is noticeable only at times  $t \sim 1$ . For  $t \gg 1$ , the difference in the initial size shows only as an offset in initial time. The growth of hot spots at later times depends only on the gravity  $\rho$ . And, depending on  $\rho$ , we observe three kinds of solutions.

When the gravity is small the left boundary of the hot spot moves to the left with a constant speed (see Figure 4). The solution at large negative  $x$  resembles a traveling wave. The right boundary is extremely sharp and moves to the right accelerating. We will from now on refer to the right boundary of the hot region as "the shock" – because that is what it is when the viscous term  $u_{xx}$  is not present. As both boundaries move in opposite directions the hot spots at small gravities never quench.

As the gravity is increased, the solution becomes more complicated (Figure 5). The right boundary is still sharp, in the form of a shock, while the left boundary is stretched in the form of long tail of partially burned fluid with temperature below the ignition threshold. This part of the solution will be referred to as "the ramp" or the "wave fan". Its analogue in the inviscid case is a rarefaction wave.

In the region between the ramp and the shock (or, for lower gravities, between two opposite moving fronts) the temperature mostly exceeds the ignition threshold. This is the only region where the reaction occurs; we refer to it as the "combustion wave", or simply "the wave".

Even when both boundaries of the combustion wave move to the right, their dynamics are different and depend on the gravity. For moderate gravities, the shock moves faster than the right border of the ramp; such combustion waves do not quench. For high gravities, the ramp eventually catches up with the shock and the hot spot quenches. This kind of quenching can occur at times significantly exceeding the laminar front self-crossing time, and after the hot region reaches the sizes significantly exceeding the Kanel's length. The object of the following two subsections is to get some intuition on how it happens as well as to obtain formally some orders of magnitude.

### 2.2.1 Ramp-wave-shock structure of the velocity profile

The solution shown in Figure 5 consists of three parts: a cold stationary fluid ahead of the shock, the combustion wave with  $T$  above the ignition temperature, and the ramp where  $T < \theta$ . Initially the hot spot is located at  $x = 0$ . We denote by  $x_f$  the location of the shock and by  $x_b$  the location of the transition point between the ramp and the combustion wave. (The subscript "f" refers to the "front", and the subscript "b" refers to the "back" of combustion wave.) Similarly, we denote the local velocity at corresponding points as  $u_f \equiv u(x_f)$  and  $u_b \equiv u(x_b)$ , and the phase velocities as  $v_f \equiv \dot{x}_f$  and  $v_b \equiv \dot{x}_b$ .

Below we construct an approximate solution at the ramp, the wave, and the shock. Combining them together we find the speed of the shock,  $v_f$ , and the growth rate of the ramp,  $v_b$ . Comparing  $v_b$  and  $v_f$  in the next subsection, we estimate criterion for quenching.

**The ramp.** In Figure 5, both  $T(x)$  and  $u(x)$  appear to be linear in the ramp. In comparison with advection, dissipation effects are negligible on the scale of the ramp. Indeed, if  $L$  is the length of the ramp and  $U$  is the typical velocity in the ramp, then  $T_{xx} \sim 1/L^2 \ll uT_x \sim U/L$  and  $u_{xx} \sim U/L^2 \ll uu_x \sim U^2/L$  for large  $L$  and  $U$ . Neglecting dissipation and taking into account that




Figure 6: Rescaled profiles of temperature and velocity from Figure 5 zoomed on the ramp region. The dashed lines correspond to Eq. (2.6).

$f(T) = 0$  in the ramp, we may find approximate solutions to (2.1) as

$$T(t, x) = \frac{2x}{\rho t^2}, \quad u(t, x) = \frac{2x}{t}, \quad 0 < x < x_b. \quad (2.6)$$

In the ramp region, equations (2.6) agree with numerical simulations: see Figure 6.

We return to Figure 5. The transition between the ramp and the wave occurs at the same value of  $T = \phi_0 < \theta$  which does not depend on time. Assuming that we know  $\phi_0$  we can estimate the location of the transition to the ramp  $x_b$ , and corresponding velocities,

$$x_b = \frac{1}{2}\rho\phi_0 t^2, \quad u_b = \rho\phi_0 t, \quad v_b = \rho\phi_0 t. \quad (2.7)$$

In this case,  $u_b = v_b$ , but in general it does not have to be this way. This aspect will be investigated further in Section 5.

**The wave.** Consider now the velocity in the wave part of the solution,  $x_b < x < x_f$ . In Figure 5, the solutions at different times appear to have the same form  $u_2(x)$ , only shifted in time. We can assume that the origin is located at  $x_b$ . If we substitute velocity in the form  $u(t, x) = u_2(x - x_b) + u_b$ , where  $x_b$  and  $u_b$  are some functions of time, into (2.1), with  $T = 1 -$  we of course do not have  $T = 1$  everywhere, but this will at least give us some order of magnitude, we obtain

$$[(u_b - x'_b)u'_2 + u'_b] + u_2 u'_2 = u''_2 + \rho. \quad (2.8)$$

We see from here that  $u_2$  is time-independent only if  $u_b$  is linear in time and  $x'_b = u_b$ . And luckily  $x_b$  and  $u_b$  given by (2.7) satisfy this condition. The other possible combination is

$$x_b = ct, \quad u_b = c, \quad v_b = c, \quad (2.9)$$

which corresponds to a solution shifting with some constant velocity  $c$ .

When  $x_b$  and  $u_b$  are given by (2.7), the expression in square brackets in (2.8) is equal to  $\rho\phi_0$  and (2.8) can be solved. Neglecting the dissipation terms (the same dimensional argument as for the ramp can be applied here), we obtain  $u_2(x) = \sqrt{2\rho(1-\phi_0)x}$ . The velocity profile in the wave is thus, approximately,

$$u(t, x) = \rho\phi_0 t + \sqrt{2\rho(1-\phi_0)\left(x - \frac{1}{2}\rho\phi_0 t^2\right)}, \quad x_b < x < x_f. \quad (2.10)$$

Similarly, when  $x_b$  and  $u_b$  are given by (2.9), the velocity in the wave is

$$u(t, x) = c + \sqrt{2\rho(x-ct)}, \quad x_b < x < x_f. \quad (2.11)$$

In the numerical simulations we have examples of both types of solutions. For lower gravities, the left boundary of the combustion wave shifts to the left with constant speed; shown in Figure 4 numerical solution agrees with (2.11). For higher gravities, the left side of the combustion wave shifts to the right accelerating; in Figure 5 the numerical solution is compared to (2.10). In both cases, the numerical data is fitted with one unknown parameter — the shift velocity  $c$  in the first case, and temperature at the transition to the ramp  $\phi_0$  in the second case. Both parameters  $c$  and  $\phi_0$  are functions of gravity as seen in Figure 7.

Figure 7 also shows numerical evidence of the existence of a first critical parameter,  $\rho_{cr1}$ , such that:

- for  $\rho < \rho_{cr1}$ , the solution is of the constant shift type, with  $c < 0$ . When gravity approaches zero the left boundary of the combustion wave moves to the left with laminar speed,  $|c| = 1$ . For very small gravities  $\rho \ll 1$ , the speed is  $c = \rho - 1$ .
- For gravities  $\rho > \rho_{cr1}$  the solution is of the accelerated shift type, with  $\phi_0 > 0$ . At  $\rho = \rho_{cr1}$  both parameters  $c$  and  $\phi_0$  are equal to zero, and the solution in the wave is stationary,  $u(t, x) = \sqrt{2\rho x}$ ,  $T(t, x) = 1$ .

### The shock.

The front ahead of the combustion wave is driven by a Burgers shock, the mechanism for which is much stronger than the front propagation due to reaction. Moreover, at high shock speeds, the shock is extremely narrow; the reaction region is narrow as well, and the role of the reaction is reduced. In the vicinity of the shock we can neglect the reaction term in the temperature equation; then the solution is the classical Burgers shock of strength  $u_f$  propagating with speed  $v_f = u_f/2$ .

On the scale of the problem, the shock can be considered as a discontinuity located at  $x_f$  and moving with the speed  $v_f = \dot{x}_f$ . Then, according to (2.10), the location of the shock is given by the following differential equation:

$$\frac{dx_f}{dt} = \frac{1}{2}u(x_f, t) = \frac{1}{2} \left[ \rho\phi_0 t + \sqrt{2\rho(1-\phi_0)\left(x_f - \frac{1}{2}\rho\phi_0 t^2\right)} \right]. \quad (2.12)$$




Figure 7: Parameters  $\phi_0$  and  $c$  as functions of  $\rho$ . Dashed lines correspond to the fits of the form  $\phi_0(\rho) = a + b/(c - \rho)$ . Symbols in the insert diagram represent numerical solutions in different regimes while the lines are schematic borders between regimes.




Figure 8: Velocity at the shock as function of time for different gravities. Dashed lines show velocity at the shock  $u_f = \frac{1}{2}\rho t$  for  $\rho = 1/4$  and  $\rho = 1/2$ , velocity  $u_f = \frac{1}{2}\rho t(1 - \phi_0)$  for  $\rho = 1$  and  $\rho = 2$ , and velocity given by solving (2.12) for  $\rho = 4$  and  $\rho = 8$ .

We compare the expected  $u_f$  to the results on numerical simulations in Figure 8. The agreement between the analytical approximation for  $u_f$  and the numerical simulations, while not perfect, shows the correct qualitative behavior of the analytical prediction. Notice that the decrease of velocity (slowing down of the shock) and abrupt termination of the curves (quenching) are observed only for high gravities. Thus, the numerical simulations suggest the existence of a critical value  $\rho_{cr2}$  such that, for  $\rho < \rho_{cr2}$ , the shock solution exists for all time. We also may infer that

- For gravities  $\rho > \rho_{cr2}$  the temperature drops below the ignition threshold  $\theta$  everywhere in space in a finite time. After this reaction ceases and the solution is quenched.

Notice that in construction of our approximate solution we do not rely on the functional form of reaction rate  $f(\phi)$ . This is not surprising. The reaction rate is non-zero at only two narrow regions in the vicinity of  $x_f$  and  $x_b$ . As we discussed earlier, the reaction is negligible in the shock region because of the compression. The only region where the reaction is important is the transition between the ramp and the wave, the point whose behavior is controlled by  $\phi_0$  and  $c$ . Although the reaction rate does not appear in the discussion, it is implicitly present in the model in the form of empirical parameters  $\phi_0$  and  $c$ .

### 2.2.2 Burning or quenching?

In order to identify the regimes of burning and quenching we compare the speed of the transition point between the ramp and the wave,  $v_b$ , and the shock speed,  $v_f$ , in the limit  $t \rightarrow \infty$ . When  $\rho < \rho_{cr1}$ , the end of the ramp moves with the speed  $v_b = c < 0$ , and the velocity at the shock  $u_f$  can be approximated as  $u_f \approx \frac{1}{2}\rho t$ . The shock speed,  $v_f = \frac{1}{2}u_f$ , is positive. Two sides of the combustion wave move in the opposite directions and quenching never happens.

When  $\rho > \rho_{cr1}$ , the ramp extends with the speed  $v_b = \rho t \phi_0$ . The distance between the end of the ramp and the shock is then  $x_f - x_b := y_f$ , and an equation for  $y_f$  is, from (2.12):

$$\dot{y}_f = \frac{1}{2} \left( \sqrt{2\rho(1 - \phi_0)y_f - \rho\phi_0 t} \right), \quad (2.13)$$

where  $\phi_0$  is the value of the temperature at the transition point. Note first that equation (2.13) has no global in time solution if  $\phi_0 > 1/9$ . Here is the reason: if we write  $y_f = \rho t^2 z$ , then we have

$$2z + t\dot{z} = \frac{1}{2} \left[ \sqrt{2(1 - \phi_0)z - \phi_0} \right].$$

Changing the time variable  $\tau = \ln t$  (for  $t \geq 1$ ) this becomes

$$\begin{aligned} \frac{dz}{d\tau} &= \frac{1}{2} q(\sqrt{z}), \\ q(s) &= \sqrt{2(1 - \phi_0)s - \phi_0} - 4s^2. \end{aligned} \tag{2.14}$$

An elementary study of  $q(s)$  on for  $s > 0$  reveals that

- For  $\phi_0 < 1/9$ , there is  $z_{\phi_0} > 0$  such that  $q > 0$  on  $[0, z_{\phi_0})$  and  $q < 0$  on  $(z_{\phi_0}, +\infty)$ ;
- for  $\phi_0 > 1/9$  we have  $q(s) < 0$  for  $s > 0$ .

In order for (2.14) to have a global solution we need  $z$  to be non-negative and  $q(s)$  not to be uniformly negative for all  $s \geq 0$ . This is true only as long as  $2(1 - \phi_0) - 16\phi_0 > 0$ , or  $\phi_0 < 1/9$ . Therefore, (2.14) has no global in time solutions and quenching occurs if  $\phi_0(\rho) > 1/9$ . The transition temperature  $\phi_0$  increases with the gravity  $\rho$ , and for large  $\rho$  exceeds the critical value  $1/9$ . More precisely, as we show in Theorem 4.1 below,  $\phi_0$  approaches the ignition threshold  $\theta$  as  $\rho$  tends to infinity. Therefore, quenching happens for a sufficiently large  $\rho$  provided that  $\theta > 1/9$ .

**Dependence on the ignition threshold  $\theta$ .** To illustrate the above quenching/propagation analysis, we show in Figure 7 the dependence of  $\phi_0(\rho)$  and  $c(\rho)$  for three values of the ignition threshold  $\theta$  in the reaction rate (2.2).

As  $\theta$  decreases, a stronger force (larger  $\rho$ ) is needed to reach the quenching value  $\phi_0 = 1/9$ . Since  $\phi_0 < \theta$ , reactions with  $\theta < 1/9$  cannot be quenched by any force. The dashed lines in Figure 7 are functions of the form  $\phi_0(\rho) = a + b/(c - \rho)$  fitted to the data. For the case of  $\theta = 1/10$  the fit is bounded by asymptote  $a = 0.0677 < 1/9$ , which suggests that the quenching is impossible for this threshold value.

The value  $\rho_{cr1}$  with  $\phi_0 = 0$  corresponds to the transition to a stationary left front; further reduction of  $\rho$  results in the opposite propagation of fronts, characterized by the speed of the left-going front  $c(\rho)$  rather than by the value of  $\phi_0$ . The speed  $c < 0$  decreases with  $\rho$  down to  $c = -1$  at  $\rho = 0$ , which corresponds to the speed of the undisturbed reaction-diffusion shock. It is interesting that the stationary left shock, when both  $c(\rho) = 0$  and  $\phi_0 = 0$ , is observed for a range of  $\rho$ , rather than for a single value  $\rho_{cr1}$ .

To summarize, with the increase of forcing (see insert diagram in Figure 7) the system with a particular reaction rate can exhibit the following regimes: (i) two shocks moving in opposite directions,  $u_b < 0 < u_f$ , no quenching; (ii) stationary left shock and right-propagating right shock,  $u_b = 0 < u_f$ , no quenching; (iii) both shocks move to right, first shock faster,  $0 < u_b < u_f$ , no quenching; (iv) both shocks move to right, second shock faster,  $0 < u_f < u_b$ , quenching.

In the rest of the paper we construct the building blocks of the observed numerical solutions – the ramp (the wave fan), the combustion wave and the shock – and show that they may be used to build an asymptotic solution. Moreover, we show that for a sufficiently large gravity, a solution that has reached the wave fan-combustion wave-shock structure will ultimately quench.

### 3 Wave fans

In this section we search for non-reactive solutions of (1.5), that is, solutions of

$$\begin{aligned} T_t - T_{xx} + uT_x &= 0 \\ u_t + uu_x &= \rho T. \end{aligned} \quad (3.1)$$

If we additionally suppress the temperature diffusion, the sought-after solution would be an analogue of a rarefaction wave. The correct self-similarity scaling is

$$T(t, x) = \frac{1}{\rho t^{3/2}} \phi\left(\frac{x}{\sqrt{t}}\right), \quad u(t, x) = \frac{1}{t^{1/2}} \psi\left(\frac{x}{\sqrt{t}}\right). \quad (3.2)$$

Setting the self-similar variable  $\eta = x/\sqrt{t}$ , we obtain the following system satisfied by  $\phi(\eta)$  and  $\psi(\eta)$ :

$$\begin{aligned} -\phi'' + \left(\psi - \frac{\eta}{2}\right)\phi' - \frac{3\phi}{2} &= 0 \\ \left(\psi - \frac{\eta}{2}\right)\psi' - \frac{\psi}{2} &= \phi. \end{aligned} \quad (3.3)$$

We require  $\phi$  to be zero at  $(-\infty)$  – remember that, without the heat diffusion, the rarefaction wave is equal to zero for large negative  $\eta$ . In addition, as we will need to match it with a traveling wave at large positive  $\eta$ , the function  $\phi$  should have a bounded derivative on the whole domain. Also, we want  $\phi$  and  $\psi$  to be positive. We will see that positivity plus a strong decay at infinity implies integrability plus a global Lipschitz bound.

The equation for the function  $\psi$  may be rewritten as

$$\left(\frac{\psi^2 - \eta\psi}{2}\right)' = \phi,$$

which implies the following quadratic equation for  $\psi$ :

$$\psi^2 - \eta\psi = 2 \int_{-\infty}^{\eta} \phi(\eta') d\eta'. \quad (3.4)$$

Choosing the positive root in the above equation we get

$$\psi(\eta) = \frac{\eta}{2} + \frac{1}{2} \sqrt{\eta^2 + 8 \int_{-\infty}^{\eta} \phi(\eta') d\eta'}. \quad (3.5)$$

Using this expression in the first equation in (3.3) leads to the following problem for the function  $\phi(\eta)$  which we are now going to investigate:

$$\begin{aligned} -\phi'' + \frac{1}{2} \left[ \eta^2 + 8 \int_{-\infty}^{\eta} \phi(\xi) d\xi \right]^{\frac{1}{2}} \phi' - \frac{3\phi}{2} &= 0 \\ \phi > 0, \quad \phi \in L^1(\mathbb{R}_-), \quad \phi' \in L^\infty(\mathbb{R}). \end{aligned} \quad (3.6)$$

First, a definition: we say that  $\phi$  has a Gaussian decay at  $-\infty$  with exponent  $\lambda$  if  $\eta \mapsto e^{\lambda\eta^2} \phi(\eta)$  has at most polynomial growth as  $\eta \rightarrow -\infty$ . We have the following result for (3.6).

**Theorem 3.1** *Equation (3.6) has at least one positive solution  $\phi(\eta)$ , having Gaussian decay at  $-\infty$  with any exponent  $\lambda \in (0, 1/4)$ . For any such solution, there exists a number  $b \in \mathbb{R}$  such that we have in addition, as  $\eta \rightarrow +\infty$ :*

$$\phi(\eta) = 2\eta + b(1 + o(1))\eta^{1/3}, \quad \phi'(\eta) = 2 + O(\eta^{-2/3}). \quad (3.7)$$



We note here that the Gaussian decay has to be imposed. At this stage we do not know whether there are waves that have no Gaussian decay. On the other hand, there is no need to impose any bound on the growth of  $\phi$  at  $+\infty$ : the Gaussian decay plus positivity implies a global Lipschitz bound, with a linear growth that will allow matching with another elementary solution.

Let us now explain why imposing a Gaussian decay is relevant. Equation (3.6), linearized around the rest state  $\phi_0 = 0$  at  $\eta = -\infty$  becomes

$$-\phi'' - \frac{\eta}{2}\phi' - \frac{3\phi}{2} = 0 \quad (3.8)$$

This equation has two integrable solutions:

$$h(\eta) = \left(\frac{\eta^2}{4} - \frac{1}{2}\right) e^{-\eta^2/4} = \frac{d^2}{d\eta^2} \left(e^{-\eta^2/4}\right), \quad k(\eta) = -\frac{1}{5} \frac{d^2}{d\eta^2} \left(e^{-\eta^2/4} \int_0^\eta e^{\zeta^2/4} d\zeta\right) \sim -\eta^{-3}. \quad (3.9)$$

A solution  $\phi$  of (3.6), having no Gaussian decay, would satisfy

$$\phi(\eta) \sim Ak(\eta) \quad \text{as } \eta \rightarrow -\infty. \quad (3.10)$$

Using (3.4) with the fact that  $\psi \geq 0$  in mind, we see that then

$$\psi(\eta) \sim \frac{\eta \pm \sqrt{\eta^2 + A/2\eta^2}}{2} \sim \frac{A}{8|\eta|^3} \quad \text{as } \eta \rightarrow -\infty.$$

But then, let us come back to the functions  $T(t, x)$  and  $u(t, x)$  defined by (3.2) – we have, by (3.10):

$$T(t, x) \sim \frac{1}{\rho|x|^3}, \quad u(t, x) \sim \frac{t}{\rho|x|^3}. \quad (3.11)$$

However, this would mean that for large  $t$ , the flow grows in any compact region in  $x$ , which contradicts the numerics in Section 2. On the other hand, solutions which have a Gaussian decay of  $\phi$  as  $\eta \rightarrow -\infty$  and grow linearly in  $\eta$  as  $\eta \rightarrow +\infty$  do not have this problem. It is interesting to note that the Gaussian decay is impossible without the presence of diffusion in the temperature equation.

The proof of Theorem 3.1 is via a shooting argument: we first construct a solution  $\phi_\delta$  to (3.3) on a half-line of the form  $(-\infty, -\eta_1)$  for some large  $\eta_1$ , having a given value  $\delta > 0$  at the end-point:  $\phi_\delta(-\eta_1) = \delta$ . The real number  $\delta$  is then adjusted to get  $\phi_\delta(\eta) = O(\eta)$  as  $\eta \rightarrow +\infty$ .

**Step 1. Solution on a left half line.** Take any  $\eta_1 \geq \sqrt{6}$ . We claim that, given  $\delta > 0$ , the Dirichlet problem

$$-\phi_\delta'' + \frac{1}{2} \left[ \eta^2 + 8 \int_{-\infty}^\eta \phi_\delta(\xi) d\xi \right]^{\frac{1}{2}} \phi_\delta' - \frac{3\phi_\delta}{2} = 0 \quad \text{on } (-\infty, -\eta_1), \quad \phi_\delta(-\eta_1) = \delta \quad (3.12)$$

with an additional constraint that  $\phi_\delta$  has a Gaussian decay as  $\eta \rightarrow -\infty$  has a positive solution. Indeed, let  $h(\eta)$  be defined by (3.9) and define the operator

$$L_0 = -\frac{d^2}{d\eta^2} - \frac{\eta}{2} \frac{d}{d\eta} - \frac{3}{2}.$$

We have  $L_0 h = 0$ . Moreover, on the half-line  $\mathbb{R}_-$ , the function  $h$  is positive and increasing on  $(-\infty, -\sqrt{6})$ , decreasing on  $(-\sqrt{6}, 0)$  and negative on  $(-\sqrt{2}, 0)$ . Now, take  $r > 0$  and  $\eta_1 \geq \sqrt{6}$ . Let  $\phi_r$  be the solution of the boundary value problem

$$\begin{aligned} -\phi_r'' + \frac{1}{2} \left[ \eta^2 + 8 \int_{-r}^\eta |\phi_r(\xi)| d\xi \right]^{\frac{1}{2}} \phi_r' - \frac{3\phi_r}{2} &= 0 \quad \text{on } (-r, -\eta_1) \\ \phi_r(-r) &= \frac{\delta h(-r)}{h(-\eta_1)}, \quad \phi_r(-\eta_1) = \delta \end{aligned} \quad (3.13)$$

Let us show that (3.13) has a nonnegative solution that satisfies

$$0 \leq \phi_r(\eta) \leq \bar{h}(\eta) = \frac{\delta h(\eta)}{h(-\eta_1)}. \quad (3.14)$$

Given a function  $q(\eta) \in C([-r, -\eta_1])$  define the nonlinear mapping  $\phi = \mathcal{M}q$ , where  $\phi$  is the unique solution of the linear boundary value problem

$$\begin{aligned} -\phi'' + \frac{1}{2} \left[ \eta^2 + 8 \int_{-r}^{\eta} |q(\xi)| d\xi \right]^{\frac{1}{2}} \phi' - \frac{3\phi}{2} &= 0 \text{ on } (-r, -\eta_1), \\ \phi(-r) &= \frac{\delta h(-r)}{h(-\eta_1)}, \quad \phi(-\eta_1) = \delta. \end{aligned} \quad (3.15)$$

In order to see that (3.15) indeed has a unique solution we write  $\phi = \bar{h} + w$  and observe that existence of a strictly positive solution of  $L_0 h = 0$  implies that the operator  $L_0$ , defined on the space  $W_r = \{w \in C((-r, -\eta_1)) : w(-r) = w(-\eta_1) = 0\}$  with the standard domain  $D(L_0)$  is invertible with a compact inverse. Upon defining  $\zeta(\eta) = e^{\eta^2/8} \phi(\eta)$  and

$$a(\eta, q) = \frac{1}{2} \left[ \eta^2 + 8 \int_{-a}^{\eta} |q(\xi)| d\xi \right]^{\frac{1}{2}}$$

we obtain :

$$-\zeta'' + \left( a(\eta, q) - \frac{\eta}{2} \right) \zeta' + \left( \frac{\eta^2}{16} + \frac{|\eta|}{4} a(\eta, q) - \frac{7}{4} \right) \zeta = 0. \quad (3.16)$$

Note that, for  $|\eta| \geq 6$  we have

$$\frac{\eta^2}{16} + \frac{|\eta|}{4} a(\eta, q) - \frac{7}{4} \geq \frac{\eta^2}{8}. \quad (3.17)$$

Then the maximum principle implies that  $\zeta$  can not attain a negative minimum inside the interval  $(-r, -\eta_1)$  and hence  $\zeta \geq 0$ , which, in turn, implies that  $\phi \geq 0$ .

Let us now show that in addition  $\phi$  satisfies  $\phi \leq \bar{h}(\eta)$ . The function  $\bar{h}(\eta)$  is monotonically increasing and is thus a super-solution to (3.15) in the sense that:

$$-\bar{h}'' + \frac{1}{2} \left[ \eta^2 + 8 \int_{-r}^{\eta} |q(\xi)| d\xi \right]^{\frac{1}{2}} \bar{h}' - \frac{3\bar{h}}{2} \geq 0 \text{ on } (-r, -\eta_1), \quad \bar{h}(-r) = \frac{\delta h(-r)}{h(-\eta_1)}, \quad \bar{h}(-\eta_1) = \delta. \quad (3.18)$$

Given any  $M \geq 1$  the difference  $w_M = M\bar{h} - \phi$  satisfies

$$\begin{aligned} -w_M'' + \frac{1}{2} \left[ \eta^2 + 8 \int_{-r}^{\eta} |q(\xi)| d\xi \right]^{\frac{1}{2}} w_M' - \frac{3w_M}{2} &\geq 0 \text{ on } (-r, -\eta_1), \\ w_M(-r) &\geq 0, \quad w_M(-\eta_1) \geq 0. \end{aligned} \quad (3.19)$$

Another consequence of the maximum principle is that  $w_M$  cannot attain an interior minimum in  $(-r, -\eta_1)$  at a point where  $w_M = 0$ . This, combined with the fact that  $w_M > 0$  for a sufficiently large  $M$ , and decreasing  $M$  until we do not have  $w_M > 0$ , yields that

$$\bar{M} := \inf\{M > 0 : w_M(\eta) \geq 0 \text{ for all } \eta \in (-r, -\eta_1)\} = 1.$$

Therefore, we have  $0 \leq \phi(\eta) \leq \bar{h}(\eta)$  for all functions  $q(x) \in C([-r, -\eta_1])$ . As a consequence, the nonlinear operator  $\mathcal{M}$  sends the closed set  $E = \{\phi \in C([-r, -\eta_1]) : 0 \leq \phi(\eta) \leq \bar{h}(\eta)\}$  to itself. The elliptic regularity theory implies that the mapping  $\mathcal{M}$  is compact. The Schauder fixed point

theorem implies that it has a fixed point in  $E$  which is a solution of (3.13). In addition, the limit has to satisfy (3.14).

An unpleasant fact in the construction of  $\phi_\delta$  is that – as is usual with applications of the Schauder theorem – it yields no information on the uniqueness of the solution  $\phi_\delta$ , or on the continuity of  $\phi_\delta$  with respect to  $\delta$ . This inconvenient will be fixed later in the course of the proof of the proposition, by adjusting the shooting point  $\eta_1$ .

**Step 2. Some estimates for  $\phi_\delta$ .** Still assume that  $\eta_1 \geq 6$  is fixed. Let us consider the functions  $h(\eta)$  and  $k(\eta)$  defined by (3.9) and introduce the following quantities:

$$u(\eta) = \int_{-\infty}^{\eta} \phi_\delta(\xi) d\xi, \quad a(\eta) = \frac{1}{2} \sqrt{\eta^2 + 8u(\eta)}, \quad (3.20)$$

The starting point of this step is the following

**Lemma 3.2** *For every  $\eta_1 \geq 6$ , any solution  $\phi$  of (3.12) with a Gaussian decay satisfies*

$$\phi(\eta) \leq \delta \frac{h(\eta)}{h(-\eta_1)}. \quad (3.21)$$

**Proof.** Upon defining  $\zeta(\eta) = e^{\eta^2/8} \phi(\eta)$ , as in Step 1, we obtain – recall that  $a(\eta)$  is defined by (3.20):

$$\tilde{L}\zeta := -\zeta'' + \left(a(\eta) - \frac{\eta}{2}\right) \zeta' + \left(\frac{\eta^2}{16} + \frac{|\eta|}{4} a(\eta) - \frac{7}{4}\right) \zeta = 0. \quad (3.22)$$

Note again that, for  $|\eta| \geq 6$ , we have

$$\frac{\eta^2}{16} + \frac{|\eta|}{4} a(\eta) - \frac{7}{4} \geq \frac{\eta^2}{8}. \quad (3.23)$$

For every  $\varepsilon > 0$ , the function

$$\bar{\zeta}_\varepsilon(\eta) = \left( \delta \frac{h(\eta)}{h(-\eta_1)} + \varepsilon k(\eta) \right) e^{\eta^2/8}$$

satisfies  $\tilde{L}\bar{\zeta}_\varepsilon \geq 0$ . Moreover, we have, because of the Gaussian decay of  $\phi$ :  $\bar{\zeta}_\varepsilon(\eta) - \zeta(\eta) > 0$  for a sufficiently large negative  $\eta$ . Therefore, if it gets negative inside  $(-\infty, -\eta_1)$  it has to reach a minimum – a situation precluded by (3.23). Hence, we have  $\bar{\zeta}_\varepsilon(\eta) - \zeta(\eta) > 0$  for all  $\eta < -\eta_1$  and all  $\varepsilon > 0$  – thus, (3.21) holds.  $\square$

Let us now examine what happens to  $\phi_\delta$  as  $\delta$  becomes large. It follows from the upper bound in (3.14) that

$$\int_{-\infty}^{\eta} \phi_\delta \leq \sqrt{6}\delta/2 < 2\delta$$

and  $\phi'_\delta(-\eta_1) > 0$ . Therefore, as  $\phi_\delta > 0$  and

$$\phi'_\delta(\eta) = \phi'(-\eta_1) \exp(A(-\eta_1) - A(\eta)) + \frac{3}{2} \int_{\eta}^{-\eta_1} \exp(A(\xi) - A(\eta)) \phi_\delta(\xi) d\xi \geq 0,$$

where

$$A'(\eta) = -\frac{1}{2} \left[ \eta^2 + 8 \int_{-\infty}^{\eta} \phi(\xi) d\xi \right]^{\frac{1}{2}},$$

the function  $\phi_\delta$  is increasing. Hence, we have

$$-\phi_\delta'' + \frac{1}{2}(4\sqrt{\delta} - \eta)\phi_\delta' - \frac{3}{2}\phi_\delta \geq -\phi_\delta'' + \frac{1}{2}\sqrt{\eta^2 + 16\delta}\phi_\delta' - \frac{3}{2}\phi_\delta \geq 0.$$

Then, just as in Lemma 3.2, we may get a lower bound  $\phi_\delta \geq \underline{\phi}_\delta$ , where

$$\begin{aligned} -\underline{\phi}_\delta'' + \frac{1}{2}(4\sqrt{\delta} - \eta)\underline{\phi}_\delta' - \frac{3\underline{\phi}_\delta}{2} &= 0 \\ \underline{\phi}_\delta(-\eta_1) &= \delta. \end{aligned} \tag{3.24}$$

The function  $\underline{\phi}_\delta$  is given explicitly by

$$\underline{\phi}_\delta(\eta) = \frac{\delta h(\eta - 4\sqrt{\delta})}{h(-\eta_1 - 4\sqrt{\delta})}.$$

Hence, for  $\eta \in [-\eta_1 - 1, -\eta_1]$  we have for a sufficiently large  $\delta > 0$ :

$$\int_{-\infty}^{-\eta} \underline{\phi}_\delta(\xi) d\xi \sim \frac{C\delta(\eta + 4\sqrt{\delta})}{(\eta_1 + 4\sqrt{\delta})^2} \sim C\sqrt{\delta}, \text{ as } \delta \rightarrow +\infty,$$

where  $C > 0$  is independent of  $\delta$ . Using this information, we integrate (3.12) on  $(-\infty, -\eta_1]$ , and obtain

$$\begin{aligned} \phi_\delta'(-\eta_1) &= \frac{1}{2} \int_{-\infty}^{-\eta_1} \left( \eta^2 + 8 \int_{-\infty}^{\eta} \phi_\delta(\xi) d\xi \right)^{\frac{1}{2}} \phi_\delta'(\eta) d\eta - \frac{3}{2} \int_{-\infty}^{-\eta_1} \phi_\delta(\eta) d\eta \\ &\geq C_1\delta^{5/4} - C_2\delta \sim C_1\delta^{5/4} \text{ as } \delta \rightarrow +\infty. \end{aligned}$$

The following estimates are therefore true for large  $\delta$ :

$$C_1\sqrt{\delta} \leq \int_{-\infty}^{-\eta_1} \phi_\delta \leq C_2\delta, \quad \phi_\delta(-\eta_1) = \delta, \quad \phi_\delta'(-\eta_1) \geq C_3\delta^{5/4}. \tag{3.25}$$

**Step 3. Extension of  $\phi_\delta$  and its behavior for  $\eta > -\eta_1$ .** Estimates (3.25) and equation (3.20) enable us to extend  $\phi_\delta$  past  $-\eta_1$ , on a maximal interval  $[-\eta_1, \eta_\infty^\delta)$  – with, possibly,  $\eta_\infty^\delta = +\infty$ . Let us define the sets

$$\begin{aligned} X_-^{\eta_1} &= \{\delta > 0 : \exists \eta_2 > -\eta_1 \text{ such that } \phi_\delta(\eta_2) = 0\} \\ X_+^{\eta_1} &= \{\delta > 0 : \phi_\delta > 0 \text{ and } \limsup_{\eta \rightarrow \eta_\infty^\delta} \phi_\delta'(\eta) = +\infty\}. \end{aligned} \tag{3.26}$$

It is clear that  $\phi_\delta$  also depends on  $\eta_1$ , but this dependence is not going to be indicated by a sub or a subscript, to keep the notations readable.

We begin the analysis for  $\delta \notin X_-^{\eta_1}$  with the following lemma.

**Lemma 3.3** *Assume that  $\delta \notin X_-^{\eta_1}$ , then  $\phi_\delta' > 0$ .*

**Proof.** Let us recall that if  $\phi_\delta \geq 0$  (which is the case for  $\delta \notin X_-^{\eta_1}$ ) then the inequality

$$\phi_\delta'' - a(\eta)\phi_\delta' \leq 0,$$

holds, with the function  $a(\eta)$  defined in equation (3.20) above. It follows that

$$\left( \exp\left(-\int a(\eta) d\eta\right) \phi_\delta' \right)' \leq 0. \tag{3.27}$$

Therefore, for any  $\eta_2$  and  $\eta_3$  larger than  $\eta_1$ , with  $\eta_3 > \eta_2$ , we have

$$\exp\left(-\int_{\eta_2}^{\eta_3} a(\xi)d\xi\right) \phi'_\delta(\eta_3) \leq \phi'_\delta(\eta_2).$$

It follows that if there exists a point  $\eta_2$  where  $\phi'_\delta(\eta_2) < 0$  then  $\phi'_\delta(\eta) \leq \phi'_\delta(\eta_2) < 0$  for all  $\eta \geq \eta_2$ . Hence  $\phi_\delta(\eta)$  has to vanish at some point. This contradicts the assumption that  $\delta \notin X_-^{\eta_1}$  and finishes the proof of Lemma 3.3.  $\square$

The main result of this step is the characterization of  $X_+^{\eta_1}$ .

**Lemma 3.4** *Let  $\delta \in X_+^{\eta_1}$ , then  $\eta_\infty^\delta < +\infty$ .*

**Proof.** Consider  $\delta \in X_+^{\eta_1}$ ; it is convenient to work with the logarithmic derivative of  $\phi_\delta$ :

$$\xi_\delta = \frac{\phi'_\delta}{\phi_\delta}. \quad (3.28)$$

Let us drop the subscript  $\delta$  for the moment. The equation for  $\xi$  is

$$\xi' = a(\eta)\xi - \xi^2 - \frac{3}{2}. \quad (3.29)$$

The term  $(-\xi^2)$  would, in principle, prevent a blow-up; it is the role of the – seemingly linear – term  $a(\eta)\xi$  to force it. Assume, therefore, that  $\eta_\infty = +\infty$ , and let us try to reach a contradiction.

**Case 1.** Assume that there exists a sequence  $(\eta_n)_n$  going to  $+\infty$  such that  $\lim_{n \rightarrow +\infty} \xi(\eta_n) = +\infty$ .

We claim that then  $\xi'(\eta) > 0$  for all sufficiently large  $\eta > 0$ . Indeed, there exists  $\eta_0 > 0$  such that  $\xi'(\eta_0) > 0$ . If  $h := \xi'$  we have, by Lemma 3.3

$$h' + (-a(\eta) + 2\xi)h = \frac{1}{2} \frac{\eta + 4\phi}{a(\eta)} \xi > 0, \quad h(\eta_0) > 0. \quad (3.30)$$

This implies  $h(\eta) > 0$  for  $\eta \geq \eta_0$ .

To prove that we have blow-up, we use an elementary numerical analysis procedure: pick  $\eta_0 > 0$  so that  $\xi' > 0$  on  $(\eta_0, +\infty)$  and  $\phi_\delta(\eta_0) \geq 10$ . The value of  $\xi(\eta_0)$  may be taken arbitrarily large, because  $\xi(\eta_n) \rightarrow +\infty$ . Let  $\lambda_n = 1/(n+1)^2$  and set  $\zeta_n = \eta_0 + \sum_{k \leq n} \lambda_k$ . First, we have

$$\xi(\zeta_{n+1}) - \xi(\zeta_n) = \int_{\zeta_n}^{\zeta_{n+1}} [a(\zeta)\xi(\zeta) - \xi^2(\zeta)] d\zeta - \frac{3\lambda_n}{2}.$$

The function  $a(\zeta)$  for  $\zeta \in (\zeta_n, \zeta_{n+1})$  may be bounded as

$$\begin{aligned} a(\zeta) &= \frac{1}{2} \left( \zeta^2 + 8 \int_{-\infty}^{\zeta} \phi_\delta(\xi)d\xi \right)^{1/2} \geq \frac{1}{2} \left( 8 \int_{\zeta_n}^{\zeta} \phi_\delta(\xi)d\xi \right)^{1/2} \\ &\geq \left( 2 \int_{\zeta_n}^{\zeta} \phi_\delta(\eta_0) \exp\left(\int_{\eta_0}^{\zeta'} \xi(x)dx\right) d\zeta' \right)^{1/2}. \end{aligned}$$

Therefore, using positivity and monotonicity of  $\psi$  we have

$$\begin{aligned}
\xi(\zeta_{n+1}) - \xi(\zeta_n) &\geq \int_{\zeta_n}^{\zeta_{n+1}} \left( \sqrt{2\phi_\delta(\eta_0)} \left[ \int_{\zeta_n}^{\zeta} \exp \left( \int_{\eta_0}^{\zeta'} \xi(x) dx \right) d\zeta' \right]^{1/2} \xi(\zeta) - \xi^2(\zeta) \right) d\zeta - \frac{3\lambda_n}{2} \\
&\geq \int_{\zeta_n}^{\zeta_{n+1}} \sqrt{2\phi_\delta(\eta_0)} \left[ \int_{\zeta_n}^{\zeta} \exp \left( \int_{\zeta_{n-1}}^{\zeta_n} \xi(\zeta_{n-1}) dx \right) d\zeta' \right]^{1/2} \xi(\zeta) d\zeta - \lambda_n \xi^2(\zeta_{n+1}) - \frac{3\lambda_n}{2} \\
&= \sqrt{2\phi_\delta(\eta_0)} \xi(\zeta_n) e^{(\zeta_n - \zeta_{n-1})\xi(\zeta_{n-1})/2} \int_{\zeta_n}^{\zeta_{n+1}} \sqrt{\zeta - \zeta_n} d\zeta - \lambda_n \xi^2(\zeta_{n+1}) - \frac{3\lambda_n}{2} \\
&\geq \lambda_n^{3/2} \xi(\zeta_n) e^{\lambda_n \psi(\zeta_{n-1})/8} - \lambda_n \xi(\zeta_{n+1})^2 - \frac{3\lambda_n}{2}.
\end{aligned}$$

As  $\lambda_n \leq 1/2$  and we may take  $\xi(\eta_0) \geq 10$  (and thus  $\xi(\zeta_n) \geq 10$  for all  $n$ ) it follows that

$$\xi(\zeta_{n+1})^2 \geq C \lambda_n^{3/2} \xi(\zeta_n) e^{\lambda_n \xi(\zeta_{n-1})/8},$$

or

$$\xi(\zeta_{n+1}) \geq \frac{e^{\xi(\zeta_{n-1})/(16(n+1)^2)}}{(n+1)^{3/2}}.$$

Now, choose  $r > 0$  so that  $e^{r(n-1)^4/[16(n+1)^2]}/(n+1)^{3/2} \geq r(n+1)^4$  for all  $n \in \mathbb{N}$ . An easy induction shows that, if  $\xi(\eta_0)$  is large enough we have  $\xi(\zeta_n) \geq rn^4$ . This contradicts the assertion  $\eta_\infty = +\infty$ .

**Case 2.** Assume that  $\xi$  is bounded. Then (3.29) may be integrated from  $+\infty$  to yield

$$\xi(\eta) = \int_\eta^{+\infty} \left( \frac{3}{2} + \psi^2 \right) e^{-\int_\eta^\zeta a(\zeta') d\zeta'} d\zeta,$$

which, as  $\xi \leq C$ , implies  $\xi \leq Ca(\eta)^{-1}$ , where  $C$  does not depend on  $\eta$ . This implies in turn

$$0 \leq \frac{\phi'_\delta}{\phi_\delta} \leq \frac{C}{\sqrt{\int_0^\eta \phi_\delta(\zeta) d\zeta}}, \quad (3.31)$$

which, after integration, yields

$$0 \leq \phi_\delta(\eta) \leq C \left( 1 + \sqrt{\int_0^\eta \phi_\delta(\zeta) d\zeta} \right). \quad (3.32)$$

Integrating (3.32) we obtain  $\int_0^\eta \phi_\delta(\zeta) d\zeta \leq C(1 + \eta^2)$ , which by (3.32) again, translates into the bound  $\phi_\delta(\eta) \leq C(1 + \eta)$ . But now, we may start again from the inequality  $\xi \leq C/a(\eta)$ , use the definition of  $\xi$  and the just obtained information: it follows that

$$\phi'_\delta(\eta) \leq \frac{C\phi}{a(\eta)} \leq \frac{C(1 + \eta)}{\eta} \leq C.$$

This contradicts the fact that  $\delta \in X_+^{\eta_1}$ . We conclude that  $\eta_\infty^\delta < +\infty$  for all  $\delta \in X_+^{\eta_1}$ .  $\square$

One important consequence of Lemma 3.4 is the following

**Corollary 3.5** *There exists  $\delta_0 > 0$  such  $[\delta_0, +\infty) \subset X_+^{\eta_1}$ .*

**Proof.** Let us recall that the logarithmic derivative  $\xi = \phi'_\delta/\phi_\delta$  satisfies equation (3.29). Moreover, if  $\phi_\delta(\eta') \geq 0$  for all  $\eta' < \eta$  then  $a(\eta) \geq a_0 = a(-\eta_1)$ . It follows that under this assumption and if  $\xi(\eta) > 0$  we have

$$\xi' \geq a_0\xi - \xi^2 - \frac{3}{2}. \quad (3.33)$$

In addition, for large enough  $\delta > 0$  we have, by estimate (3.25) of Step 2:

$$\xi(-\eta_1) = \frac{\phi'_\delta(-\eta_1)}{\phi_\delta(-\eta_1)} \geq C\delta^{1/4}.$$

As the smallest root  $q_0$  of the right side of (3.33) is smaller than  $\xi(-\eta_1)$  and  $\phi_\delta$  may not become negative before so does the function  $\xi$ , it follows from the above that  $\xi(\eta) > q_0$  for all  $\eta > -\eta_1$ . As a consequence, we have  $\phi'_\delta > q_0\phi_\delta$  and thus  $\phi_\delta$  blows up at infinity (or at a finite distance) together with its derivative and so  $\delta \in X_+^{\eta_1}$ .  $\square$

**Step 4. Choice of the shooting point.** Take, for definiteness,  $\eta_1 = 7$ . Corollary 3.5 implies the existence of  $\delta_0 > 0$  such that: if  $\phi$  is a solution of (3.6) with Gaussian decay at  $-\infty$ , then  $\phi(-7) \leq \delta_0$ . By Lemma 3.2 we have  $\phi(\eta) \leq \delta_0 h(\eta)/h(-7)$ , a quantity that decays to 0 as  $\eta \rightarrow -\infty$ . Pick any  $\lambda_0 \in (1/8, 1/4)$ , that will remain fixed until the end of the proof of Theorem 3.1. By elementary elliptic regularity we may find a constant  $\eta_0 > 7$  such that: if  $\phi$  is a solution of (3.6) with Gaussian decay at  $-\infty$ , then

$$\forall \eta \leq -\eta_0, \quad 0 \leq \phi(\eta), \quad \phi'(\eta) \leq e^{-\lambda_0\eta^2}. \quad (3.34)$$

For  $\eta_1 > \eta_0$ , let us go back to problem (3.12). We may now prove the uniqueness and continuity with respect to  $\delta$  that were lacking.

**Lemma 3.6** *If  $\eta_1 > 0$  is large enough, and  $\delta \in [0, e^{-\lambda_0\eta_1^2}]$ , the problem (3.12) has exactly one solution, that we still call  $\phi_\delta$ .*

**Proof.** Let  $\phi_1$  and  $\phi_2$  two such solutions; define  $\zeta(\eta) = e^{\eta^2/8}(\phi_1(\eta) - \phi_2(\eta))$ , and

$$a_i(\eta) = \frac{1}{2} \sqrt{\eta^2 + 8 \int_{-\infty}^{\eta} \phi_i}. \quad (3.35)$$

The equation for  $\zeta$  is

$$-\zeta'' + \left(a_1(\eta) - \frac{\eta}{2}\right) \zeta' + \left(\frac{\eta^2}{16} + \frac{|\eta|}{4}a_1(\eta) - \frac{7}{4}\right) \zeta = e^{\eta^2/8}(a_2 - a_1)(\eta)\phi'_2(\eta). \quad (3.36)$$

Note again that, for  $\eta \geq 6$ , we have

$$\frac{\eta^2}{16} + \frac{|\eta|}{4}a_1(\eta) - \frac{7}{4} \geq \frac{\eta^2}{8}. \quad (3.37)$$

By the definition (3.35) of the  $a_i$ 's, we have

$$4e^{\eta^2/8} |a_2(\eta) - a_1(\eta)| = e^{\eta^2/8} \frac{\left| \int_{-\infty}^{\eta} (\phi_2 - \phi_1) dx \right|}{a_1(\eta) + a_2(\eta)} \leq \frac{e^{\eta^2/8} \int_{-\infty}^{\eta} e^{-x^2/8} dx}{a_1(\eta) + a_2(\eta)} \|\zeta\|_\infty \leq 4 \frac{\|\zeta\|_\infty}{\eta_1^3}.$$

Combining the above inequality with estimate (3.34) and inequality (3.37), we obtain from (3.36)

$$\|\zeta\|_\infty \leq \frac{8e^{-\lambda_0\eta_1^2}}{\eta_1^5} \|\zeta\|_\infty.$$

This implies  $\zeta \equiv 0$  as soon as  $\eta_1$  is large enough.  $\square$

**Step 5. Existence of the wave fan.** We have to prove two things: first, the existence of a solution to (3.12) with Gaussian decay; second, the asymptotic behavior of the constructed solution. Let us first worry about the existence: for this we fix any  $\eta_1$  large enough such that

- Lemma 3.6 holds, and
- $e^{-\lambda_0 \eta_1^2} \in X_+^{\eta_1}$ .

The second condition above is realizable because if  $\delta_0 \in X_+^{\eta_1}$  then, as  $\phi_\delta$  satisfies the Gaussian decay bound  $\phi_\delta(\eta) \leq \delta \bar{h}(\eta)/\bar{h}(-\eta_1)$ , then when we increase  $\eta_1$  the "critical"  $\delta_0$  from Corollary 3.5 is approximately multiplied by the factor  $e^{-\lambda_0 \eta_1^2}$ .

We now redefine the sets  $X_\pm^{\eta_1}$  by restricting the values of  $\delta$  to the interval  $[0, e^{-\lambda_0 \eta_1^2}]$ . For small  $\delta > 0$ , the function  $\phi_\delta$  is close on compact intervals to  $\delta h(\eta)/h(-\eta_1)$ . Hence, it vanishes at some point  $\eta < 0$  close to  $(-\sqrt{2})$  – the negative point where  $h(\eta)$  vanishes. This says that the set  $X_-^{\eta_1}$  is nonempty. Moreover, the functions  $\phi_\delta$  may not attain a local minimum equal to zero. Therefore, the continuity of  $\delta \mapsto \phi'_\delta(\eta)$  on compact sets implies that  $X_-^{\eta_1}$  is open. On the other hand, we know that  $X_+^{\eta_1}$  is nonempty. By the arguments in the proof of Case 1 in Lemma 3.4 and the continuity of  $\delta \mapsto \phi_\delta(\eta)$  on compact intervals, it is also open. Consequently, there exists  $\delta \in [0, e^{-\lambda_0 \eta_1^2}] \setminus (X_+^{\eta_1} \cup X_-^{\eta_1})$ . This  $\delta$  generates our desired solution  $\phi(\eta)$  of (3.12).

**Step 6: Behavior of  $\phi$  at  $+\infty$ : the first term in expansion (3.7).** If  $u(\eta)$  and  $a(\eta)$  are defined by the expressions (3.20) above, then the equation for  $u$  is

$$-u''' + a(\eta)u'' - \frac{3}{2}u' = 0,$$

and there exists  $C > 0$  so that

$$u(\eta) \leq C\eta^2 \text{ for } \eta > 0 \quad (3.38)$$

as  $\phi' \in L^\infty(\mathbb{R})$ . This implies by integration from  $\eta$  to  $+\infty$ , with  $\eta > 0$ :

$$\begin{aligned} u''(\eta) &= \frac{3}{2} \int_\eta^{+\infty} \exp\left(-\int_\eta^\xi a(\zeta) d\zeta\right) u'(\xi) d\xi \\ &= \frac{3u'(\eta)}{2a(\eta)} + \frac{3}{2} \int_\eta^{+\infty} \exp\left(-\int_\eta^\xi a(\zeta) d\zeta\right) \left(\frac{u'(\xi)}{a(\xi)}\right)' d\xi := \frac{3u'(\eta)}{2a(\eta)} - f(\eta), \end{aligned} \quad (3.39)$$

the last line being obtained by integration by parts. The uniform bound for  $\phi' = u''$  and positivity of  $u$  imply that  $C_1\eta \leq a(\eta) \leq C_2\eta$ , and  $f(\eta)$  satisfies for  $\eta > 0$ :

$$\begin{aligned} |f(\eta)| &\leq C \int_\eta^{+\infty} \exp\left\{-C \int_\eta^\xi \zeta d\zeta\right\} \frac{d\xi}{\xi} = \int_\eta^{+\infty} \exp\{-C(\xi^2 - \eta^2)\} \frac{d\xi}{\xi} \\ &= Ce^{C\eta^2} \int_\eta^{+\infty} \exp(-C\xi^2) \frac{\xi d\xi}{\eta^2} \leq \frac{C}{\eta^2}, \end{aligned}$$

so that  $f(\eta) = O(\eta^{-2})$  as  $\eta \rightarrow +\infty$ .

Let us show that

$$u(\eta) = \eta^2 + o(\eta^2), \quad (3.40)$$

where  $C$  is a constant depending on  $u$ . Note that the function  $\phi(\eta)$  is increasing since it satisfies

$$-\phi'' + a(\eta)\phi' = \frac{3\phi}{2},$$



and integrating this equation from  $\eta$  to  $+\infty$  we obtain

$$\phi'(\eta) = \frac{3}{2} \int_{\eta}^{+\infty} \exp\left(-\int_{\eta}^{\xi} a(\zeta) d\zeta\right) \phi(\xi) d\xi \geq 0.$$

It follows that

$$\lim_{\eta \rightarrow +\infty} u(\eta) = +\infty. \quad (3.41)$$

In order to improve this estimate to (3.40) we start with the inequality

$$u''(\eta) \geq \frac{3u'(\eta)}{\sqrt{8u(\eta)}} - \frac{C}{\eta^2},$$

which follows from (3.39) and holds for  $\eta > 0$ , and integrate it from 1 to  $\eta$ :

$$u'(\eta) \geq \frac{3\sqrt{u(\eta)}}{\sqrt{2}} - C.$$

Using (3.41) we conclude that  $u(\eta) \geq C\eta^2$  with  $C > 0$  and in particular

$$l = \liminf_{\eta \rightarrow +\infty} \frac{u(\eta)}{\eta^2} > 0.$$

Then for any  $\delta > 0$  we can find  $\eta(\delta)$  so that  $u(\eta) \geq (l - \delta)\eta^2$  for all  $\eta > \eta(\delta)$ . Going back to (3.39) we observe that for  $\eta > \eta(\delta)$  we have

$$u''(\eta) \geq \frac{3u'(\eta)}{\sqrt{\eta^2 + 8u}} - f(\eta) \geq \frac{3u'(\eta)}{\sqrt{\frac{\eta^2}{l - \delta} + 8u}} - \frac{C}{\eta^2} \geq \frac{3u'(\eta)}{\sqrt{\frac{u}{l - \delta} + 8u}} - \frac{C}{\eta^2}.$$

Integrating this inequality between  $\eta(\delta)$  and  $\eta$  we obtain for  $\eta$  sufficiently large:

$$u'(\eta) \geq \frac{6\sqrt{u(\eta)}}{\sqrt{8 + \frac{1}{l - \delta}}} - C(\delta) \geq \frac{(6 - \delta)\sqrt{u(\eta)}}{\sqrt{8 + \frac{1}{l - \delta}}}. \quad (3.42)$$

We used (3.41) in the last step above. Therefore, we have

$$u(\eta) \geq \left[ \frac{(6 - \delta)}{2\sqrt{8 + \frac{1}{l - \delta}}} \eta - C(\delta) \right]^2,$$

as  $\eta \rightarrow +\infty$ . It follows that for any  $\delta > 0$  we have

$$l \geq \frac{(6 - \delta)^2}{4\left(8 + \frac{1}{l - \delta}\right)}.$$

Passing to the limit  $\delta \rightarrow 0$  we see that

$$l \geq \frac{9}{\left(8 + \frac{1}{l}\right)},$$

and hence  $l \geq 1$ .

On the other hand, it follows from (3.38) that

$$L = \limsup_{\eta \rightarrow +\infty} \frac{u(\eta)}{\eta^2} < +\infty.$$

Then (3.39) implies for any  $\delta > 0$  and  $\eta > \eta(\delta)$ :

$$u''(\eta) = \frac{3u'(\eta)}{\sqrt{\eta^2 + 8u}} - f(\eta) \leq \frac{3u'(\eta)}{\sqrt{\frac{u}{L+\delta} + 8u}} + \frac{C}{\eta^2}.$$

Integrating this inequality between 1 and  $\eta$  we obtain

$$u'(\eta) \leq \frac{6\sqrt{u(\eta)}}{\sqrt{\frac{1}{L+\delta} + 8}} + C(\delta) \leq \frac{(6+\delta)\sqrt{u(\eta)}}{\sqrt{\frac{1}{L+\delta} + 8}} \quad (3.43)$$

for  $\eta > \eta(\delta)$ . Therefore, we have

$$u(\eta) \leq \left[ \frac{(6+\delta)\eta}{2\sqrt{\frac{1}{L+\delta} + 8}} + C(\delta) \right]^2.$$

In the limit  $\eta \rightarrow +\infty$  we obtain

$$L \leq \frac{(6+\delta)^2}{4\left(\frac{1}{L+\delta} + 8\right)},$$

which in the limit  $\delta \rightarrow 0$  becomes  $L \leq 1$ . As  $1 \geq L \geq l \geq 1$ , we conclude that  $L = l$  and

$$\lim_{\eta \rightarrow +\infty} \frac{u(\eta)}{\eta^2} = 1,$$

so that (3.40) indeed holds. Moreover, as  $\phi = u'$ , it follows now from (3.42) and (3.43) that

$$\lim_{\eta \rightarrow +\infty} \frac{\phi(\eta)}{\eta} = 2.$$

**Step 7: The second term in expansion (3.7).** Going back to  $u(\eta)$  defined by the expression (3.20), and  $f(\eta)$  defined by (3.39), we set

$$u(\eta) = \eta^2 + v(\eta). \quad (3.44)$$

The equation for  $v$  is

$$v'' - \frac{v'}{\eta} + \frac{8v}{9\eta^2} = -\frac{4vv'}{9\eta^3} - f(\eta) + \frac{v' + 2\eta}{\eta} g\left(\frac{v}{\eta^2}\right), \quad (3.45)$$

where  $g(\eta)$  is a smooth function such that  $g(0) = g'(0) = 0$ . An asymptotic equation for (3.44) is

$$v'' - \frac{v'}{\eta} + \frac{8v}{9\eta^2} = 0, \quad (3.46)$$

which has two solutions:  $\eta^{4/3}$  and  $\eta^{2/3}$ . We expect the function  $v$  to be asymptotic to  $\eta^{4/3}$  and  $v' = \phi - 2\eta$  to be asymptotic to  $4\eta^{1/3}/3$ , which would essentially finish the proof: the behavior of  $\phi'$  would be obtained by looking at (3.44).

**Lemma 3.7** *If  $v = u(\eta) - \eta^2$  is our sought for solution of (3.45), consider the differential equation with unknown  $w$ :*

$$w'' - \left(1 + g\left(\frac{v}{\eta^2}\right)\right) \frac{w'}{\eta} + \frac{8}{9\eta^2} \left(1 + \frac{v'}{2\eta}\right) w = 0. \quad (3.47)$$

*For every small enough  $\varepsilon > 0$ , there exists  $\eta^\varepsilon \geq \eta_0$  and a solution of (3.47), denoted by  $w^\varepsilon(\eta)$ , such that the following inequalities hold on  $[\eta^\varepsilon, +\infty)$ :*

$$\eta^{4/3-\varepsilon} \leq |w^\varepsilon(\eta)| \leq \eta^{4/3+\varepsilon}, \quad |w'_\varepsilon(\eta)| \leq \eta^{1/3+\varepsilon}, \quad \left| \frac{w'_\varepsilon(\eta)}{w_\varepsilon(\eta)} - \frac{4}{3\eta} \right| \leq \frac{\varepsilon}{\eta}. \quad (3.48)$$

**Proof.** We drop for simplicity the subscripts and superscripts, and use once again the (slightly adjusted) logarithmic derivative of  $w$ :  $q(\eta) = w'(\eta)/w(\eta) - 4/(3\eta)$ . We have, using (3.47):

$$q' + \frac{5 - 3g(v/\eta^2)}{3\eta} q = -q^2 + \frac{4}{3\eta^2} g\left(\frac{v}{\eta^2}\right) - \frac{4}{9\eta^2} \frac{v'}{\eta} := -q^2 + \frac{h_0(\eta)}{\eta^2}. \quad (3.49)$$

If  $h$  were identically equal to zero we could take  $q = 0$  (recall that we are simply looking for one solution  $w$  of (3.47)) – this is, of course, not the case. Let us pick  $\varepsilon > 0$ . From Step 6 we know that there exists  $\eta^\varepsilon > 0$  such that

$$\forall \eta \geq \eta^\varepsilon, \quad 3 \left| g\left(\frac{v}{\eta^2}\right) \right| + |h_0(\eta)| \leq \varepsilon^2. \quad (3.50)$$

Let us look for a solution of (3.47) which has  $q(\eta_\varepsilon) = 0$ , then there exists a constant  $C > 0$  independent of  $\varepsilon$  such that

$$|q(\eta)| \leq C \int_{\eta^\varepsilon}^{\eta} \left(\frac{\eta'}{\eta}\right)^{5/3-C\varepsilon^2} \left(q^2 + \frac{\varepsilon^2}{\eta'^2}\right) d\eta'. \quad (3.51)$$

We conclude by a classical stability argument: let  $\eta_1$  be the first point where the inequality  $q(\eta) \leq \sqrt{\varepsilon}\eta^{-1}$  is violated; if  $\eta_1 < +\infty$  equation (3.49) implies – for  $\varepsilon$  small enough:  $q(\eta_1) \leq C\varepsilon/\eta_1$ , a contradiction. Therefore  $\eta_1 = +\infty$ , proving the desired inequality for  $w$ . The estimate for  $w'$  is obtained in a similar way.  $\square$

A similar argument shows that the other fundamental solution of (3.47) satisfies the estimate  $|\tilde{w}_\varepsilon(\eta)| \leq C_\varepsilon \eta^{2/3+\varepsilon}$  – we leave the details for the reader.

**End of the proof of Theorem 3.1.** We apply Lemma 3.7 twice, by suitably varying the right hand side and the coefficients in the equation (3.45). Let us first observe that, if  $h \in C(\mathbb{R}_+)$  and a small  $\varepsilon > 0$ , and  $\eta_\varepsilon > 0$  are given so that Lemma 3.7 holds, the problem

$$w'' - \left(1 + g\left(\frac{v}{\eta^2}\right)\right) \frac{w'}{\eta} + \frac{8}{9\eta^2} \left(1 + \frac{v'}{2\eta}\right) w = h(\eta), \quad w(\eta_\varepsilon) \text{ and } w'(\eta_\varepsilon) \text{ given}, \quad (3.52)$$

has a unique solution of the form

$$w(\eta) = w_{\text{free}}(\eta) + w^\varepsilon(\eta) \int_{\eta_\varepsilon}^{\eta} \int_{\eta_\varepsilon}^{\eta'} \exp\left(\int_{\eta'}^{\eta''} \left(2 \frac{w'_\varepsilon(\zeta)}{w_\varepsilon(\zeta)} - \frac{1 + g(v/\zeta^2)}{\zeta}\right) d\zeta\right) \frac{h(\eta'')}{w_\varepsilon(\eta'')} d\eta'' d\eta', \quad (3.53)$$

where  $w_{\text{free}}(\eta)$  is the solution of (3.47) with the data  $(w(\eta_\varepsilon), w'(\eta_\varepsilon))$ , and  $w_\varepsilon$  is the particular solution found in Lemma 3.7. However, if  $h = f$  we have  $w = v$  and  $h(\eta) = O(\eta^{-2})$ . From Lemma 3.7 we have

$$2 \frac{w'_\varepsilon(\zeta)}{w_\varepsilon(\zeta)} - \frac{1 + g(v/\zeta^2)}{\zeta} \geq \left(\frac{5}{3} - C\varepsilon\right) \frac{1}{\zeta},$$

with a constant  $C > 0$  independent of  $\varepsilon$ . Lemma 3.7 and the remark following its proof imply that we have  $|w_{\text{free}}(\eta)| \leq C_\varepsilon \eta^{4/3+\varepsilon}$ . Now, it follows from (3.53) with  $h = f$  that

$$|v(\eta)| \leq C_\varepsilon \eta^{4/3+\varepsilon} + \eta^{4/3+\varepsilon} \int_{\eta_\varepsilon}^\eta \int_{\eta_\varepsilon}^{\eta'} \left( \frac{\eta''}{\eta'} \right)^{5/3-C\varepsilon} \frac{|f(\eta'')|}{\eta''^{4/3-\varepsilon}} d\eta'' d\eta' \leq C_\varepsilon \eta^{4/3+C\varepsilon}. \quad (3.54)$$

In a similar way we may obtain

$$|v'(\eta)| \leq C_\varepsilon \eta^{1/3+C\varepsilon}. \quad (3.55)$$

We set now

$$h_1(\eta) = -\frac{4vv'}{9\eta^3} - f(\eta) + \frac{v' + 2\eta}{\eta} g \left( \frac{v}{\eta^2} \right).$$

We have just proved the existence of  $C > 0$  such that, for every small  $\varepsilon > 0$  there are two large constants  $\eta_\varepsilon$  and  $C_\varepsilon$  for which we have

$$\forall \eta \geq \eta_\varepsilon, \quad |h_1(\eta)| \leq \frac{C_\varepsilon}{\eta^{4/3-C\varepsilon}}. \quad (3.56)$$

Fix now  $\varepsilon > 0$  once and for all such that

$$\frac{5}{3} - C\varepsilon > 1, \quad \text{i.e.} \quad \varepsilon < \frac{2}{3C}. \quad (3.57)$$

We have:

$$\begin{aligned} v(\eta) &= v_{\text{free}}(\eta) + \eta^{4/3} \int_{\eta_\varepsilon}^\eta \int_{\eta_\varepsilon}^{\eta'} \left( \frac{\eta''}{\eta'} \right)^{5/3} \frac{h_1(\eta'')}{\eta''^{4/3}} d\eta'' d\eta' \\ v'(\eta) &= v'_{\text{free}}(\eta) + \frac{4}{3} \eta^{1/3} \int_{\eta_\varepsilon}^\eta \int_{\eta_\varepsilon}^{\eta'} \left( \frac{\eta''}{\eta'} \right)^{5/3} \frac{h_1(\eta'')}{\eta''^{4/3}} d\eta'' d\eta' + \eta^{4/3} \int_{\eta_\varepsilon}^\eta \left( \frac{\eta'}{\eta} \right)^{5/3} \frac{h_1(\eta')}{\eta'^{4/3}} d\eta', \end{aligned} \quad (3.58)$$

where  $v_{\text{free}}(\eta)$  is the solution of (3.46), with  $(v_{\text{free}}(\eta_\varepsilon), v'_{\text{free}}(\eta_\varepsilon)) = (v(\eta_\varepsilon), v'(\eta_\varepsilon))$ ; hence it is a linear combination of  $\eta^{4/3}$  and  $\eta^{2/3}$ . As for the integral term, we observe that

$$\left( \frac{\eta''}{\eta'} \right)^{5/3} \frac{h_1(\eta'')}{\eta''^{4/3}} = O(\eta'^{-5/3} \eta''^{C\varepsilon-1}).$$

Hence, because of (3.57), the integral

$$\int_{\eta_\varepsilon}^{+\infty} \int_{\eta_\varepsilon}^{\eta'} \left( \frac{\eta''}{\eta'} \right)^{5/3} \frac{h(\eta'')}{\eta''^{4/3}} d\eta'' d\eta'$$

is finite; call it  $I$ . We have therefore  $v(\eta) - v_{\text{free}}(\eta) = (I + o(1))\eta^{4/3}$ , and a similar identity may be proved for  $v'(\eta) - v'_{\text{free}}(\eta)$  by examining the equation for  $v'(\eta) = \phi(\eta) - 2\eta$  in (3.58). This ends the proof of Theorem 3.1.  $\square$

## 4 Combustion waves

In this section we seek traveling wave profiles that will play the role of the inner waves observed in Section 2. Recalling that  $\theta > 0$  is the ignition temperature, let us pick  $\phi_0 \in [0, \theta)$  and investigate the following differential system, with unknowns  $(c, \phi, \psi)$ :

$$\begin{aligned} -\phi'' + (c + \psi)\phi' &= f(\phi) \\ (c + \psi)\psi' &= \rho(\phi - \phi_0) \end{aligned} \quad (4.1)$$

$$\begin{aligned} \phi(-\infty) &= \phi_0, & \phi(+\infty) &= 1 \\ \psi(-\infty) &= 0, & \psi(+\infty) &= +\infty \end{aligned}$$

In Theorem 4.1 below we'll present the two cases ( $c > 0, \phi_0 = 0$ ) and ( $c = 0, \phi_0 > 0$ ). The first case, described in part (i) of the theorem, represents the left-going traveling waves that were observed numerically in Section 2 when the gravity  $\rho \in (0, \rho_{cr1})$  is sufficiently small. The second case corresponds to the numerically observed profiles that connect the wave fan on the left to a shock on the right, for  $\rho \in (\rho_{cr1}, \rho_{cr2})$ . The critical threshold  $\rho_{cr2}$  appears in the numerical simulations because the initial data vanishes far on the right: this leads to a shock, and opens the way to quenching for large  $\rho$ . If the initial data for temperature have the value  $T \rightarrow 1$  as  $x \rightarrow +\infty$  at  $t = 0$ , the solutions would have the form of a wave fan followed by a traveling wave for all  $\rho > \rho_{cr1}$ . This is reflected in part (ii) of the following theorem.

**Theorem 4.1** (i). *Assume that  $\phi_0 = 0$ . Then there exists  $\rho_{max} > 0$  such that system (4.1) has no solution  $(c, \phi, \psi)$  with  $c \geq 0$  for all  $\rho \geq \rho_{max}$ . If  $\rho$  is small enough, there exists  $c > 0$  such that the system (4.1) has a solution  $(\phi, \psi)$ .*

(ii). *Assume that  $c = 0$ . If  $\rho > 0$  is large enough, there exists  $\phi_0 \in (0, \theta)$  such that system (4.1) has a solution. If  $\phi_0(\rho)$  is the smallest of all  $\phi_0 \in (0, \theta)$  such that system (4.1) has a solution, then we have*

$$\lim_{\rho \rightarrow +\infty} \phi_0(\rho) = \theta. \quad (4.2)$$

The second statement in part (ii) is essential for the quenching phenomenon – if the wave fan does catch up with the shock, the temperature drops below the value  $\phi_0$  everywhere and it is important that this value be close to  $\theta$ .

Note that if the smooth reaction term is replaced by the Dirac mass  $\delta_{\phi=1}$ , the proof of existence of a traveling wave is much simpler. Recall that this regime (see, for instance, [3]) is the limit of a sequence of reaction terms with high activation energies. Then, we have an explicit expression for a travelling wave

$$\phi(x) = \begin{cases} \phi_0 \rho^{-1/2} \phi_\lambda(x) & \text{on } \mathbb{R}_- \\ 1 & \text{on } \mathbb{R}_+ \end{cases},$$

where  $\lambda = (16/[(\theta - \phi_0)\sqrt{\rho}])^{1/3}$ , the family  $\phi_\lambda(x)$  is defined by (4.4) below, while  $\phi_0$  is adjusted to satisfy the derivative jump  $[\phi'](0) = 1$ . We will not pursue this direction. The price to pay for this very simple existence proof is indeed a more difficult study of the quenching – where we crucially use the fact that the reaction term is globally Lipschitz.

Before we start proving anything about (4.1), let us note that any of its solutions satisfies  $\psi' > 0$ , hence, it may be reduced to

$$-\phi'' + \left[ c^2 + 2\rho \int_{-\infty}^x (\phi(y) - \phi_0) dy \right]^{\frac{1}{2}} \phi' = f(\phi) \quad (4.3)$$

$$\phi(-\infty) = \phi_0, \quad \phi(+\infty) = 1.$$

As in the proof of Proposition 3.1 in the previous section, we proceed in several steps.

**Step 1. Nonexistence.** Our primary concern here is what happens as  $x \rightarrow -\infty$  in (4.3), for different values of  $\phi_0$ . For this we need a Liouville type lemma.

**Lemma 4.2** *Consider the family of functions  $(\phi_\lambda^-)_{\lambda \in \mathbb{R}}$  – the dependence on  $\rho$  is omitted, for simplicity:*

$$\forall x < \lambda, \quad \phi_\lambda^-(x) = -\frac{16}{(x - \lambda)^3}. \quad (4.4)$$

*The only increasing solutions  $\phi$  of the equation*

$$-\phi'' + \left( 2\rho \int_{-\infty}^x \phi(y) dy \right)^{\frac{1}{2}} \phi' = 0, \quad \phi(-\infty) = 0 \quad (4.5)$$

have the form  $\rho^{-1/2}\phi_\lambda^-$ .

**Proof.** It suffices to set  $\rho = 1$ , the complete result then follows by scaling. Set

$$u(x) = \int_{-\infty}^x \phi(y) dy, \quad \phi(x) = \eta(u(x)), \quad (4.6)$$

where  $\phi$  is a solution of (4.5). An equation for  $\eta(u)$  is

$$-\frac{d}{du} \left( \eta \frac{d\eta}{du} \right) + \sqrt{2u} \frac{d\eta}{du} = 0, \quad \eta(0) = 0, \quad \frac{d\eta}{du} > 0. \quad (4.7)$$

An explicit solution of (4.7), derived from  $\phi_\lambda^-$ , is  $\eta(u) = u^{3/2}/\sqrt{2}$ . Inspired by this explicit solution we introduce the new unknown

$$p(u) = \eta(u)^{2/3}, \quad (4.8)$$

which in turn satisfies

$$-(p^2 p')' + \sqrt{2u} p^{1/2} p' = 0, \quad p(0) = 0, \quad p' > 0. \quad (4.9)$$

**Claim 1. The function  $p'$  is locally bounded on  $\mathbb{R}_+$ .** To see this, observe that since  $\phi$  is increasing, and  $\phi' > 0$ , we have  $\phi'' \leq C\phi'$  for  $x < 0$ , with  $C = \left(2 \int_{-\infty}^0 \phi(y) dy\right)^{1/2}$ . This implies  $\phi' \leq C\phi$  or, equivalently,  $\sqrt{p}p' \leq C$ . Equation (4.9) may then be integrated from 0 to yield

$$p^2 p' = \int_0^u \sqrt{2v} p p' dv. \quad (4.10)$$

Because  $p' > 0$  it follows that

$$p^2 p' \leq \sqrt{2u} p \int_0^u p' dv = \sqrt{2u} p^{3/2}. \quad (4.11)$$

Hence, we have  $p^{3/2} \leq Cu^{3/2}$ , or  $p \leq Cu$ . We may use this information in (4.10) and infer that

$$p^2 p' \geq C \int_0^u p p' dv = \frac{3}{4\sqrt{2}} p^2,$$

so that  $p' \geq C > 0$  and  $p(u) \geq Cu$ . Now, we may go back to (4.11) and conclude that  $p' \leq C_2 < +\infty$ .

**Claim 2. The derivative  $p'(u)$  has a limit as  $u \rightarrow 0$ .** Expand (4.9) to get

$$-\frac{pp''}{2} - p'^2 + \sqrt{\frac{u}{2p}} p' = 0, \quad p(0) = 0, \quad (4.12)$$

and set

$$\underline{l} = \liminf_{u \rightarrow 0} p'(u), \quad \bar{l} = \limsup_{u \rightarrow 0} p'(u).$$

Assume that  $\underline{l} < \bar{l}$ ; then there exist two sequences  $\underline{u}_n$  and  $\bar{u}_n$ , going to 0 as  $n \rightarrow +\infty$ , such that (i) we have  $\lim_{n \rightarrow +\infty} p'(\underline{u}_n) = \underline{l}$ , and  $\lim_{n \rightarrow +\infty} p'(\bar{u}_n) = \bar{l}$ ; (ii)  $\bar{u}_n$  and  $\underline{u}_n$  are, respectively, a local maximum and a local minimum for  $p'$ . Equation (4.12) implies

$$p'(\underline{u}_n) = \sqrt{\frac{\underline{u}_n}{2p(\underline{u}_n)}}, \quad p'(\bar{u}_n) = \sqrt{\frac{\bar{u}_n}{2p(\bar{u}_n)}}.$$

By the mean value theorem, there is  $(\underline{v}_n, \bar{v}_n)$  such that  $0 \leq \underline{v}_n \leq \underline{u}_n$ ,  $0 \leq \bar{v}_n \leq \bar{u}_n$ , so that

$$p'(\underline{u}_n) = \sqrt{\frac{1}{2p'(\underline{v}_n)}}, \quad p'(\bar{u}_n) = \sqrt{\frac{1}{2p'(\bar{v}_n)}},$$

yielding in turn

$$\underline{l} \geq \sqrt{\frac{1}{2\bar{l}}}, \quad \bar{l} \leq \sqrt{\frac{1}{2\underline{l}}}$$

and, finally  $-\underline{l} \geq \bar{l}$ , a contradiction. As a consequence, we have  $\underline{l} = \bar{l} = p'(0) = 2^{-1/3}$ .

**Claim 3. We have**  $p(u) = u/2^{1/3}$ . Assume first  $p'$  to have both a global minimum  $\underline{u}_0$  and a global maximum  $\bar{u}_0$  on an interval  $(0, r)$ . Equation (4.12) and the mean value theorem imply that

$$p'(\underline{u}_0) \geq \sqrt{\frac{1}{2p'(\bar{u}_0)}}, \quad p'(\bar{u}_0) \leq \sqrt{\frac{1}{2p'(\underline{u}_0)}}$$

and thus  $p'(\underline{u}_0) = p'(\bar{u}_0)$ . If  $p'$  has neither a minimum nor a maximum on intervals of the form  $(0, r)$ , then  $p''$  has a constant sign; assume  $p'' \geq 0$ . Equation (4.12) then implies

$$p'(u) \leq \sqrt{\frac{u}{2p(u)}} \leq \sqrt{\frac{1}{2p'(0)}} = p'(0).$$

Hence, in this case we have  $p'(u) = p'(0)$ . The case  $p'' \leq 0$  is treated in the same fashion. The only cases that remain to be treated are (i) when  $p'$  has a global minimum but no global maximum, and (ii) the converse case. Assume for instance that (i) holds. Then there exists  $u_1 > 0$  such that  $p'' \leq 0$  on  $(0, u_1)$  and  $p'' \geq 0$  on  $\{u \geq u_1\}$ . Once again, we apply at that point both equation (4.9) and the mean value theorem: there exists  $u_2 \in (0, u_1)$  such that

$$p'(u_1) = \sqrt{\frac{1}{2p'(u_2)}} \geq 2^{-1/3} = p'(0).$$

Consequently,  $p'$  is constant on  $[0, u_1]$ , hence everywhere else.  $\square$

**Proof of the nonexistence part of Theorem 4.1.** Let us first take  $\phi_0 = 0$  and prove that when  $\rho > 0$  is sufficiently large, equation (4.3) has no solution for any  $c > 0$ . If  $\phi$  is a solution, note that we have, classically:  $\phi' > 0$ . This is seen by the standard integration of (4.3) from  $x$  to  $+\infty$ , taking  $f(\phi)$  as a non-negative right side.

Let us multiply (4.3) by  $\phi'$  and integrate on  $\mathbb{R}$ . This yields

$$\int_{-\infty}^0 \left( c^2 + 2\rho \int_{-\infty}^x \phi(y) dy \right)^{1/2} \phi'^2(x) dx \leq \int_0^1 f(\phi) d\phi := M. \quad (4.13)$$

We always may assume that  $\phi(0) = \theta$ . Therefore, as  $\phi$  is increasing,  $\phi(x) \leq \theta$  for  $x \leq 0$  so that  $f(\phi) = 0$  there and we have:

$$-\phi'' + \left( c^2 + 2\rho \int_{-\infty}^x \phi(y) dy \right)^{1/2} \phi' = 0, \quad \text{for } x \leq 0 \text{ and } \phi(0) = \theta. \quad (4.14)$$

Hence

$$\left( c^2 + 2\rho \int_{-\infty}^x \phi(y) dy \right)^{1/2} = \frac{\phi''}{\phi'}$$

and inequality (4.13) becomes:

$$(\phi'(0))^2 \leq 2M. \quad (4.15)$$

We therefore have to estimate  $\phi'(0)$  in terms of  $\rho$ . Let us first note that  $\phi(x) \leq \theta e^{cx}$ , which, as  $\phi(0) = \theta$ , implies

$$\phi'(0) \geq c\theta, \quad \text{hence, by (4.15):} \quad c \leq \sqrt{\frac{2M}{\theta}}. \quad (4.16)$$

To proceed further let us make the following change of the unknown variable, as in the proof of Lemma 4.2:

$$u(x) = \int_{-\infty}^x \phi(y) dy, \quad \phi(x) = \eta(u(x)) \quad (4.17)$$

where  $\phi$  is a solution of (4.14). An equation for  $\eta(u)$  is

$$-\frac{d}{du} \left( \eta \frac{d\eta}{du} \right) + (c^2 + 2\rho u)^{1/2} \frac{d\eta}{du} = 0, \quad \eta(0) = 0, \quad \frac{d\eta}{du} > 0. \quad (4.18)$$

Then we set

$$p(u) = \eta(u)^{2/3}, \quad (4.19)$$

which in turn satisfies

$$-(p^2 p')' + ((c^2 + 2\rho u)p)^{1/2} p' = 0, \quad p(0) = 0, \quad p' > 0. \quad (4.20)$$

As in the proof of Claim 1 in Lemma 4.2, we have

$$p^2 p' = \int_0^v \sqrt{(c^2 + 2\rho w)p(w)} p'(w) dw, \quad (4.21)$$

and hence – by the same argument as in the proof of Claim 1 in Lemma 4.2 we get

$$p^2 p' = \int_0^u \sqrt{(c^2 + 2\rho w)p(w)} p'(w) dw \leq \sqrt{(c^2 + 2\rho u)p(u)} p(u).$$

It follows that

$$p(u)^{3/2} \leq K \int_0^u \sqrt{c^2 + 2\rho w} dw \leq K\sqrt{\rho} \left[ \left( \frac{c^2}{2\rho} + u \right)^{3/2} - \left( \frac{c^2}{2\rho} \right)^{3/2} \right] \leq \frac{K}{\rho} (c^2 + 2\rho u)^{3/2}. \quad (4.22)$$

As a consequence, we have

$$p(u) \leq \frac{K}{\rho^{2/3}} (c^2 + 2\rho u).$$

This, in turn, implies

$$p^2 p' = \int_0^u \sqrt{(c^2 + 2\rho w)p(w)} p'(w) dw \geq K\rho^{1/3} \int_0^u p(w) p'(w) dw = K\rho^{1/3} p^2(u)$$

so that  $p(u) \geq K\rho^{1/3}u$ . The function  $\phi(x)$  thus satisfies the inequality

$$\phi(x) \geq K\rho^{1/2}u^{3/2},$$

which may be re-written as

$$u'(x) \geq K\rho^{1/2}u^{3/2}. \quad (4.23)$$



The second inequality in (4.22) implies also the following upper bound for  $u'(x)$ :

$$u'(x) \leq \frac{K}{\rho} \left[ (c^2 + 2\rho u)^{3/2} - c^3 \right]. \quad (4.24)$$

Recall that  $\phi(0) = u'(0) = \theta$  – hence, we have from (4.23):

$$u(0) \leq K \left( \frac{\theta}{\rho^{1/2}} \right)^{2/3} \leq \frac{K}{\rho^{1/3}}, \quad (4.25)$$

while from (4.24) we obtain

$$u(0) \geq \frac{K}{\rho^{1/3}} - \frac{c^2}{2\rho} \geq \frac{K}{2\rho^{1/3}} \quad (4.26)$$

for  $\rho > \rho_0$ . We used (4.16) in the last step above. Another consequence of (4.23) is that for  $x \leq 0$  we have

$$\frac{1}{\sqrt{u(x)}} - \frac{1}{\sqrt{u(0)}} \geq K\rho^{1/2}|x|,$$

so that

$$u(x) \leq \frac{u(0)}{(1 + \sqrt{u(0)}K\rho^{1/2}|x|)^2}.$$

With the help of (4.25) and (4.26) this becomes

$$u(x) \leq \frac{K}{\rho^{1/3}} \left( 1 + \frac{K}{\rho^{1/6}}\rho^{1/2}|x| \right)^{-2} \leq \frac{K}{(\rho^{1/6} + \rho^{1/2}|x|)^2}.$$

As a consequence, we have  $u(-\rho^{-m}) \leq K/\rho^{1-2m}$  for  $m \in (0, 1/3)$  and thus we get from (4.24):

$$\phi(\rho^{-m}) \leq \frac{K}{\rho}\rho^{3m} = K\rho^{3m-1} \ll 1.$$

As  $\phi(0) = \theta$ , there exists a point  $\xi \in (-\rho^{-m}, 0)$  so that  $\phi'(\xi) \geq K\rho^m$ . However, the function  $\phi$  is convex, thus  $\phi'(0) \geq K\rho^m$  which contradicts (4.15) if  $\rho$  is large enough. This contradiction shows that no solution may exist for a sufficiently large  $\rho$ .

**Step 2. Existence for small  $\rho$ .** The reason why solutions may exist when  $\rho$  is small is quite easy to understand: when  $\rho = 0$ , (4.3) reduces to

$$-\phi'' + c\phi' = f(\phi), \quad \phi(-\infty) = 0, \quad \phi(+\infty) = 1,$$

which, of course, has a unique solution  $(c_0, \bar{\phi})$ , up to translation, when the nonlinearity  $f$  is of the ignition type. The idea is that the standard non-degeneracy property of this solution allows its continuation for tiny values of  $\rho$ . However, there is a technical point that makes the program not completely trivial: the term  $\rho \int_{-\infty}^x \phi(y) dy$  grows linearly as  $x \rightarrow +\infty$ , whereas we would like to treat it as a small perturbation. This forces us to work a little more with the a priori estimates. The main element that will make a perturbation argument work is the following

**Lemma 4.3** *There exists  $\rho_0 > 0$  such that, for all  $\rho \leq \rho_0$ , for every  $h(x) \in BUC(\mathbb{R}_+)$  – the space of all bounded, uniformly continuous functions on  $\mathbb{R}_+$  – there exists a unique  $u(x) \in BUC(\mathbb{R}_+)$  satisfying the boundary value problem*

$$L_\rho u := -u'' + \left( c_0^2 + 2\rho \int_{-\infty}^x \bar{\phi}(y) dy \right)^{\frac{1}{2}} u' - f'(\bar{\phi})u = h \quad (x > 0), \quad u(0) = 0. \quad (4.27)$$

Moreover there exists  $C > 0$  independent of  $\rho$  such that

$$\|u\|_\infty \leq C\|h\|_\infty, \quad (4.28)$$

and the map  $\rho \mapsto (u(0), u'(0))$  is continuous on  $[0, \rho_0]$ . Finally, if in addition  $h \in L^1(\mathbb{R}_+)$ , we have

$$\|u\|_{W^{1,1}(\mathbb{R}_+) \cap W^{1,\infty}(\mathbb{R}_+)} \leq C\|h\|_{L^1(\mathbb{R}_+) \cap L^\infty(\mathbb{R}_+)}. \quad (4.29)$$

**Proof.** What really matters is the estimate (4.28). Once this property is at hand, (4.27) may be approximated by the following sequence of problems

$$L_n u := -u'' + \left( c_0^2 + 2\rho \int_{-\infty}^{\max(x,n)} \bar{\phi}(y) dy \right)^{\frac{1}{2}} u' - f'(\bar{\phi})u = h \quad (x > 0), \quad u(0) = 0. \quad (4.30)$$

The operator  $L_n$  is Fredholm, and estimate (4.28) is – as will become clear in the proof of the lemma – still valid for the solutions of (4.30). One then concludes by a standard compactness argument. The continuity of the map  $\rho \mapsto (u(0), u'(0))$  is also inferred from compactness. The estimate (4.29), which is the main result, will easily follow.

Let us therefore assume that we have constructed a solution  $u(x)$  to (4.27), and let us estimate it. Let us re-write (4.27) as

$$-u'' + c_0 u' - f'(\bar{\phi})u = l[u] := h - \frac{2\rho \int_{-\infty}^x \bar{\phi}(y) dy}{c_0 + \sqrt{c_0^2 + 2\rho \int_{-\infty}^x \bar{\phi}(y) dy}} u' \quad (x > 0), \quad u(0) = 0.$$

This problem has an explicit solution

$$u(x) = \int_0^x \frac{\bar{\phi}'(x)}{\bar{\phi}'(y)} \int_y^{+\infty} \frac{\bar{\phi}'(z)}{\bar{\phi}'(y)} e^{c_0(y-z)} l[u](z) dy dz.$$

Using integration by parts to eliminate the derivative of  $u$  in the function  $l$  and the exponential decay of  $\bar{\phi}'(x)$ , we deduce that there exists a constant  $C > 0$  independent of  $\rho$  such that

$$|u(x)| \leq C(\|h\|_\infty + \sqrt{\rho x} \|u\|_\infty). \quad (4.31)$$

Consider now  $x_0 > 0$  and  $\delta > 0$  such that  $-f'(\bar{\phi}) \geq \delta$  on  $[x_0, +\infty)$ . On that interval equation (4.27) and the maximum principle yield that  $u(x)$  can not attain a maximum at a point where its value is larger than  $\|h\|_\infty/\delta$ . On the other hand, if  $u(x)$  is monotonic on an infinite half-interval we set  $u_\varepsilon(x) = u(x) - \varepsilon \bar{u}(x)$ , where  $\bar{u}(x)$  is

$$\bar{u}(x) = \int_0^x \exp\left(\int_0^y a(z) dz\right) dy, \quad a(x) = \left(c_0^2 + 2\rho \int_{-\infty}^x \bar{\phi}(y) dy\right)^{\frac{1}{2}}.$$

As the function  $a(y)$  tends to  $+\infty$  as  $x \rightarrow +\infty$ , it follows that  $u_\varepsilon(+\infty) = -\infty$  and thus has to attain a local maximum. Applying the maximum principle to  $u_\varepsilon$  and passing to the limit  $\varepsilon \rightarrow 0$  we conclude that

$$\forall x \geq x_0, \quad |u(x)| \leq |u(x_0)| + \frac{\|h\|_\infty}{\delta}.$$

Combining this with (4.31) yields

$$\|u\|_{L^\infty([x_0, +\infty))} \leq C(\|h\|_\infty + \sqrt{\rho x_0} \|u\|_\infty) + \frac{\|h\|_\infty}{\delta}. \quad (4.32)$$

But then, (4.31) implies that

$$\|u\|_{L^\infty([0,x_0])} \leq C(\|h\|_\infty + \sqrt{\rho x_0}\|u\|_\infty). \quad (4.33)$$

Adding up (4.32) and (4.33), then choosing  $\rho$  so that  $C\sqrt{\rho x_0} < 1$  yields (4.28). We note that the above argument is valid for the family of operators  $L_n$  given by (4.30): inequality (4.32) does not change as long as  $x_0 \leq n$ , and this inequality does not use any bound on the first order term of  $L_n$ .

Finally, let us assume in addition that  $h \in L^1(\mathbb{R}_+)$ . We set for convenience

$$a(x) = \sqrt{c_0^2 + 2\rho \int_{-\infty}^x \bar{\phi}(y) dy};$$

then we have

$$|a'(x)| \leq \rho c_0^{-1}. \quad (4.34)$$

With this fact in mind, we multiply (4.27) by  $\text{sgn}u(x)$  and integrate it over  $(0, +\infty)$ . We get, after integration by parts:

$$\int_{x_0}^{+\infty} (\delta - a'(x))|u(x)| dx \leq \|h\|_{L^1} + \int_0^{x_0} |f'(\bar{\phi})u| dx.$$

Here  $x_0$  and  $\delta$  are chosen as in the previous step of the proof. The upper bound (4.34) and the  $L^\infty$  bound for  $u$  that we have already obtained imply that for a sufficiently small  $\rho$  we have an  $L^1$ -bound for  $u$ :  $\|u\|_{L^1} \leq C\|h\|_{L^1 \cap L^\infty}$ . In order to improve it to an  $W^{1,1}$  bound we note that the  $L^1$ -estimate for  $u$  and the fact that  $h \in L^1$  imply that

$$u'(x) = - \int_x^{+\infty} g(y) \exp\left(- \int_x^y a(z) dz\right) dy,$$

with a function  $g \in L^1$ . As  $a(x)$  is uniformly bounded from below by a positive constant, it follows that  $u' \in L^1 \cap L^\infty$ .  $\square$

The construction of a solution  $(c, \phi)$  to (4.3) can now be done: it is a classical derivative matching problem. First, let us add to (4.3) the normalization condition  $\phi(0) = \theta$ . We know that a solution  $\phi(x)$  of (4.3) has to be increasing – therefore,  $f(\phi) \equiv 0$  for  $x \leq 0$ . The equation for  $\phi$  is thus

$$-\phi'' + \left(c^2 + 2\rho \int_{-\infty}^x \phi(y) dy\right)^{\frac{1}{2}} \phi' = 0 \quad \text{for } x < 0; \quad \phi(-\infty) = 0, \quad \phi(0) = \theta. \quad (4.35)$$

We pick any  $\mu \in (0, c_0/5)$ , and  $c$  such that  $|c - c_0| < 2\mu$ . The implicit function theorem in the space  $\{u \in BUC(\mathbb{R}), e^{-\mu x}u \in BUC(\mathbb{R}_-)\}$  yields, for  $\rho > 0$  small enough, the existence of a unique solution  $\phi_{c,\rho}^-$  to (4.35). Moreover we have

$$\frac{d\phi_{c,\rho}^-}{dx}(0) = c\theta + O(\rho). \quad (4.36)$$

The details are standard and are therefore omitted.

Let us turn to the problem on the right half line:

$$\begin{aligned} -\phi'' + \left[c^2 + 2\rho \int_{-\infty}^x (\phi(y) - \phi_0) dy\right]^{\frac{1}{2}} \phi' &= f(\phi) \\ \phi(0) = \theta, \quad \phi(+\infty) &= 1, \end{aligned} \quad (4.37)$$

with the additional constraint  $\phi'(0) = \phi_{c,\rho}^-(0)$ . We look for a solution  $(c, \phi)$  to (4.37) in the form  $(c_0 + d, \bar{\phi} + u)$ . We also extend  $\phi$  to  $\phi_{c,\rho}^-$  on  $\mathbb{R}_-$  – this is only necessary to assign a value to the integrals between  $-\infty$  and  $x$  in (4.37). Write the problem as

$$\begin{aligned} L_\rho u &= g[u], \\ u(0) &= 0, \quad u \in W^{1,1}(\mathbb{R}_+), \end{aligned} \tag{4.38}$$

with the operator  $L_\rho$  defined in (4.27),

$$\begin{aligned} g[u] &= K[\phi]u^2 - \frac{(c + c_0)d + 2\rho \int_{-\infty}^x u(y) dy}{\sqrt{c_0^2 + 2\rho \int_{-\infty}^x \bar{\phi}(y) dy} + \sqrt{c^2 + 2\rho \int_{-\infty}^x \bar{\phi}(y) dy}} (u' + \bar{\phi}') \\ &\quad - \frac{2\rho \int_{-\infty}^x u(y) dy}{\sqrt{c_0^2 + 2\rho \int_{-\infty}^x \bar{\phi}(y) dy} + \sqrt{c^2 + 2\rho \int_{-\infty}^x \bar{\phi}(y) dy}} \bar{\phi}' \end{aligned}$$

and  $K[\phi]u^2 = f(\bar{\phi} + u) - f(\bar{\phi}) - f'(\bar{\phi})u$ . Lemma 4.3 asserts that  $L_\rho$  is invertible, and that  $L_\rho^{-1}$  sends  $L^1(\mathbb{R}_+) \cap L^\infty(\mathbb{R}_+)$  to  $W^{1,1}(\mathbb{R}_+) \cap W^{1,\infty}(\mathbb{R}_+)$  and thus equation (4.38) is equivalent to

$$u = L_\rho^{-1}(g[u]). \tag{4.39}$$

The Banach fixed point theorem yields the existence of two positive numbers  $\rho_0$  and  $\delta_0$  such that, for each  $(d, \rho) \in [-\delta_0, \delta_0] \times [0, \rho_0]$ , equation (4.39) has a unique solution  $u_{c,\rho}^+$  of the size  $|d| + \rho$  – and therefore the resulting  $\phi_{c,\rho}^+ = \bar{\phi} + u_{c,\rho}^+$  is  $|d| + \rho$ -close to  $\bar{\phi}$ .

Now, the problem (4.3) reduces to the following equation: given  $\rho$  in some subinterval of  $[0, \rho_0]$  containing 0, find  $c$  close to  $c_0$  so that the equation

$$\frac{d\phi_{c,\rho}^+}{dx}(x=0) = \frac{d\phi_{c,\rho}^-}{dx}(x=0). \tag{4.40}$$

Now, a well-known Melnikov-type computation gives

$$\left. \frac{\partial^2 \phi_{c,\rho}^+}{\partial c \partial x} \right|_{x=0, \rho=0, c=c_0} = -\frac{1}{c_0 \theta} \int_0^{+\infty} e^{-c_0 x} \bar{\phi}^2(y) dy.$$

This, combined with (4.36), implies that (4.40) can be solved uniquely, provided that  $\rho$  is chosen small enough. This ends the small  $\rho$  construction part and the proof of part [i] of Theorem 4.1.

**Step 3. Proof of part [ii] of Theorem 4.1: the large values of  $\rho$ .** Recall that we are looking for a pair  $(\phi_0, \phi)$  satisfying (4.3) with  $c = 0$ . We may impose the normalization condition  $\phi(0) = \theta$ , and the solution  $\phi$  is increasing on  $\mathbb{R}$ . This implies, by Lemma 4.2 that

$$\phi = \phi_0 + \rho^{-1/2} \phi_\lambda \text{ for } x \leq 0, \tag{4.41}$$

with  $\lambda = (16/(\theta - \phi_0)\sqrt{\rho})^{1/3}$ . In particular, we have

$$\int_{-\infty}^0 [\phi(y) - \phi_0] dy = 2^{1/3}(\theta - \phi_0)^{2/3} \rho^{-1/6} := c_\rho(\phi_0), \quad \phi'(0^-) = \frac{3}{2^{4/3}}(\theta - \phi_0)^{4/3} \rho^{1/6}. \tag{4.42}$$

The strategy is, once again, a shooting argument: we are going to solve the following Cauchy problem:

$$\begin{aligned} -\phi'' + \left( c_\rho(\phi_0) + 2\rho \int_0^x \phi \right)^{\frac{1}{2}} \phi' &= f(\phi) \\ \phi(0) = \theta, \quad \phi'(0^+) &= \frac{3}{2^{4/3}}(\theta - \phi_0)^{4/3} \rho^{1/6} \end{aligned} \tag{4.43}$$

and adjust  $\phi_0$  so that

$$\lim_{x \rightarrow +\infty} \phi(x) = 1. \quad (4.44)$$

For all  $\rho > 0$  and  $\phi_0 > 0$  the Cauchy problem (4.43) has a unique maximal solution  $\phi_{\rho, \phi_0}$  defined on an interval of the form  $[0, x_{max}(\rho, \phi_0))$ . Exactly as in Section 3, for each  $\rho > 0$  we define the following subsets of  $[0, \theta]$ :

$$\begin{aligned} X_-^\rho &= \{\phi_0 \in [0, \theta] : \exists x_0 > 0 \text{ such that } \phi_{\rho, \phi_0} = \theta\} \\ X_+^\rho &= \{\phi_0 \in [0, \theta] : \phi_{\rho, \phi_0} > 0 \text{ and } x_{max}(\rho, \phi_0) < +\infty\}, \end{aligned} \quad (4.45)$$

**Lemma 4.4** *For every  $\rho > 0$  there exists  $\bar{\phi}_0(\rho) > 0$  such that every  $\phi_0 \in (\bar{\phi}_0(\rho), \theta]$  belongs to  $X_-^\rho$ .*

**Proof.** Given  $\rho > 0$ , the Cauchy Problem

$$\begin{aligned} -\phi'' &= f(\phi) \\ \phi(0) &= \theta, \quad \phi'(0^+) = \frac{3}{2^{4/3}}(\theta - \phi_0)^{4/3} \rho^{1/6} \end{aligned} \quad (4.46)$$

has – by an easy explicit computation – a unique solution  $\underline{\phi}$  which is larger than  $\theta$  exactly on a finite interval  $(0, \underline{x}(\rho, \phi_0))$  provided that  $\phi_0 > 0$  is close enough to  $\theta$ . The difference between  $\underline{\phi}$  and  $\phi_{\rho, \phi_0}$  is easily estimated via the Gronwall lemma for large  $\rho$  since both  $c_\rho(\phi_0)$  and  $\underline{x}(\rho, \phi_0)$  are small.  $\square$

Because a solution  $\phi$  to (4.3) is increasing in  $x$ , we are not really interested in the values of  $f$  outside the interval  $(0, 1)$ . Therefore we may extend  $f$  by 0 outside  $(0, 1)$ .

**Lemma 4.5** *For every  $\phi_0 < \theta$  there exists  $\rho_0 > 0$  so that we have  $\phi_0 \in X_+^\rho$  for all  $\rho > \rho_0$ .*

**Proof.** Quite the same as in Lemma 3.4. The logarithmic derivative of  $\phi = \phi_{\rho, 0}$ , denoted by  $\zeta$ , satisfies

$$\zeta' = a(x)\zeta - \zeta^2 - \frac{f(\phi)}{\phi}. \quad (4.47)$$

where we have suppressed the subscripts, and where we have set

$$a(x) = \sqrt{c_\rho(\phi) + 2\rho \int_0^x \phi(y) dy}. \quad (4.48)$$

This equation, due to the boundedness of  $f(\phi)/\phi$ , is essentially the same as (3.29). In particular, if  $\zeta(0)$  is large enough we have  $\zeta > 0$  on its existence interval, and  $x_{max}(\rho, 0) < +\infty$ . This implies that  $\phi_0 \in X_+^\rho$  if  $\zeta(0)$  is sufficiently large. However, as  $\zeta(0)$  is proportional to  $\rho^{1/6}$ , this is the case for a sufficiently large  $\rho$ .  $\square$

**End of the proof of Theorem 4.1.** Take  $\rho > 0$  large enough so that Lemma 4.5 holds. As opposed to the construction of the wave fan-rarefaction wave, where the fact that the sets  $X_\pm$  were open was nontrivial, it is here very easy to infer from the continuity of the solution of the Cauchy Problem (4.43) with respect to its initial values, that the sets  $X_\pm^\rho$  are open and non-empty. Consequently, there exists at least one  $\phi_0$  not in  $X_+^\rho \cup X_-^\rho$  – we need now to show that the solution generated by  $\phi_0$  tends to one as  $x \rightarrow +\infty$ . The strong maximum principle implies that the corresponding solution  $\phi$  cannot have a local minimum in  $\mathbb{R}_+$  and hence it is increasing. Assume that  $\phi$  goes over the value 1 and let  $x_0$  be such that  $\phi(x_0) = 1$ . Setting  $\bar{c}_\rho(\phi_0) = c_\rho(\phi_0) + 2\rho \int_0^{x_0} \phi(y) dy$ , the equation for  $\phi$  is

$$\begin{aligned} -\phi'' + \left( \bar{c}_\rho(\phi_0) + 2\rho \int_{x_0}^x \phi \right)^{\frac{1}{2}} \phi' &= 0 \\ \phi(x_0) &= 1, \quad \phi' > 0. \end{aligned} \quad (4.49)$$

Equation (4.49) implies that  $\phi' > 0$  and  $\phi$  is convex on  $(x_0, +\infty)$ . Hence, the derivative  $\phi'(x)$  has a limit  $l > 0$  as  $x \rightarrow +\infty$ . This implies, once again by (4.49):

$$\phi''(x) \sim l\sqrt{\rho l x}$$

and thus  $l$  can not be finite. Therefore, repeating the argument in the proof of Lemma 3.4 we conclude that  $\phi$  becomes infinite at a finite distance, which is a contradiction. Therefore, we have  $\phi < 1$ , hence, (4.44) is true. Finally, Lemma 4.5 implies that (4.2) holds.  $\square$

### Asymptotic behavior of the traveling wave at $+\infty$

We end this section by additional information on the behavior of a traveling wave solution of (4.3). The last lemma of this section is an estimate of how the wave solution converges to its rest state at  $+\infty$  which we will need in the construction of an asymptotic solution when we match the traveling wave to the back of the shock. Due to the linear growth of the advection term, it is not a standard version of the stable manifold theorem. We could, at a not too high cost, derive a precise asymptotic expansion. The following weaker version will be sufficient for our purpose.

**Lemma 4.6** *Let  $\phi(x)$  be a solution of (4.3) and set  $\alpha_0 = (-f'(1))\sqrt{2/\rho}$ . There exists  $B(\rho) > 0$  so that for each  $\varepsilon > 0$ , there exist  $C_\varepsilon(\rho), C'_\varepsilon(\rho) > 0$  such that*

$$1 - C_\varepsilon \exp(-(\alpha_0 - B\varepsilon)\sqrt{x}) \leq \phi(x) \leq 1 - C'_\varepsilon \exp(-(\alpha_0 + B\varepsilon)\sqrt{x}) \quad (4.50)$$

and

$$0 \leq \phi'(x) \leq B e^{-\alpha_0 \sqrt{x}/2}.$$

**Proof.** The function  $q(x) = 1 - \phi(x)$  satisfies

$$NL(q) := -q'' + a(x)q' + f(1 - q) = 0, \quad q(-\infty) = 1, \quad q(+\infty) = 0, \quad (4.51)$$

where

$$a(x) = \left( c^2 + 2\rho \int_{-\infty}^x \phi(y) dy \right)^{1/2}.$$

Let us choose  $x_\varepsilon > 1/\varepsilon^2$  so that for all  $x \geq x_\varepsilon$  we have

$$(-f'(1) - \varepsilon)q \leq f(1 - q) \leq (-f'(1) + \varepsilon)q,$$

and

$$\sqrt{2\rho x}(1 - \varepsilon) \leq a(x) \leq \sqrt{2\rho x}(1 + \varepsilon).$$

Let us find a supersolution  $\bar{q}(x) \geq 0$  such that  $\bar{q}(x_\varepsilon) = q(x_\varepsilon)$ ,  $\bar{q}' \leq 0$ ,  $\bar{q}(+\infty) = 0$  and  $NL[\bar{q}] \geq 0$ . This will imply that  $q(x) \leq \bar{q}(x)$  and thus provide the lower bound on  $\phi(x)$  in (4.50). For the last condition to hold it is sufficient to require that

$$\bar{L}[\bar{q}] := -\bar{q}'' + \sqrt{2\rho x}(1 + \varepsilon)\bar{q}' + (-f'(1) - \varepsilon)\bar{q} \geq 0.$$

For a function of the form  $s(x) = q(x_\varepsilon) \exp(-\alpha\sqrt{x - x_\varepsilon})$  we have

$$\bar{L}[s] = s(x) \left( O\left(\frac{1}{x}\right) + (-f'(1) - \varepsilon) - \frac{\alpha\sqrt{\rho}}{\sqrt{2}}(1 + \varepsilon) \right).$$

Hence, for such function to be a supersolution it is sufficient to take

$$\alpha < \sqrt{\frac{2}{\rho}} \left( \frac{-f'(1) - \varepsilon}{1 + \varepsilon} \right), -\varepsilon$$

so  $\alpha \leq \alpha_0 - B(\rho)\varepsilon$  with  $B(\rho)$  sufficiently large will suffice. The upper bound on  $\phi$  in (4.50) is proved similarly. The bound on the derivative  $\phi'$  in Lemma 4.6 is obtained by differentiating (4.51):

$$-(q')'' + a(x)(q')' + (-f'(1 - q) + a'(x))q' = 0.$$

As  $a'(x) = O(1/\sqrt{x})$  for large  $x$ , the exponential bound for  $q' = -\phi'$  follows by the same construction of subsolutions and supersolutions.  $\square$

## 5 Large-time evolution: asymptotic solutions

The numerical simulations of Section 2 indicate that, if the support of the initial data for temperature  $T(0, x)$  – or, at least, the measure of the set where it is above ignition – is very large, the solution has the following structure: it has the form of a ramp on the left, followed by a combustion wave, which is itself terminated by a shock that brings back both temperature and velocity to their rest states. This structure appears after some transient behavior that we will not study in this paper, and remains valid almost all the time before quenching occurs. In this section we derive an asymptotic relation for the shock position, and discuss the time interval on which the above picture is valid. What happens after this time is the subject of Section 6.

It is clear from the numerical simulations that quenching will be provoked by the dissipation at the accelerated shock, and that the shock location is really what will eventually tell us the dynamics of our solution. In order to get an equation for the shock motion we have to construct the whole asymptotic solution, gluing together asymptotically the ramp, combustion wave and the shock constructed in the previous sections. First, we should identify the small parameter that controls the asymptotics – this is the object of Section 5.1. In Section 5.2, we will see how to glue the ramp to the combustion wave; the role of the ramp being played by the selfsimilar solution constructed in Section 3, and the role of the combustion wave will be played by the traveling wave constructed in Section 4. In Section 5.3 we will place the shock, thus terminating the description of the asymptotic solution.

### 5.1 Devising a length and time scale

Let us recall that if  $y_f(t)$  is the position of the shock relative to the end point of the ramp, that is the transition point in the temperature profile between the ramp and the combustion wave, an asymptotic equation for  $y_f(t)$  is given by (2.13):

$$\dot{y}_f = \frac{1}{2}(\sqrt{2\rho(1 - \phi_0)y_f} - \rho\phi_0 t). \quad (5.1)$$

Here  $\phi_0$  is the value of the temperature at the transition. We will assume that  $\phi_0 > 1/9$  – recall that this ensures that (5.1) has no global in time solution  $y_f(t)$ , hence  $y_f(t)$  reaches zero in a finite time – this is the time when quenching occurs since the transition value  $\phi_0$  is below the ignition temperature  $\theta$ . Let us now worry about how large we should choose the support of the temperature for the solution to maintain the "ramp-wave-shock" structure for a long time – this length will be our large parameter with respect to which we shall expand our solution.

Let us set

$$Q(z) = 2 \int_0^z \frac{dz'}{f(\sqrt{z'})};$$

and choose a pair of large positive numbers:  $(t_0, x_0)$ . We specify the initial datum of  $y_f$  as  $y_f(t_0) = x_0$ . The reason why we wish to start the integration of (5.1) from a large time  $t_0$  will become clear in Section 5.3. In a few words: we want to make sure that the transition layer (the shock width) in which the temperature goes from 1 to 0 is very narrow. From (2.14) we have, for  $t \geq t_0$ , using expression (2.14) for  $f(\sqrt{z})$ :

$$y_f(t) = \rho t^2 Q^{-1}(\ln \frac{t}{t_0} + Q(\frac{x_0}{\rho t_0^2})) \sim_{x_0/(\rho t_0)^2 \rightarrow +\infty} \rho t^2 Q^{-1}(\ln \frac{t}{\sqrt{\rho^{-1}x_0}}). \quad (5.2)$$

Thus, for  $t \sim \sqrt{\rho^{-1}x_0}$  we have  $y_f(t) \sim 0$ , meaning that the shock has been caught up by the ramp - thus, presumably, that quenching has occurred around that time. The parameter  $x_0$  will from then be the large parameter; we call it  $\varepsilon^{-1}$ , with  $\varepsilon > 0$ . The time interval over which we want to construct an approximate solution to (1.5) runs from  $t_0$  to approximately  $(\rho\varepsilon)^{-1/2}$ . Recall that we want  $t_0$  also very large; call it  $\delta^{-1}$  and  $\delta$  will be another small parameter. Our sole requirement for the moment is  $x_0/\rho t_0^2 \gg 1$ ; hence  $\delta \gg \sqrt{\varepsilon}$ .

Before we proceed to the actual construction of the asymptotic solution, let us set the following definitions and notations: from now on, *and until the end of this section*, let us give the following names to the wave fan and travelling wave solutions to (1.5):

- A selfsimilar solution of (3.1) will be denoted by  $t^{-3/2}(\phi_-, \psi_-)(x/\sqrt{t})$ . Hence, the pair  $(\phi_-, \psi_-)$  is a solution of (3.2).
- We assume that the ignition threshold  $\theta > 1/9$  and  $\rho$  is sufficiently large. Then, according to Theorem 4.1, a traveling wave solution of (1.5) with  $c = 0$  and the temperature that converges to a value  $\phi_0 \in (0, \theta)$  as  $x \rightarrow -\infty$  exists, and will be denoted by  $(\phi_+, \psi_+)(x)$ . Recall that  $(\phi_+, \psi_+)$  is a solution of

$$\begin{aligned} -\phi_+'' + \psi_+\phi_+' &= f(\phi_+) \\ \phi_+(-\infty) &= \phi_0, \quad \phi_+(+\infty) = 1 \\ \psi_+(x) &= \sqrt{2\rho \int_{-\infty}^x (\phi(y) - \phi_0) dy}. \end{aligned} \quad (5.3)$$

Theorem 4.1 also implies that for a sufficiently large  $\rho$  we have  $\phi_0 > 1/9$ , since  $\theta$  satisfies this strict inequality.

- We denote by  $(T^{app}(t, x), u^{app}(t, x))$  the approximate solution that we wish to construct. It will therefore NOT be an exact solution to (1.5); it will satisfy (1.5) up to a small error - typically, of the order  $O(\sqrt{\varepsilon} + \sqrt{\delta})$  over a time interval  $O(\sqrt{\rho^{-1}\varepsilon})$ .

We now have all the ingredients: a set of elementary pieces (wave fan, traveling wave, shock) and two small parameters.

## 5.2 Gluing a wave fan to a combustion wave

Recall that we are not interested at this stage in the transients leading to the development of our composite wave. Hence, we let a pure wave fan of the nonreactive equation (3.1) evolve until we are satisfied with the size of its support and of its derivatives. Then, we translate the profile so that the



temperature at  $x = 0$  is equal (perhaps up to some  $\delta$ -correction) to  $\phi_0$ . This will provide our initial datum to the left. To glue it to a combustion wave at later times, we proceed as follows: we resume the evolution of the wave fan and consider the place where it reaches the value  $\phi_0$  – modulo, once again, a  $\delta$ -correction. If the  $\delta$ -correction is chosen carefully enough, it will be possible to translate the wave profile, then slightly modify the temperature, in such a way that the temperature component of the modified combustion wave matches exactly the wave fan and its slope. The velocity will then be set according to the equation for  $\psi_+$  in (5.3). This will provide a  $\delta$ -approximate solution to (1.5), over a large time interval.

**The reference frame.** Let us therefore consider a solution  $(\phi_-, \psi_-)$  of (3.2). Choose  $\tau_- > 0$  such that

$$\tau_- \geq \delta^{-1} \quad \text{and} \quad \forall \tau \geq \tau_-, \forall x, \quad \frac{1}{\rho\tau^{3/2}}\phi'_-\left(\frac{x}{\sqrt{\tau}}\right) \leq \min(\delta^2, \tau^{-2}). \quad (5.4)$$

This is possible since  $\phi'_- \in L^\infty(\mathbb{R})$ . For  $\tau \geq \tau_-$ , set

$$\tilde{T}_-(\tau, x) := (\rho\tau^{3/2})^{-1}\phi_-(x/\sqrt{\tau}), \quad \tilde{u}_-(\tau, x) = \tau^{-1/2}\psi_-(x/\sqrt{\tau}). \quad (5.5)$$

We consider the places where  $\tilde{T}_-$  reaches values close to  $\phi_0$ . For that obviously we must have  $x/\sqrt{\tau}$  very large. This implies  $T_-(\tau, x) \sim 2x/\rho\tau^2$  – hence, the gluing point  $x \sim \rho\phi_0\tau^2/2$ . We could try to directly consider the moving point  $\tilde{x}_b(\tau) = \rho\phi_0\tau^2/2$ . This is not, however, the most convenient choice. Instead, let us set:

$$\tilde{x}_b(\tau) = \frac{\rho\phi_0\tau^2}{2} + \alpha\tau.$$

The subscript "b" above stands for "back". Let us choose  $\alpha$  according to the strategy that we have proposed above. The function  $\phi_-$  comes with a function  $\psi_-$  accounting for the velocity, given by (3.5)

$$\psi_-(\eta) = \frac{\eta}{2} + \frac{1}{2}\sqrt{\eta^2 + 8 \int_{-\infty}^{\eta} \phi_-(\eta') d\eta'}. \quad (5.6)$$

By the expansion (3.7) of Theorem 3.1, we have, with  $\eta = \tilde{x}_b/\sqrt{\tau}$ :

$$\begin{aligned} \tilde{u}_-(\tau, \tilde{x}_b(\tau)) &= \frac{1}{2\sqrt{\tau}} \left[ \eta + \sqrt{\eta^2 + 8 \left( \eta^2 + \frac{3}{4}a\eta^{4/3} + o(\eta^{4/3}) \right)} \right] = \frac{1}{2\sqrt{\tau}} \left[ 4\eta + a\eta^{1/3} + o(\eta^{1/3}) \right] \\ &= \frac{1}{2\sqrt{\tau}} \left[ \frac{4}{\sqrt{\tau}} \left( \frac{\rho\phi_0\tau^2}{2} + \alpha\tau \right) + \frac{a}{\tau^{1/6}} \left( \frac{\rho\phi_0\tau^2}{2} + \alpha\tau \right)^{1/3} + o(\tau^{1/2}) \right]. \end{aligned}$$

We conclude that

$$\begin{aligned} \tilde{u}_-(\tau, \tilde{x}_b(\tau)) &= \rho\phi_0\tau + 2\alpha + \frac{a}{2} \left( \frac{\rho\phi_0}{2} \right)^{1/3} + o(1) \\ \partial_\tau \tilde{u}_-(\tau, \tilde{x}_b(\tau)) &= \rho\phi_0 + O\left(\frac{1}{\tau}\right). \end{aligned} \quad (5.7)$$

We choose  $\alpha$  to ensure that

$$\tilde{u}_-(\tau, \tilde{x}_b(\tau)) = \dot{\tilde{x}}_b(\tau) + o(1), \quad (5.8)$$

hence  $\alpha = -a(\rho\phi_0/16)^{1/3}$ .

Let us now set the initial time to be  $t = 1$ ; then for all  $t \geq 1$  we choose:

$$x_b(t) = \tilde{x}_b(\tau_- + t) = \frac{\rho\phi_0(\tau_- + t)^2}{2} - a \left( \frac{\rho\phi_0}{16} \right)^{1/3} (\tau_- + t). \quad (5.9)$$

Then, we change the reference frame by setting  $x = x_b(t) + x'$  and drop the prime in order to alleviate the notations: set  $x' := x$ . *This will be our reference frame until the end of this section.* The system (1.5) becomes, in this new reference frame:

$$\begin{aligned} T_t - T_{xx} + (u - \dot{x}_b)T_x - f(T) &= 0 \\ u_t + (u - \dot{x}_b)u_x - \rho T &= 0 \end{aligned} \quad (5.10)$$

**The asymptotic solution on the left.** Let  $\mu$  be a smooth nonnegative function, equal to 1 on the interval  $[2\phi_0/3, 1]$  and equal to 0 on  $[0, \phi_0/2]$ . Our choice for  $(T^{app}, u^{app})$  for  $x < 0$  (in the new moving frame) is:

$$T^-(t, x) = \tilde{T}_-(\tau_- + t, x_b(t) + x) + \gamma(t)\mu\left(\tilde{T}_-(\tau_- + t, x_b(t) + x)\right) \quad (5.11)$$

$$u^-(t, x) = \tilde{u}_-(\tau_- + t, x_b(t) + x) + \dot{x}_b(\tau_- + t). \quad (5.12)$$

The function  $\gamma$  is a correction of the order  $o(1)$ , to be chosen in a more precise fashion below and  $\tilde{T}_-$  is defined in (5.5). The multiplicative correction  $\mu(\tilde{T})$  is non-zero only in the region where we have  $\tilde{T}_-(\tau_- + t, x_b(t) + x) \in (\phi_0/2, \phi_0]$ . We have chosen to multiply the already small term  $\gamma(t)$  by the cut-off  $\mu$  in order to keep the correction of the same order as the main term  $T^-$  for large negative  $x$  where  $T^-$  decays as a Gaussian.

Let us define

$$NL[T, u] = (NL_1, NL_2) := (T_t - T_{xx} + (u - \dot{x}_b)T_x - f(T), u_t + (u - \dot{x}_b)u_x - \rho T).$$

Then, as  $\phi_0 < \theta$ , we have  $f(T_-) = 0$  for  $x < 0$  and  $\delta$  sufficiently small, so that

$$\begin{aligned} NL_1(T^-, u^-) &= T_t^- - T_{xx}^- + (u^- - \dot{x}_b)T_x^- \\ &= \tilde{T}_t^- - \tilde{T}_{xx}^- + \tilde{u}^- \tilde{T}_x^- + \gamma \mu'(\tilde{T})[\tilde{T}_t^- - \tilde{T}_{xx}^- + \tilde{u}^- \tilde{T}_x^-] - \gamma \mu''(\tilde{T}^-)(\tilde{T}_x^-)^2 + \mu(\tilde{T})\dot{\gamma} \\ &= -\gamma \mu''(\tilde{T}^-)(\tilde{T}_x^-)^2 + \mu(\tilde{T})\dot{\gamma} \end{aligned}$$

and

$$NL_2(T^-, u^-) = u_t^- + (u^- - \dot{x}_b)u_x^- - \rho T^- = \tilde{u}_t^- + \tilde{u}^- \tilde{u}_x^- - \rho \tilde{T}^- - \rho \gamma(t)\mu(\tilde{T}^-) = -\rho \tilde{T}^- - \rho \gamma(t)\mu(\tilde{T}^-).$$

We have therefore:

$$\forall x \leq 0, \forall t \geq 1: \quad NL[T, u](x) = (\dot{\gamma}(t)\mu(T) - \gamma \mu''(\tilde{T}^-)(\tilde{T}_x^-)^2, -\rho \gamma(t)\mu(\tilde{T}^-)).$$

Provided that  $\gamma$  is  $o(1)$  as announced, the pair  $(T_-, u_-)$  is then an asymptotic solution on the left.

**The asymptotic solution on the right.** Let us now consider what happens for  $x \in \mathbb{R}_+$ . We seek our solution in the form  $(T^{app}, u^{app})(t, x) = (T_+, u_+)(t, x) = (T_+, u_-(t, 0) + v_+(t, x))$  where  $(T_+, u_+)$  satisfies (5.10) up to a small error, and in addition satisfies the matching conditions

$$\begin{aligned} T_+(t, 0) &= T_-(t, 0) = \phi_0 + \gamma(t) + O((\delta^{-1} + t)^{-1}), \\ \partial_x T_+(t, 0) &= \partial_x T_-(t, 0) = 2(\delta^{-1} + t)^{-2} + o((\delta^{-1} + t)^{-2}) \\ v_+(t, 0) &= 0. \end{aligned} \quad (5.13)$$

Recall that, because of (5.7) and (5.8), the system for  $(T_+, v_+)$  that we wish to satisfy approximately is – we drop the subscripts for convenience:

$$\begin{aligned} T_t - T_{xx} + (v + o(1))T_x &= f(T) \\ v_t + (v + o(1))v_x &= \rho(T - \phi_0 + o(1)). \end{aligned}$$

Hence, it is enough to find a pair  $(T_+, v_+)$  satisfying

$$\begin{aligned}\overline{NL}_1[T, v] &:= T_t - T_{xx} + vT_x - f(T) \\ \overline{NL}_2[T, v] &:= v_t + vv_x - \rho(T - \phi_0),\end{aligned}\tag{5.14}$$

up to an  $o(1)$  error. Now, if  $(\phi_+, \psi_+)$  is a solution of (5.3) normalized so that  $\phi_+(0) = \theta$ , we look for  $T_+$  in the form

$$T_+(t, x) = \phi_+(x - x_+(t)),\tag{5.15}$$

the shift  $x_+(t)$  being adjusted to satisfy the boundary condition for  $\partial_x T_+(t, 0)$  in (5.13). Using representation (4.41) this equation reduces to

$$\phi'_+(-x_+(t)) = 2(\delta^{-1} + t)^{-2} + o((\delta^{-1} + t)^{-2}) = \frac{48}{\sqrt{\rho}(x_+(t) + (16\theta\sqrt{\rho})^{-1/3})^4},$$

which defines a unique  $x_+(t)$  satisfying

$$x_+(t) \sim \left( \frac{24}{\sqrt{\rho}}(t + \delta^{-1}) \right)^{1/2}.\tag{5.16}$$

Let us recall from Section 4 that we have

$$\text{if } \phi_+ \leq \theta, \quad \phi_+ - \phi_0 = \frac{2}{3^{3/4}}\rho^{-1/8}(\phi'_+)^{3/4} := h(\phi'_+).\tag{5.17}$$

Now, the function  $\gamma(t)$  is chosen to satisfy the first equation in (5.13), namely:

$$\gamma(t) = h\left( \frac{48}{\sqrt{\rho}(x_+(t) + (16\theta\sqrt{\rho})^{-1/3})^4} \right) + o(1) = o(1).$$

This fully determines  $T_+(t, x)$ . Now,  $v_+(t, x)$  is just computed as

$$v_+(t, x) = \sqrt{2 \int_0^x (T_+(t, y) - \phi_0) dy}.\tag{5.18}$$

We have  $\overline{NL}[T_+, v_+] = o(1)$ , and thus  $NL[T_+, u_+] = o(1)$ . This ends the construction of the right solution  $(T_+, u_+)$ .

### 5.3 Terminating the combustion wave with a shock

The numerical simulations of Section 2, show that the combustion wave terminates on the right by a hydrodynamic shock, that is, a moving point  $y_f(t)$  across which the unknown  $u$  jumps from a large value  $u_+(t, y_f(t))$  to approximately 0. The temperature profile is slaved to the velocity profile, and undergoes a transition from (approximately) 1 to 0 inside a  $\delta$ -wide layer. Let us recall that at time  $t_0 = \delta^{-1}$  the shock is located at a position  $x_0 = \varepsilon^{-1}$  with the restriction  $\delta \gg \sqrt{\varepsilon}$ . Note that this condition also ensures that the shock is far removed initially from the wave fan to combustion wave transition which has initial width  $x_+ = O(\delta^{-1/2}) \ll \varepsilon^{-1}$  – consequently the shock does not interact with the wave fan at  $t = 1$ .

Let us now find the shock location at later times. From (5.18) we have

$$v_+(t, y_f(t)) = \sqrt{2 \int_0^{y_f(t)} (T_+(t, y) - \phi_0) dy} = \sqrt{2(1 - \phi_0)y_f(t)} + O(1);$$

the quantity  $O(1)$  referring to the time  $t$  and the small parameter  $\varepsilon$ , and coming from the integrability of  $1 - T$  ensured by Lemma 4.6. Assuming that 0 is a good approximation of  $u(t, x)$  to the right of  $y_f(t)$ , the Rankine-Hugoniot condition for the equation for  $u$  yield

$$\dot{y}_f(t) = \frac{1}{2} \left( \sqrt{2 \int_0^{y_f(t)} (T_+(t, y) - \phi_0) dy} - \dot{x}_b(t) \right) = \frac{1}{2} \left[ \sqrt{2(1 - \phi_0)y_f(t)} - \rho\phi_0(\delta^{-1} + t) \right] + O(1). \quad (5.19)$$

This is almost the same equation as (5.1), were it not for the  $O(1)$  term. This is, however, not such a problem: a time shift  $t + \delta^{-1} \rightarrow t$ , change of the unknown  $y_f(t) = \rho t^2 z(t)$ , and of the independent variable  $t = e^\tau$  yields the equation

$$\frac{dz}{d\tau} = \frac{1}{2} q(\sqrt{z}) + O(e^{-2\tau}),$$

which has exactly the same dynamics as (2.14).

The above analysis completes, in principle, the asymptotic analysis, because it describes the dynamics of the two transition layers: the first connects the wave fan to the combustion wave, and the second is the hydrodynamic shock. However, because we wish to pursue our analysis in Section 6 beyond the time of validity of the fan-wave-shock picture, we also have to say something about the temperature profile. The crucial zone is around the shock: in this area we look for an expression for  $T^{app}(t, x)$  in the form

$$T^{app}(t, x) = T_s(t, (\delta^{-1} + t)(x - y_f(t))).$$

Define  $v(t, x)$  as  $v_+(t, x)$  to the left of the shock, that is, for  $x < y_f(t)$ , and as  $-\dot{x}_b(t)$  to the right of the shock, for  $x > y_f(t)$ . The equation to be satisfied by  $T_s(t, y)$  is

$$-\partial_{yy} T_s + \frac{v(t, y_f(t) + (\delta^{-1} + t)^{-1}y) - \dot{y}_f}{\delta^{-1} + t} \partial_y T_s = \frac{f(T_s) - \partial_t T_s - y(\delta^{-1} + t)^{-1} \partial_y T_s}{(\delta^{-1} + t)^2}. \quad (5.20)$$

An approximate equation for  $T_s$  is, therefore,

$$-\partial_{yy} T_s + c(t, y) \partial_y T_s = 0, \quad (5.21)$$

with the function  $c(t, y)$  which is an odd function of  $y$ , discontinuous at  $y = 0$ , and which takes values  $c_-(t)$  for  $y < 0$  and  $c_+(t) = -c_-(t)$  for  $y > 0$ . To the left of the shock, that is, for  $y < 0$ , we have, using expression (5.2) for  $y_f(t)$ :

$$c(t, y) = \frac{\dot{x}_b + \sqrt{2(1 - \phi_0)y_f}}{2(\delta^{-1} + t)} = \frac{1}{2} \left( 2\rho(1 - \phi_0)Q^{-1} \left[ \ln \frac{\delta^{-1} + t}{\sqrt{\rho^{-1}x_0}} \right] \right)^{1/2} + \frac{\rho\phi_0}{2} := c_\delta(t)$$

An important feature of  $c_\delta(t)$  is that we have

$$\dot{c}_\delta(t) = O(\delta^{-1} + t)^{-1}. \quad (5.22)$$

If we additionally impose the conditions  $T_s(t, -\infty) = 1$  and  $T_s(t, +\infty) = 0$ , an expression for  $T_s(t, y)$  is

$$T_s(t, y) = 1 - \frac{c_\delta(t)}{2} \int_{-\infty}^y e^{-c_\delta(t)|z|} dz. \quad (5.23)$$

The final expression for the approximate temperature in the reference frame of the location  $x_b(t)$  of the wave fan is therefore taken as

$$T^{app}(t, x) = \begin{cases} T_-(t, x) & \text{for } x < 0 \\ \inf (T_+(t, x), T_s(t, (\delta^{-1} + t)(x - y_f(t)))) & \text{if } 0 < x < y_f(t) \\ T_s(t, (\delta^{-1} + t)(x - y_f(t))) & \text{if } x > y_f(t) \end{cases} \quad (5.24)$$

and for approximate velocity as

$$u^{app}(t, x) = \begin{cases} u_-(t, x) & \text{for } x < 0 \\ u_-(t, 0) + v_+(t, x) & \text{if } 0 < x < y_f(t) \\ 0 & \text{if } x > y_f(t) \end{cases} \quad (5.25)$$

The function  $T_-(t, x)$  in (5.24) is given by (5.11) and  $T_+(t, x)$  by (5.15), while in (5.25) the function  $u_-(t, x)$  is given by (5.12) and  $v_+(t, x)$  by (5.18).

Since  $\partial_x T_+ > 0$  and  $\partial_x T_s < 0$ , there is only one point where both coincide, and this is a point of discontinuity for  $T_x$ . However, it occurs at a point  $x_d(t)$  where  $T$  is very close to 1. Hence, we know from Lemma 4.6 that the jump in  $T_x$  at  $x_d(t)$  is exponentially small: there exist two positive constants  $k_1$  and  $k_2$  such that:

$$k_1 e^{-k_1 \sqrt{x_d(t)}} \leq e^{(t+\delta^{-1})c_\delta(t)(x_d(t)-y_f(t))} \leq k_2 e^{-k_2 \sqrt{x_d(t)}}.$$

This implies that  $y_f - x_d = O(\sqrt{y_f})$ , thus the jump in  $T_{xx}$  produces a negligible Dirac mass – which one even may regularize by modifying  $T_+$  and  $T_s$  by suitable cut-offs near  $x_d(t)$  where solution is very close to a constant. Therefore, we have

$$T_t^{app} - T_{xx}^{app} + v^{app} T_x^{app} - f(T^{app}) = O((t + \delta^{-1})^{-1}),$$

except in an  $O((t + \delta^{-1})^{-1})$  layer around  $y_f(t)$ , where  $T$  is neither close to 1 nor to 0, and where therefore  $f(T)$  is not close to 0. We have set here  $v^{app} = u^{app} - u_-(t, 0)$ . In the same fashion, we have

$$v_t^{app} + v^{app} v_x^{app} - \rho(T^{app} - \phi_0) = O((t + \delta^{-1})^{-1}),$$

once again except in the same layer where  $T^{app}$  is neither close to 1 nor to 0. However we may write for all  $t \in [0, \sqrt{\rho^{-1}\varepsilon}]$ :

$$\|\overline{NL}(T^{app}(t, \cdot), v^{app}(t, \cdot))\|_{L^1([y_f(t)-1, y_f(t)+1])} = O((t + \delta^{-1})^{-1}). \quad (5.26)$$

Thus we still get an approximate solution albeit not in the pointwise sense. This analysis is valid as long as the transition layers between the wave fan and the combustion wave, and the wave and the rest state are well separated. In the next section we will consider what happens to the "wave fan-combustion wave-shock" solution when the wave starts catching up with the shock.

## 6 The final quenching

The analysis of the previous sections shows that after a long time solution consists of a wave fan on the left, followed by a combustion wave, which in turn ends with a shock. Here we consider such profile as initial data and show that it can quench in a certain regime even when the data is large. We make the following assumptions on the initial data:

**Assumptions on  $T_0$ .** We assume that  $T_0(0) = \phi_0$  – this is the value "in the back of the combustion wave", where transition from the ramp on the left to the wave on the right occurs. To the left of  $x = 0$  the initial data for  $T_0$  looks like a ramp, that is, we assume that

$$0 \leq T_0'(x) \leq \frac{1}{\beta^2} \text{ for all } x \leq 0, \quad (6.1)$$

with some  $\beta \gg \rho$ . The parameter  $\beta$  plays the role of the time it took the original solution to reach the profile that we are now taking as the initial data. The function  $T_0$  looks like a combustion wave,

connecting the values  $T = \phi_0$  on the left and  $T = 1$  on the right, between the points  $x = 0$  and  $x_0 = O(\beta^{\delta_0})$  with  $0 \leq \delta_0 < 1/2$ , where the shock is located, and falls off over a distance  $l_f$  after  $x_0$ :

$$T'_0 > 0 \text{ on } (0, x_0); \quad T'_0 < 0, \text{ on } (x_0, +\infty); \quad T_0(x_0) = 1 - O(e^{-\rho^\alpha}), \quad T_0(x_0 + l_f) \leq \frac{1}{\beta} \quad (6.2)$$

with some  $\alpha > 0$ . We assume that  $l_f \leq C\beta^{\gamma_f}$  with  $\gamma_f < \delta_0 < 1/2$ .

**Assumptions on  $u_0$ .** We assume that to the left of  $x = 0$  the flow profile looks like a ramp and we have  $u'_0(x) \sim O(\rho/\beta)$  for  $x < 0$ , while  $u_0(0) = \beta$ . The function  $u_0(x)$  grows as in the combustion wave between  $x = 0$  and  $y_0 = x_0 + C_0\beta^{-1}$ , where  $u_0(x)$  has its shock, so that

$$|u'_0(x)| \leq C < +\infty, \quad u'_0 > 0 \text{ on } (0, y_0); \quad u_0 = 0 \text{ on } (y_0, +\infty), \quad (6.3)$$

and, moreover

$$u_{max} = u_0(y_0) = \beta + O(\sqrt{\rho y_0}). \quad (6.4)$$

Recall that  $u$  and  $T$  satisfy

$$\begin{aligned} T_t - T_{xx} + uT_x &= f(T) \\ u_t + uu_x &= \rho T. \end{aligned} \quad (6.5)$$

Let us solve the Cauchy problem for (6.5) with the initial data  $(T_0, u_0)$  satisfying the above assumptions. The main result of this section is the following

**Theorem 6.1** *Under the above assumptions on  $T_0$  and  $u_0$ , let  $(T, u)$  be the solution of (6.5) with the Cauchy data  $(T_0, u_0)$ . There exists  $\bar{t} > 0$  such that  $\|T(t, \cdot)\|_\infty \leq \theta$  for  $t \geq \bar{t}$ .*

In particular, assumptions on  $T_0$  and  $u_0$  in Theorem 6.1 are satisfied if we take the approximate solution  $(T^{app}, u^{app})$  constructed in Section 5 at the time  $t_0 = \varepsilon^{-1/2} - \varepsilon^{-1/3}$  and set  $\beta \sim t_0$ .

The strategy is the following: first, we prove that  $u(t, \cdot)$  is well approximated by a time-shift of the solution of the pure Burgers equations, at least for a time much larger than  $1/\beta$ . This property of  $u$  is then exploited in the structure of the equation for  $T$ , which is proved to be quenched in a time of order

$$\bar{t} = K\beta^{\delta_0-1},$$

except in a zone of very small size. Quenching by diffusion is finally proved in this very small zone.

Before starting the construction we change the reference frame: we set  $x' = x - \frac{\rho\phi_0 t^2}{2}$  so that equations become

$$\begin{aligned} T_t - \rho\phi_0 t T_{x'} + u T_{x'} &= T_{x'x'} + f(T) \\ u_t - \rho\phi_0 t u_{x'} + uu_{x'} &= \rho T. \end{aligned}$$

Next, we set

$$v(t, x') = u(t, x') - \rho\phi_0 t \quad (6.6)$$

and drop the primes to get the following equations in the new frame

$$\begin{aligned} T_t + vT_x &= T_{xx} + f(T) \\ v_t + vv_x &= \rho[T - \phi_0]. \end{aligned}$$

The functions  $v$  and  $T$  in the new variables have the same initial values  $u_0(x)$  and  $T_0(x)$ .

**Step 1. An explicit approximation for  $v(t, x)$ .** The maximum principle for entropy solutions of the inviscid Burgers equations implies that  $\underline{v}(t, x) \leq v(t, x) \leq \bar{v}(t, x)$  with the functions  $\underline{v}$  and  $\bar{v}$  that satisfy

$$\underline{v}_t + \frac{1}{2}(\underline{v}^2)_x = -\rho, \quad \bar{v}_t + \frac{1}{2}(\bar{v}^2)_x = \rho, \quad (6.7)$$

with the initial data  $\underline{v}(0, x) = \bar{v}(0, x) = u_0(x)$ . Let us also introduce  $w(t, x)$  which is the entropy solution of the unforced Burgers equation

$$w_t + \frac{1}{2}(w^2)_x = 0, \quad w(0, x) = u_0(x). \quad (6.8)$$

Observe that we have

$$\bar{v}(t, x) = \rho t + w\left(t, x - \frac{\rho t^2}{2}\right), \quad \underline{v}(t, x) = -\rho t + w\left(t, x + \frac{\rho t^2}{2}\right).$$

Therefore, the function  $v(t, x)$  is bounded above and below as follows:

$$-\rho t + w\left(t, x + \frac{\rho t^2}{2}\right) \leq v(t, x) \leq \rho t + w\left(t, x - \frac{\rho t^2}{2}\right),$$

and for small times the problem is essentially reduced to understanding the behavior of  $w(t, x)$ .

As  $u_0(x)$  is smooth and increasing on the interval  $(-\infty, y_0)$  and is equal to zero for  $x > y_0$  the function  $w(t, x)$  remains smooth on an interval  $(-\infty, y_f(t))$  and is equal to zero for  $x > y_f(t)$ , where  $y_f(t)$  is the shock location for the function  $w$  at the time  $t \geq 0$ . The Rankine-Hugoniot condition implies that

$$\dot{y}_f = \frac{1}{2}w(t, y_f^-(t)).$$

We define the characteristics on the left of the shock:

$$\dot{X}(t; x) = w(t, X(t; x)), \quad X(0) = x.$$

The map  $X(t; x)$  is well-defined and increasing both in  $t$  and  $x$  as long as  $x < y_0$  and until the characteristic hits the shock. In addition we have  $w(t, X(t; x)) = u_0(x)$  and therefore

$$X(t; x) = x + tu_0(x).$$

For  $x < y_f(t)$  we may define the inverse map  $y(t; x) = X^{-1}(t, x)$  so that  $x = y + tu_0(y)$ . Now, we can compute almost explicitly the shock location: set  $y_s(t) = X^{-1}(t, y_f(t))$ , then

$$\dot{y}_f = \frac{1}{2}u_0(y_s(t)), \quad y_f(0) = y_0 \quad (6.9)$$

$$y_f(t) = y_s(t) + tu_0(y_s(t)). \quad (6.10)$$

Differentiating (6.10) in time we obtain

$$\dot{y}_s = -\frac{u_0(y_s)}{2(1 + tu_0'(y_s))}, \quad y_s(0) = y_0. \quad (6.11)$$

Note that  $y_s(t) < y_0$  for all  $t > 0$  and  $u_0'(x) > 0$  for  $x < y_0$  so that the solution of (6.11) exists for all  $t > 0$ . Moreover, as  $|u_0'| \leq C$  and  $0 \leq t \leq \bar{t} = K\beta^{\delta_0-1}$  we have

$$-\frac{u_0(y_s)}{2} \leq \dot{y}_s \leq -\frac{u_0(y_s)}{2(1 + C\beta^{\delta_0-1})},$$

so that for  $0 \leq t \leq \bar{t}$  we have  $y_s(t) \geq y_0 - Cu_{max}\beta^{\delta_0-1}$ . It follows that (at the expense of increasing the constant  $C$  in the last inequality below)

$$u_0(y_0) \geq u_0(y_s(t)) \geq u_0\left(y_0 - Cu_{max}\beta^{\delta_0-1}\right) \geq u_0\left(y_0 - C(\beta + C\sqrt{\rho y_0})\beta^{\delta_0-1}\right) \geq u_0\left(y_0 - C\beta^{\delta_0}\right).$$

Now, as  $|u'_0(x)| \leq C\rho/\beta$  for  $x < 0$  we conclude that (again, at the expense of increasing the constant)

$$\beta + C\sqrt{\rho\beta^{\delta_0}} \geq u_0(y_0) \geq u_0(y_s(t)) \geq \beta - \frac{C\rho}{\beta^{1-\delta_0}} \quad \text{for } 0 \leq t \leq \bar{t}.$$

As a consequence, we obtain that

$$\frac{\beta}{2} - \frac{C\rho}{\beta^{1-\delta_0}} \leq \dot{y}_f(t) \leq \frac{\beta}{2} + C\sqrt{\rho\beta^{\delta_0}} \quad \text{for } 0 \leq t \leq \bar{t}. \quad (6.12)$$

To summarize, we have shown that  $v(t, x) \leq \rho t$  for  $x \geq y_f(t) + \rho t^2/2$  and  $v(t, x) \geq w(t, x) - \rho t$  for  $x \leq y_f(t) - \rho t^2/2$  with  $\dot{y}_f$  satisfying (6.12).

Furthermore, for all  $x < y_f(t)$  we have  $w(t, x) = u_0(y(t, x))$  and  $0 \leq x - y(t, x) \leq u_{max}t$ . Therefore,  $w(t, x)$  is large for all  $-\sqrt{\beta} \leq x \leq y_f(t)$  and  $0 \leq t \leq \bar{t}$ :

$$w(t, x) \geq u_0\left(-\sqrt{\beta} - Cu_{max}\beta^{\delta_0-1}\right) \geq u_0\left(-\sqrt{\beta} - C\beta^{\delta_0}\right) \geq \beta - \frac{C\rho}{\beta}\sqrt{\beta} \geq \beta - \frac{C\rho}{\sqrt{\beta}}.$$

We have proved the following lemma.

**Lemma 6.2** *There exists a “shock location” function  $y_f(t)$  satisfying (6.12) with  $y_f(0) = y_0$  such that for any  $K > 0$  there exists  $\beta_0 > 0$  and  $C > 0$  so that for all  $0 \leq t \leq \bar{t} = K\beta^{\delta_0-1}$  we have for  $\beta > \beta_0$ : (i)  $v(t, x) \leq \rho t$  for  $x \geq y_f(t) + \rho t^2/2$ , and (ii)  $v(t, x) \geq \beta - C\rho\beta^{-1/2} - \rho t$  for  $-\sqrt{\beta} \leq x \leq y_f(t) - \rho t^2/2$ .*

**Step 2. A uniform bound for temperature on the left.** Now, in order to establish quenching we consider the coordinate system that moves with the speed  $\dot{y}_f(t)$ : set  $x'' = x - y_f(t)$ . The temperature equation takes the form (we drop the primes):

$$T_t + [v - \dot{y}_f(t)]T_x = T_{xx} + f(T), \quad T(0, x) = T_0(x). \quad (6.13)$$

The first step is to bound the temperature “far on the left”.

**Lemma 6.3** *In the new coordinate system for any  $K > 0$  there exists  $C(K) > 0$  so that we have  $T(t, -\sqrt{\beta}) \leq \phi_0 + C(K)/\sqrt{\beta}$  for  $0 \leq t \leq \bar{t} = K\beta^{\delta_0-1}$ .*

**Proof.** First, note that  $f(T) \leq \Lambda T$  and thus  $T(t, x) \leq \Phi(t, x)e^{\Lambda t}$  with the function  $\Phi$  that satisfies

$$\Phi_t + [v - \dot{y}_f]\Phi_x - \Phi_{xx} = 0, \quad \Phi(0, x) = T_0(x). \quad (6.14)$$

We can write  $T(t, x) = E\{T_0(X(t; x))\}$  where the process  $X(t; x)$  solves

$$dZ = (\dot{y}_f - v)dt + \sqrt{2}dW.$$

However, as  $|v| \leq u_{max} \leq \beta + C\sqrt{\rho y_0}$  and  $|\dot{y}_f(t)| \leq \beta$ , it follows that the probability that  $Z(t)$  starting at  $x = -\sqrt{\beta}$  exits the (very long) interval  $(-3\sqrt{\beta}/2, -\sqrt{\beta}/2)$  before the (very short) time  $K\beta^{\delta_0-1}$  with a sufficiently large  $K > 0$  is smaller than the probability that  $\max_{0 \leq t \leq K\beta^{\delta_0-1}} W(t) \geq K\sqrt{\beta}$ ,



which is exponentially small in  $\beta$  as  $\delta_0 \in (0, 1/2)$ . The claim of Lemma 6.3 now follows since  $T_0(x) \leq \phi_0 + C/\beta^{3/2}$  for all  $x \in (-3\sqrt{\beta}/2, -\sqrt{\beta}/2)$ .  $\square$

**Step 3. Quenching in the middle and on the right.** Lemmas 6.2 and 6.3 are really the piece of information that will lead us to quenching. Clearly, they do not use all the information provided by the construction of our subsolutions and supersolutions for  $v$ ; however they will be sufficient – and it is not obvious that they could have been obtained in a simpler way.

By Lemma 6.3 and the maximum principle, we have  $T(t, x) \leq \phi_0 + C(K)/\sqrt{\beta} \leq \theta$  for the points  $x \in (-\infty, -\beta^{1/2})$ , as long as  $t \leq \bar{t} = K\beta^{\delta_0-1}$ . We want to prove that  $T$  falls under  $\theta$  on the interval  $(-\beta^{1/2}, +\infty)$  at some time  $\tau \leq \bar{t}$ . We will split this interval into three sub-intervals:  $(-\beta^{1/2}, -N/\beta)$ ,  $(-N/\beta, N/\beta)$  and  $(N/\beta, +\infty)$  with a sufficiently large  $N$ .

**1. The interval  $(N/\beta, +\infty)$ .** We use the fact that the advection term in equation (6.13) is less than  $-\beta/4$  for  $x > 1/\beta$  and  $0 \leq t \leq \bar{t}$  by part (i) in Lemma 6.2 since  $\beta^{2\delta_0-2} \ll 1/\beta$  because  $\delta_0 \in (0, 1/2)$ . Let

$$A(t, x) = \frac{1}{\beta} + \exp(-\mu(x + \beta(t - \bar{t})/16 - z_0))$$

with  $z_0 = 1/\beta$  be the exponentially decaying solution of the problem

$$A_t - A_{xx} - \frac{3\beta}{16}A_x = \Lambda \left[ A - \frac{1}{\beta} \right], \quad -\mu^2 + \frac{\beta}{8}\mu = \Lambda,$$

with the constant  $\Lambda$  chosen so that  $\Lambda(s - 1/\beta) \geq f(s)$  for  $s \geq 1/\beta$  – this is possible as  $1/\beta \leq \theta/2$  for a sufficiently large  $\beta$ . The constant  $\mu$  is chosen so that

$$\mu = \frac{\beta}{16} + \frac{1}{2}\sqrt{\frac{\beta^2}{64} - 4\Lambda} \geq \frac{\beta}{16}. \quad (6.15)$$

Note that, since  $A_x(t, x) \leq 0$  and  $v \leq C\rho\beta^{\delta_0-1}$  for  $x > z_0$  and  $0 \leq t \leq \bar{t}$ , we have for  $x \geq z_0$ :

$$A_t - A_{xx} + [v - \dot{y}_f]A_x \geq A_t - A_{xx} - \frac{\beta}{8}A_x = \Lambda(A - \beta^{-1}) \geq f(A). \quad (6.16)$$

Moreover, at the endpoint  $x = z_0$  we have

$$T(t, z_0) \leq 1 \leq \exp(-\mu\beta(t - \bar{t})/16) \leq A(t, z_0) \quad (6.17)$$

for all  $0 \leq t \leq \bar{t}$ . In order to apply the maximum principle we compare the initial data  $T_0(x)$  and  $A(0, x)$ . First, for  $x \geq x_0 - y_0 + l_f$  we have  $T_0(x) \leq 1/\beta \leq A(0, x)$ . Now, at  $x = x_0 - y_0 + l_f$  we have

$$A(0, x_0 - y_0 + l_f) = \frac{1}{\beta} + \exp(-\mu(x_0 - y_0 + l_f - \beta\bar{t}/16 - z_0)) \geq 1,$$

provided that

$$x_0 - y_0 + l_f - \beta\bar{t}/16 - z_0 \leq 0. \quad (6.18)$$

As  $|x_0 - y_0| \leq C/\beta$ ,  $z_0 = 1/\beta$  and  $l_f \leq C\beta^{\gamma_f}$ , the inequality (6.18) indeed holds since  $\gamma_f < \delta_0$ . The function  $A(0, x)$  is decreasing in  $x$  – therefore, we also have

$$T_0(x) \leq 1 \leq A(0, x) \text{ for all } x \leq x_0 - y_0 + l_f.$$

We conclude that  $T_0(x) \leq A(0, x)$  for all  $x \geq z_0$ . The maximum principle together with the inequalities (6.16) and (6.17) implies that  $T(t, x) \leq A(t, x)$  for all  $x \geq z_0$  and all  $0 \leq t \leq \bar{t}$ . Therefore, at the points  $x \geq z_0 + (N - 1)/\beta$  we have at the time  $\bar{t}$ :

$$T(\bar{t}, x) \leq A(\bar{t}, x) \leq A(\bar{t}, z_0 + (N - 1)/\beta) = \frac{1}{\beta} + \exp(-\mu(N - 1)/\beta) \leq \frac{\theta + \phi_0}{2}, \quad (6.19)$$

for a sufficiently large  $N > 0$  since  $\mu \geq \beta/16$ .

**2. The interval**  $(-\sqrt{\beta}, -N/\beta)$ . We set  $z_1 = -1/\beta$  and construct a supersolution for temperature on the interval  $(-\sqrt{\beta}, z_1)$  using part (ii) of Lemma 6.2 in a similar fashion. Take  $\mu$  as in (6.15) and set

$$B(t, x) = e^{\mu(x-\beta(t-\bar{t})/16-z_1)} + \phi_0 + \frac{C}{\sqrt{\beta}}$$

so that

$$B_t - B_{xx} + \frac{3\beta}{16}B_x = \Lambda \left( B - \phi_0 - \frac{C}{\sqrt{\beta}} \right)$$

with  $\Lambda > 0$  chosen so that  $\Lambda(s - \phi_0 - C/\sqrt{\beta}) \geq f(s)$  for  $s \geq \phi_0$  – such  $\Lambda$  exists for a sufficiently large  $\beta > 0$  because  $\phi_0 < \theta$ . Then, as  $B(t, x)$  is increasing in  $x$  for all  $t \geq 0$  and  $v(t, x) \geq 3\beta/4$  for  $x \leq z_1$  and  $0 \leq t \leq \bar{t}$ , we have

$$B_t - B_{xx} + [v - \dot{y}_f]B_x \geq B_t - B_{xx} + \frac{3\beta}{16}B_x = \Lambda \left( B - \phi_0 - \frac{C}{\sqrt{\beta}} \right) \geq f(B) \text{ for } -\sqrt{\beta} \leq x \leq z_1$$

for all  $0 \leq t \leq \bar{t}$ . At the two endpoints:  $x_1 = -\sqrt{\beta}$  and  $x_2 = z_1$  we have  $T(t, x_j) \leq B(t, x_j)$ ,  $j = 1, 2$  for all  $0 \leq t \leq \tau$  simply because  $T(t, x_1) \leq \phi_0 + C/\sqrt{\beta} \leq B(t, x_1)$  according to Lemma 6.3, and  $T(t, x_2) \leq 1 \leq B(\bar{t}, x_2) \leq B(t, x_2)$  since  $B$  is decreasing in  $t$ . Moreover, at the time  $t = 0$  we have

$$B(-y_0, 0) = (1 - \phi_0)e^{\mu(-y_0+1/\beta+\beta\bar{t}/16)} + \phi_0 + \frac{C}{\sqrt{\beta}} > 1,$$

as soon as  $K$  is sufficiently large, since  $y_0 = C\beta^{\delta_0}$  and  $\bar{t} = K\beta^{\delta_0-1}$ . As the function  $B(0, x)$  is increasing in  $x$  it follows that we have  $T(0, x) \leq 1 < B(0, x)$  for all  $x \geq -y_0$ . However, for  $-\sqrt{\beta} \leq x \leq -y_0$  we have  $T(0, x) \leq \phi_0 < B(0, x)$  – we conclude that  $T(0, x) \leq B(0, x)$  for all  $x \geq -\sqrt{\beta}$ . Therefore,  $B(t, x)$  is a supersolution for  $T(t, x)$  and  $T(t, x) \leq B(t, x)$  for all  $0 \leq t \leq \tau$  and all  $x \in (-\sqrt{\beta}, z_1)$ . However, at the time  $t = \bar{t}$  we have then for all  $x \leq -N/\beta$ :

$$\begin{aligned} T(\bar{t}, x) &\leq B(\bar{t}, x) \leq B(\bar{t}, -N/\beta) = (1 - \phi_0)e^{\mu(-N/\beta-\beta\bar{t}/16-z_1+\beta\bar{t}/16)} + \phi_0 + \frac{C}{\sqrt{\beta}} \\ &\leq (1 - \phi_0)e^{\mu(-(N-1)/\beta)} + \phi_0 + \frac{C}{\sqrt{\rho}} \leq \frac{\theta + \phi_0}{2} \end{aligned}$$

since  $\mu \geq \beta/16$  and  $\phi_0 < \theta$ .

We conclude from the above that at the time  $\bar{t}$  the function  $T(t, x)$  is below the value  $\theta$  everywhere except on the interval  $x \in (-N/\beta, N/\beta)$ . A slight generalization of that argument shows that (after increasing  $N$ ) the same statement can be proved for all  $t \in (\bar{t}/2, \bar{t})$ .

**3. The interval**  $(-N/\beta, N/\beta)$ . This is now just quenching by diffusion. It follows from the previous calculations that on the slightly larger interval  $(-2N/\beta, +2N/\beta)$  itself and for any time  $t \in (\bar{t}/2, \bar{t})$  the function  $T$  may be bounded from above as

$$T(t, x) \leq \left[ \frac{\phi_0 + \theta}{2} + \Phi(t, x) \right] e^{\Lambda t}.$$

The function  $\Phi(t, x)$  is the solution of the Dirichlet problem

$$\Phi_t + [v - \dot{y}_f]\Phi_x = \Phi_{xx}, \quad \Phi(t, -2N/\beta) = \Phi(t, +2N/\beta) = 0$$

with the Cauchy data

$$\Phi(\bar{t}/2, x) = \begin{cases} 1, & \text{for } -N/\beta \leq x \leq N/\beta \\ 0, & \text{for } -2N/\beta \leq x < -N/\beta \text{ and } N/\beta < x \leq y_0 + 2N/\beta. \end{cases}$$

As  $|v| + |\dot{y}_f| \leq C\beta$  we have the inequality

$$\bar{\Phi}_t - \bar{\Phi}_{xx} \leq C\beta|\bar{\Phi}_x|.$$

Let now  $\bar{\Phi}$  solve

$$\bar{\Phi}_t - \bar{\Phi}_{xx} = C\beta|\bar{\Phi}_x|, \quad \bar{\Phi}(t, -2N/\beta) = \bar{\Phi}(t, +2N/\beta) = 0$$

with  $\bar{\Phi}(\bar{t}/2, x) = \Phi(\bar{t}/2, x)$ . The maximum principle implies that  $\bar{\Phi}(t, x) \geq \Phi(t, x)$ . However, the function  $\bar{\Phi}$  is symmetric about  $y_0$  and solves the half-interval problem

$$\bar{\Phi}_t - \bar{\Phi}_{xx} = C\beta\bar{\Phi}_x, \quad -2N/\beta < x < 0, \quad \bar{\Phi}(t, -2N/\beta) = \bar{\Phi}_x(t, y_0) = 0.$$

Consider the principal eigenfunction  $\xi(x)$  of this problem with the eigenvalue  $\lambda(\beta)$ :

$$-\xi_{xx} = C\beta\xi_x - \lambda(\beta)\xi, \quad -2N/\beta < x < y_0, \quad \xi(-2N/\beta) = \xi_x(y_0) = 0.$$

After rescaling:  $x = z/\beta$  this becomes

$$-\beta^2\xi_{zz} = C\beta^2\xi_z - \lambda(\beta)\xi, \quad -2 < z < 0, \quad \xi(-2) = \xi_z(0) = 0.$$

and thus  $\lambda(\beta) = -\lambda_0\beta^2$  with  $\lambda_0 > 0$ . It follows that  $\bar{\Phi}(t, x) \leq C_0e^{-\lambda_0\beta^2t}$  and in particular  $\bar{\Phi}(t, x) \leq \theta$  for all  $-2N/\beta \leq x \leq 2N/\beta$  at the time  $\bar{t} = C\delta_0^{\delta_0-1}$ . The proof of Theorem 6.1 is now complete.  $\square$

## References

- [1] M. BELK AND V. VOLPERT, Modeling of heat explosion with convection, *Chaos* **14** (2004), pp. 263–273.
- [2] M. BELK, B. KAZMIERCZAK, V. VOLPERT Existence of reaction-diffusion-convection waves in unbounded cylinders, *Int. Jour. Math. Math. Sci.*, **2005:2** (2005), pp. 169–193.
- [3] H. BERESTYCKI, B. NICOLAENKO, B. SCHEURER, Travelling waves solutions to combustion models and their singular limits, *SIAM J. Math. Anal.* **16**, 6 (1985), pp. 1207–1242.
- [4] H. BERESTYCKI, P. CONSTANTIN, L. RYZHIK, Non-planar fronts in Boussinesq reactive flows, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **23** (2006), pp. 407–437.
- [5] P. CONSTANTIN, A. KISELEV, L. RYZHIK, Quenching of flames by fluid advection, *Comm. Pure Appl. Math.* **54** (2001), pp. 1320–1342.
- [6] P. CONSTANTIN, A. KISELEV AND L. RYZHIK, Fronts in reactive convection: bounds, stability and instability, *Comm. Pure Appl. Math.*, **56** (2003), pp. 1781–1803.
- [7] P. CONSTANTIN, M. LEWICKA AND L. RYZHIK, Traveling waves in 2D reactive Boussinesq systems with no-slip boundary conditions, *Nonlinearity*, **19** (2006), pp. 2605–2615.
- [8] P. CONSTANTIN, A. NOVIKOV AND L. RYZHIK, Relaxation in reactive flows, to appear in *Geom. Funct. Anal.*, 2007.
- [9] A. DE WIT, Fingering of chemical fronts in porous media, *Phys. Rev. Lett.*, **87**, 054502.
- [10] A. DE WIT AND G.M. HOMSY, Viscous fingering in reaction-diffusion systems, *J. Chem. Phys.*, **110** (1999), pp. 8663–8675.

- [11] A. FANNJIANG, A. KISELEV, L. RYZHIK, Quenching of reaction by cellular flows, *Geom. Funct. Anal.* **16** (2006), pp. 40–69.
- [12] L. KAGAN AND G. SIVASHINSKY, Flame propagation and extinction in large-scale vortical flows, *Combust. Flame*, ]**120** (2000), pp. 222–232.
- [13] L. KAGAN, P.D. RONNEY AND G. SIVASHINSKY, Activation energy effect on flame propagation in large-scale vortical flows. *Combust. Theory Modelling*, **6** (2002), pp. 479–488.
- [14] L. KAGAN AND G. SIVASHINSKY Effect of Lewis number on flame propagation through vortical flows. *Combust. Flame*, **142** (2005), pp. 235–240.
- [15] YA.I. KANEL', Certain problems of burning-theory equations, *Sov. Math. Dokl.* **2** (1961), pp 48-51.
- [16] YA.I. KANEL', Stabilization of solution of the Cauchy problem for equations encountered in combustion theory, *Mat. Sbornik* **59** (1962), pp. 245-288.
- [17] A. KISELEV, A. ZLATOŠ, Quenching of combustion by shear flows, *Duke Math. J.* **132** (2006), pp. 49–72.
- [18] M. LEWICKA, Existence of travelling waves in the Stokes-Boussinesq system for reactive flow, *Jour. Diff. Eq.*, **237** (2007), pp. 343–371.
- [19] R. TEXIER-PICARD AND V. VOLPERT, Problèmes de réaction-diffusion-convection dans des cylindres non bornés, *C. R. Acad. Sci. Paris Sér. I Math.* **333** (2001), pp. 1077-1082
- [20] R. TEXIER-PICARD AND V. VOLPERT, Reaction-diffusion-convection problems in unbounded cylinders, *Revista Matematica Complutense*, **16** (2003), pp. 233–276.
- [21] N. VLADIMIROVA, P. CONSTANTIN, A. KISELEV, O. RUCHAYSKIY, L. RYZHIK, Flame enhancement and quenching in fluid flows, *Combust. Theory Model.* **7** (2003), pp. 487–508.
- [22] N. VLADIMIROVA, R. ROSNER, Model flames in the Boussinesq limit: the effects of feedback, *Phys. Rev. E.*, **67** (2003), pp. 066305.
- [23] N. VLADIMIROVA AND R. ROSNER, Model Flames in the Boussinesq Limit: The Case of Pulsating Fronts, *Phys. Rev. E*, **71** (2005), pp. 067303
- [24] A. ZLATOŠ, Sharp transition between extinction and propagation of reaction. *J. Amer. Math. Soc.* **19** (2006), pp. 251–263