

Global Offensive Alliances in Graphs via Degree Sequences

Ararat Harutyunyan^{1,2}

*Department of Mathematics
Simon Fraser University
Vancouver, Canada*

Abstract

A global offensive alliance in a graph $G = (V, E)$ is a subset S of V such that for every vertex v not in S at least half of the vertices in the closed neighborhood of v are in S . We give an upper bound on the global offensive alliance number of a graph in terms of its degree sequence. We also study global offensive alliances of random graphs.

Keywords: Global offensive alliances, degree sequence, upper bounds, random graph.

1 Introduction

The study of alliances in graphs was first introduced by Hedetniemi, Hedetniemi and Kristiansen [7]. They introduced the concepts of defensive and offensive alliances, global offensive and global defensive alliances and studied alliance numbers of a class of graphs such as cycles, wheels, grids and complete graphs. Haynes et al. [5] studied the global defensive alliance numbers of different classes of graphs. They gave lower bounds for general graphs, bipartite graphs and trees, and upper bounds for general graphs and trees. Rodriquez-Velazquez and Sigarreta [12] studied the defensive alliance number

¹ Research supported by FQRNT (Le Fonds québécois de la recherche sur la nature et les technologies) doctoral scholarship.

² Email: aha43@sfu.ca

and the global defensive alliance number of line graphs. A characterization of trees with equal domination and global strong defensive alliance numbers was given by Haynes, Hedetniemi and Henning [6].

Offensive alliances were first studied by Favaron et. al [3], where they derived some bounds on the offensive alliance number. Rodriguez-Velazquez and Sigarreta [9] gave bounds for offensive and global offensive alliance numbers in terms of the algebraic connectivity, the spectral radius, and the Laplacian spectral radius of a graph. They also gave bounds on the global offensive alliance number of cubic graphs in [10] and the global offensive alliance number for general graphs in [11]. Some bounds on the global offensive alliances were given in [4]. Balakrishnan et al. [2] studied the complexity of global alliances. They showed that the decision problems for global defensive and global offensive alliances are both NP-complete for general graphs.

This paper further studies the global offensive alliance number of a graph.

Given a simple graph $G = (V, E)$ and a vertex $v \in V$, the *open neighborhood* of v , $N(v)$, is defined as $N(v) = \{u : uv \in E\}$. The *closed neighborhood* of v , denoted by $N[v]$, is $N[v] = N(v) \cup \{v\}$. Given a set $X \subset V$, the *boundary* of X , denoted by $\delta(X)$, is the set of vertices in $V - X$ that are adjacent to at least one member of X . A set $X \subset V$ is called a *dominating set* if $\delta(X) = V - X$.

Definition 1.1 A set $S \subset V$ is an *offensive alliance* if for every $v \in \delta(S)$, $|N[v] \cap S| \geq |N[v] - S|$. An offensive alliance S is called a *global offensive alliance* if S is also a dominating set.

Definition 1.2 A global offensive alliance S is called a *global strong offensive alliance* if for every $v \in \delta(S)$, $|N[v] \cap S| > |N[v] - S|$.

Definition 1.3 The *global (strong) offensive alliance number* of G is the cardinality of a minimum size global (strong) offensive alliance in G , and is denoted by $\gamma_o(G)$ ($\gamma_s(G)$). A minimum size global offensive alliance is called a $\gamma_o(G)$ -set.

In this paper, we study the global (strong) offensive alliance numbers of general graphs. We give an upper bound on the global (strong) offensive alliance number of general graphs. Additionally, we study the global (strong) offensive alliance number of random graphs.

The rest of the paper is organized as follows. In Section 2, we give an upper bound on the global (strong) offensive alliance number of a general graph in terms of its order and degree sequence. Using this bound, we obtain a second upper bound on the global (strong) offensive alliance number in terms of the

minimum degree of the graph. In Section 3, we study the global (strong) offensive alliance number of the random graph $G(n, 1/2)$.

2 Global Offensive Alliances in Graphs

In this section we give an upper bound on $\gamma_o(G)$ for any graph G . Our result derives an upper bound on $\gamma_o(G)$ in terms of the degree sequence of the graph G . The method of the proof is probabilistic. All the required probabilistic tools can be found in [1]. Note that $\exp(x)$ is the exponential function e^x .

Theorem 2.1 *Let $G = (V, E)$ be a graph of order n . Let $\deg(v)$ denote the degree of vertex v . Then for all $1/2 > \alpha > 0$,*

$$\gamma_o(G) \leq \left(\frac{1}{2} + \alpha\right)n + \left(\frac{1}{2} - \alpha\right) \sum_{v \in V} \exp\left(-\frac{\alpha^2}{1 + 2\alpha} \cdot \deg(v)\right)$$

Proof. We put every vertex $v \in V$ in a set S with probability p , independently. The value of p will be determined later. The random set S is going to be part of the global offensive alliance. For every vertex $v \in V$, let X_v denote the number of vertices in the neighborhood of v that are in S . Let $Y = \{v \in V : v \notin S \text{ and } X_v \leq \lfloor \frac{\deg(v)}{2} \rfloor\}$. Clearly, $S \cup Y$ is a global offensive alliance. Note that $\mathbb{E}[|S|] = np$. Now, we estimate $\mathbb{E}[|Y|]$.

It is not hard to see that X_v is a Binomial($\deg(v), p$) random variable. We use the Chernoff Bound (see, for example, Alon and Spencer [1]) to bound $\mathbb{P}[X_v \leq \frac{\deg(v)}{2}]$. The Chernoff Bound states that for any positive $\epsilon < 1$ and random variable X that has binomial distribution with probability p and mean pn ,

$$(1) \quad \mathbb{P}[|X - pn| > \epsilon pn] < 2e^{-\epsilon^2 pn/2}.$$

Let $\epsilon = 1 - \frac{1}{2p}$. Then, by the Chernoff Bound,

$$\begin{aligned} \mathbb{P}[X_v \leq \frac{\deg(v)}{2}] &= \mathbb{P}[X_v \leq (1 - \epsilon)p(\deg(v))] \\ &< e^{-\epsilon^2 \deg(v)p/2} \\ &= e^{-(1 - \frac{1}{2p})^2 \deg(v)p/2}. \end{aligned}$$

Chernoff's bound holds whenever $0 < \epsilon < 1$, or equivalently when $p > \frac{1}{2}$.

Now,

$$\mathbb{P}[v \in Y] = \mathbb{P}[\{v \notin S\} \cap \{X_v \leq \deg(v)/2\}]$$

$$\begin{aligned}
&= \mathbb{P}[v \notin S] \mathbb{P}[X_v \leq \deg(v)/2] \\
&\leq (1-p)e^{-(1-\frac{1}{2p})^2 \deg(v)p/2},
\end{aligned}$$

by independence. By linearity of expectation, we get that

$$\mathbb{E}[|Y|] \leq \sum_{v \in V} (1-p)e^{-(1-\frac{1}{2p})^2 \deg(v)p/2}.$$

Now, we have that

$$(2) \quad \mathbb{E}[|S \cup Y|] \leq np + \sum_{v \in V} (1-p)e^{-(1-\frac{1}{2p})^2 \deg(v)p/2}.$$

Therefore, there exists a global offensive alliance in G of size at most

$$(3) \quad np + \sum_{v \in V} (1-p)e^{-(1-\frac{1}{2p})^2 \deg(v)p/2}.$$

Thus, we have that

$$(4) \quad \gamma_o(G) \leq np + \sum_{v \in V} (1-p)e^{-(1-\frac{1}{2p})^2 \deg(v)p/2}.$$

The only constraint we have on p is that $p > \frac{1}{2}$. We set $p = \frac{1}{2} + \alpha$ for any $\alpha > 0$. This completes the proof. \square

A similar result can be derived for the global strong offensive alliance number of a graph.

Theorem 2.2 *Let $G = (V, E)$ be a graph of order n . Then for all $1/2 > \alpha > 0$,*

$$\gamma_o(G) \leq \left(\frac{1}{2} + \alpha\right)n + \sum_{v \in V} \exp\left(-\frac{\alpha^2}{1+2\alpha} \cdot (\deg(v) + 1)\right).$$

Due to space restrictions, the proof of Theorem 2.2 is omitted. The proof is in the same spirit as the proof of Theorem 2.1. Theorems 2.1 and 2.2 yield the following corollaries.

Corollary 2.3 *Let G be a graph of minimum degree $d \geq 2$. Then*

$$\gamma_o(G) \leq \left(\frac{1}{2} + \left(\frac{\log d}{d}\right)^{1/2} + \frac{1}{2\sqrt{d}} - \frac{\sqrt{\log d}}{d}\right)n.$$

Corollary 2.4 *Let G be a graph of minimum degree $d \geq 2$. Then*

$$\gamma_{\delta}(G) \leq \left(\frac{1}{2} + \left(\frac{\log d}{d+1} \right)^{1/2} + \frac{1}{\sqrt{d}} \right) n.$$

Note that if the minimum degree d of a graph G tends to infinity, corollaries 2.3 and 2.4 imply that $\gamma_o(G)$ and $\gamma_{\delta}(G)$ approach to $n/2$. For large minimum degree d , our results improve the following sharp bounds found in [8].

Theorem 2.5 ([8]) *For every connected graph G of order $n \geq 2$, $\gamma_o(G) \leq \frac{2n}{3}$. If the minimum degree of G is at least 2, $\gamma_{\delta}(G) \leq \frac{5n}{6}$.*

3 Global Offensive Alliances in Random Graphs

The *random graph* $G(n, 1/2)$ is the graph on n vertices where each possible edge is present with probability $1/2$, independently. It seems plausible that the random graph $G(n, 1/2)$ should have a global offensive alliance number of approximately $n/2$. In this section, we provide some evidence that this is the case. The main result of this section is the following theorem. The proof is long and is omitted.

Theorem 3.1 *Let $c < 1/2$ be any fixed constant. Then*

$$\mathbb{P}[\gamma_o(G(n, 1/2)) \leq cn] = o(1).$$

Since a global strong offensive alliance in a graph G is also a global offensive alliance in G , Theorem 3.1 immediately implies the following.

Theorem 3.2 *Let $c < 1/2$ be any fixed constant. Then*

$$\mathbb{P}[\gamma_{\delta}(G(n, 1/2)) \leq cn] = o(1).$$

On the other hand, using corollaries 2.3 and 2.4, one can obtain that for $c > 1/2$, the random graph has a global (strong) offensive alliance of size at most cn almost surely.

Theorem 3.3 *Let $c > 1/2$ be any fixed constant. Then*

$$\mathbb{P}[\gamma_{\delta}(G(n, 1/2)) \leq cn] = 1 - o(1).$$

Theorem 3.4 *Let $c > 1/2$ be any fixed constant. Then*

$$\mathbb{P}[\gamma_o(G(n, 1/2)) \leq cn] = 1 - o(1).$$

References

- [1] N. Alon, J. Spencer, “The Probabilistic Method,” 2nd Ed., Wiley, 1992.
- [2] H. Balakrishnan, A. Cami, N. Deo, and R. D. Dutton, *On the complexity of finding optimal global alliances*, J. Combinatorial Mathematics and Combinatorial Computing, Volume 58 (2006), 23–31.
- [3] O. Favaron, G. Fricke, W. Goddard, S. M. Hedetniemi, S. T. Hedetniemi, P. Kristiansen, R. C. Laskar and D. R. Skaggs, *Offensive alliances in graphs*, Discuss. Math. Graph Theory 24 (2)(2004), 263–275.
- [4] A. Harutyunyan, *Some bounds on alliances in trees*, 9th Cologne-Twente Workshop on Graphs and Combinatorial Optimization (CTW 2010), 83–86.
- [5] T. W. Haynes, S. T. Hedetniemi, and M. A. Henning, *Global defensive alliances in graphs*, Electronic Journal of Combinatorics 10 (2003), no. 1, R47.
- [6] T. W. Haynes, S. T. Hedetniemi, and M. A. Henning, *A characterization of trees with equal domination and global strong alliance numbers*, Utilitas Mathematica, Volume 66(2004), 105–119.
- [7] S. M. Hedetniemi, S. T. Hedetniemi, and P. Kristiansen, *Alliances in graphs*, Journal of Combinatorial Mathematics and Combinatorial Computing, Volume 48 (2004), 157–177.
- [8] J. A. Rodriguez-Velazquez, J.M. Sigarreta, *On the global offensive alliance number of a graph*, Discrete Applied Mathematics, Volume 157 (2006), no.2, 219–226.
- [9] J. A. Rodriguez-Velazquez, J. M. Sigarreta, *Spectral study of alliances in graphs*, Discussiones Mathematicae Graph Theory 27 (1) (2007) 143–157.
- [10] J. A. Rodriguez-Velazquez and J. M. Sigarreta, *Offensive alliances in cubic graphs*, International Mathematical Forum Volume 1 (2006), no. 36, 1773–1782.
- [11] J. A. Rodriguez-Velazquez, J.M. Sigarreta, *Global Offensive Alliances in Graphs*, Electronic Notes in Discrete Mathematics, Volume 25 (2006), 157–164.
- [12] J. A. Rodriguez-Velazquez, J. M. Sigarreta, *On defensive alliances and line graphs*, Applied Mathematics Letters, Volume 19 (12) (2006) 1345–1350.