

BÖTTCHER COORDINATES

XAVIER BUFF, ADAM L. EPSTEIN, AND SARAH KOCH

ABSTRACT. A well-known theorem of Böttcher asserts that an analytic germ $f : (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$ which has a superattracting fixed point at 0, more precisely of the form $f(z) = az^k + o(z^k)$ for some $a \in \mathbb{C}^*$, is analytically conjugate to $z \mapsto az^k$ by an analytic germ $\phi : (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$ which is tangent to the identity at 0. In this article, we generalize this result to analytic maps of several complex variables.

INTRODUCTION

An analytic germ $f : (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$ with a superattracting fixed point at 0 can be written in the form

$$f(z) = az^k + \mathcal{O}(z^{k+1}), \quad a \neq 0, \quad k \geq 2.$$

In 1904, L. E. Böttcher proved the following theorem.

Theorem (Böttcher). *If $f : (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$ has a superattracting point at 0 as above, there exists a germ of analytic map $\phi : (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$, which is tangent to the identity at 0 and conjugates f to the map $h : w \mapsto aw^k$ in some neighborhood of 0, i.e., $\phi \circ f = h \circ \phi$.*

The germ ϕ is called a *Böttcher coordinate* for the germ f .

Böttcher coordinates have been an essential tool in the study of complex dynamics in one variable; a Böttcher coordinate gives polar coordinates near the associated superattracting fixed point, which are compatible with the dynamics of f . In the case of the superattracting fixed point at infinity for a polynomial, the angular coordinate is called the *external angle* and its level curves are called *external rays*. The importance of external angles and external rays was emphasized in [DH].

It is quite natural to ask whether there is an analogue of this theorem in higher dimensions; that is, given an analytic germ $F : (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^m, 0)$ is there an analytic germ $\Phi : (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^m, 0)$ which conjugates F to its terms of lowest degree? According to Hubbard and Papadopol in [HP], “the map is not in general locally conjugate, even topologically, to its terms of lowest degree; the local geometry near such a point is much too rich for anything like that to be true.” Hubbard and Papadopol present the following example to illustrate their point.

Example 1. Consider the map

$$F : \mathbb{C}^2 \rightarrow \mathbb{C}^2 \text{ given by } (x, y) \mapsto (x^2 + y^3, y^2).$$

Let $H : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ be the map $(x, y) \mapsto (x^2, y^2)$. There is no analytic conjugacy between F and H in a neighborhood of the origin because the dynamics of the

The research of the first author was supported in part by the grant ANR-08-JCJC-0002 and the IUF.

The research of the third author was supported in part by the NSF.




FIGURE 1. For polynomials of one complex variable, the Böttcher coordinate of ∞ extends throughout the entire basin if the filled Julia set is connected. The checkerboard pattern in the pictures above highlights the external rays in each picture. On the right is the filled Julia set of the model map, $z \mapsto z^2$, and on the left is the filled Julia set for a quadratic polynomial.

maps are incompatible. Indeed, the critical locus of F consists of the the two axes, which is also the critical locus of H . But the map H fixes the two axes, whereas F fixes $y = 0$ and maps $x = 0$ to the curve $t \mapsto (t^3, t^2)$.

However, there is another explanation which is more relevant to our discussion. The critical value locus of F contains the curve $t \mapsto (t^3, t^2)$ which has a cusp; any smooth conjugacy would have to map this singular curve to a component of the critical value locus of H ; however, the critical value locus of H consists of $x = 0$ and $y = 0$, which are smooth. So certainly no analytic conjugacy exists.

Remark 1. In the example above, the map H is *homogeneous*. A homogeneous map

$$H : \mathbb{C}^{m+1} \rightarrow \mathbb{C}^{m+1}$$

descends to a map on projective space $h : \mathbb{P}^m \dashrightarrow \mathbb{P}^m$; the critical locus and the postcritical locus of H will be cones over the critical locus and postcritical locus of h .

Therefore, if F is going to be locally conjugate to a homogeneous map in a neighborhood of 0, each component of the critical locus and postcritical locus of F containing the superattracting fixed point, must be an *analytic cone*; that is, each of these components must be the image of a cone under an analytic isomorphism. This is a rather strong condition; the homogeneous maps certainly satisfy it, and we will see in section 3 that there are other families of maps which satisfy this criterion.

The following people have studied the dynamics of maps with superattracting fixed points in higher dimensions: J.H. Hubbard and P. Papadopol develop a very important theory of Green functions for maps with superattracting fixed points in \mathbb{C}^m (see [HP]), C. Favre and M. Jonsson classify *contracting rigid germs* of $(\mathbb{C}^2, 0)$ in [FJ]. Both S. Ushiki and T. Ueda have results about a Böttcher theorem in

higher dimensions: Ushiki presents a local Böttcher theorem in \mathbb{C}^2 for maps of a special form in [Us], and Ueda presents both local and global Böttcher theorems in \mathbb{C}^m for a particular family of maps in [Ue1].

In this article, we begin with an analytic germ $F : (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^m, 0)$, which has an *adapted superattracting fixed point* at 0. We present necessary and sufficient conditions for such an F to be locally conjugate to its *quasihomogeneous part* at 0. Our criteria are stated in terms of *admissible vector fields*; these vector fields detect the shape of the postcritical locus of our map F to see if the analytic cone condition in remark 1 is satisfied. The relevant definitions are given in Section 1.1.

Theorem 1 (Local Böttcher Coordinates). *Let $F : (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^m, 0)$ be a germ of an analytic map having an adapted superattracting fixed point at $0 \in \mathbb{C}^m$. Let $H : \mathbb{C}^m \rightarrow \mathbb{C}^m$ be the quasihomogeneous part of F at 0 with multidegree (k_1, \dots, k_p) . Then the following are equivalent.*

- (1) *There is a germ of an analytic map $\Phi : (\mathbb{C}^m, 0) \rightarrow (\mathbb{C}^m, 0)$ such that*

$$\Phi(0) = 0, \quad D_0\Phi = \text{id} \quad \text{and} \quad \Phi \circ F = H \circ \Phi.$$

- (2) *There is an admissible p -tuple of germs of vector fields (ξ_1, \dots, ξ_p) such that*

$$DF \circ \xi_j = k_j \cdot \xi_j \circ F$$

near 0 for all $j \in [1, p]$.

- (3) *There is an admissible p -tuple of germs of vector fields $(\zeta_1, \dots, \zeta_p)$ such that ζ_j is tangent to the germ of the postcritical set of F for all $j \in [1, p]$.*

The proof of theorem 1 is a beautiful confluence of classical results of Euler, Poincaré, Cartan, and Arnol'd with the theory of Green functions developed in [HP]. Sections 1.2–1.6 are devoted to presenting these results and adapting them to our setting.

In section 2, we present a result about extending a local Böttcher coordinate Φ to a larger domain. More precisely let Ω be a complex analytic manifold, and $F : \Omega \rightarrow \Omega$ be an analytic map with a superattracting fixed point at $a \in \Omega$. The basin of attraction of a for F is the open set of points whose orbits converge to a . The immediate basin is the connected component of the basin containing a ; we denote the immediate basin as $\mathcal{B}_a(F)$.

Theorem 2 (Global Böttcher coordinates). *Let Ω be a complex analytic manifold, $F : \Omega \rightarrow \Omega$ be a proper analytic map with a superattracting fixed point at $a \in \Omega$ and $H : T_a\Omega \rightarrow T_a\Omega$ be a quasihomogeneous map. Suppose that*

- *there is a local isomorphism $\Phi : (\Omega, a) \rightarrow (T_a\Omega, 0)$ with $\Phi \circ F = H \circ \Phi$;*
- *near a , Φ maps the postcritical set of $F : \mathcal{B}_a(F) \rightarrow \mathcal{B}_a(F)$ to the postcritical set of $H : \mathcal{B}_0(H) \rightarrow \mathcal{B}_0(H)$.*

Then, Φ extends to a global isomorphism $\Phi : \mathcal{B}_a(F) \rightarrow \mathcal{B}_0(H)$ conjugating F to H :

$$\begin{array}{ccc} \mathcal{B}_a(F) & \xrightarrow{\Phi} & \mathcal{B}_0(H) \\ \downarrow F & & \downarrow H \\ \mathcal{B}_a(F) & \xrightarrow{\Phi} & \mathcal{B}_0(H). \end{array}$$

In section 3, we apply our results to a large class of new examples of *postcritically finite endomorphisms of \mathbb{P}^n* . And finally, in section 4, we conclude with some remaining questions.

Acknowledgements. We would like to thank our colleagues for their constant support, and stimulating discussions. In particular, we thank John H. Hubbard, Curtis T. McMullen, John Milnor, and Saeed Zakeri.

1. THE LOCAL RESULT

1.1. **Set Up.** In section 1.1, we will use the following notation.

- E is a \mathbb{C} -linear space of dimension $m \geq 1$.
- $\langle \cdot | \cdot \rangle$ is a Hermitian product on E .
- $\| \cdot \|$ is the associated norm.
- $E = E_1 \oplus \cdots \oplus E_p$ is the direct sum of $p \geq 1$ linear spaces.
- $(\pi_j : E \rightarrow E_j)_{j \in [1, p]}$ are the projections associated to the direct sum.
- for $v \in E$, we denote $v_j := \pi_j(v) \in E_j$.
- For $F : (E, 0) \rightarrow (E, 0)$ an analytic germ, we denote $F_j := \pi_j \circ F : E \rightarrow E_j$.
- if $(H_j : E_j \rightarrow E_j)_{j \in [1, p]}$ are maps, then $H_1 \oplus \cdots \oplus H_p$ is the map

$$E \ni v_1 + \cdots + v_p \mapsto H_1(v_1) + \cdots + H_p(v_p) \in E.$$

- $(\pi_j : E \rightarrow E_j)_{j \in [1, p]}$ are the projections associated to the direct sum.
- k_1, \dots, k_p are integers greater than or equal to 2.

The word *adapted* in the definition 1 refers to this data.

Recall that a map $H : L \rightarrow L$ on a \mathbb{C} -linear space L is homogeneous of degree $k \geq 1$ if

$$\forall v \in L, \quad H(v) = \phi(\underbrace{v, \dots, v}_{k \text{ times}})$$

for some k -linear map $\phi : L^k \rightarrow L$. Equivalently, H is analytic and

$$\forall \lambda \in \mathbb{C}, \forall v \in L, \quad H(\lambda v) = \lambda^k H(v).$$

The homogeneous map H is nondegenerate if $H^{-1}\{0\} = \{0\}$.

Definition 1. *An analytic germ $F : (E, 0) \rightarrow (E, 0)$ has an adapted superattracting fixed point if for all $j \in [1, p]$, there is a nondegenerate homogeneous map $H_j : E_j \rightarrow E_j$ of degree $k_j \geq 2$ such that*

$$F_j(v) \underset{v \rightarrow 0}{\simeq} H_j \circ (v_j) + o(\|(v_j)\|^{k_j}).$$

The map $H_1 \oplus \cdots \oplus H_p$ is the quasihomogeneous part of F at 0 and (k_1, \dots, k_p) is its multidegree.

Note that the derivative of a map at an adapted superattracting fixed point necessarily vanishes. The particular form of F_j implies the following invariance property.

Proposition 1. *The spaces $E_j^\top := \{v_j = 0\}$ and E_j are locally totally invariant under F ; that is, there exists a neighborhood V of 0 such that*

$$F^{-1}(E_j^\top) \cap V = E_j^\top \cap V \quad \text{and} \quad F^{-1}(E_j) \cap V = E_j \cap V.$$

Proof. Let us first prove that E_j^\top is locally totally invariant. First, if $v_j = 0$, then $F_j(v) = 0$. This shows that near 0, $E_j^\top \subseteq F^{-1}(E_j^\top)$. So if E_j^\top were not locally totally invariant, we could find an analytic germ

$$\gamma : (\mathbb{C}, 0) \rightarrow (F^{-1}(E_j^\top), 0) \quad \text{with} \quad \gamma_j := \pi_j \circ \gamma \neq 0.$$

The order of vanishing m of γ_j at 0 would be finite. Since H_j is nondegenerate with degree k_j , the order of vanishing of $H_j \circ \gamma_j$ would be $k_j \cdot m$. So the order of vanishing of $F_j \circ \gamma = H_j \circ \gamma_j + o(\|\gamma_j\|^{k_j})$ would be $k_j \cdot m < \infty$ which is a contradiction since by assumption $F_j \circ \gamma \equiv 0$.

Next, note that

$$E_j = \bigcap_{i \neq j} E_i^\top.$$

Since the spaces E_i^\top are locally totally invariant, it follows that the space E_j is locally totally invariant. \square

We want to give necessary and sufficient conditions for the existence of an isomorphism $\Phi : (E, 0) \rightarrow (E, 0)$ conjugating F to its quasihomogeneous part H , *i.e.*, such that $\Phi \circ F = H \circ \Phi$. Such an isomorphism is called a Böttcher coordinate for F .

For $v \in E$, there is a canonical isomorphism between the \mathbb{C} -linear space E and the tangent space $T_v E$. Together, these isomorphisms induce a canonical bundle isomorphism between TE and $E \times E$. If $(v, w) \in E \times E$, we shall denote by $(v; w)$ the corresponding tangent vector in $T_v E$. If $F : U \subseteq E \rightarrow E$ is an analytic map, we denote $D_v F : T_v E \rightarrow T_{F(v)} E$ the derivative of F at $v \in U$. We denote by $DF : TU \rightarrow TX$ the bundle map $(v; w) \mapsto D_v F(v; w)$. We shall denote by $F'(v; w)$ the vector in E corresponding to $DF(v; w) \in T_{F(v)} E$.

Definition 2. Let ϑ_{rad} and $\vartheta_1, \dots, \vartheta_p$ be the linear vector fields $E \rightarrow TE$ defined by

$$\forall v \in E, \quad \vartheta_{\text{rad}}(v) := (v; v) \quad \text{and} \quad \vartheta_j(v) := (v; v_j).$$

Note that $\vartheta_{\text{rad}} = \vartheta_1 + \dots + \vartheta_p$.

Definition 3. A vector field ξ is asymptotically radial if ξ is defined and analytic near $0 \in E$ with $\xi(v) \underset{v \rightarrow 0}{=} \vartheta_{\text{rad}}(v) + o(\|v\|)$.

A p -tuple of vector fields (ξ_1, \dots, ξ_p) is admissible if

- for all $j \in [1, p]$, ξ_j is defined and analytic near 0 in E , ξ_j is tangent to E_j , ξ_j vanishes when $v_j = 0$ and $\xi_j(v) \underset{v \rightarrow 0}{=} \vartheta_j(v) + o(\|v\|)$, and
- for all $i \in [1, p]$ and all $j \in [1, p]$, the vector fields ξ_i and ξ_j commute.

The tangency condition in Theorem 1 requires some explanation. We say that an analytic vector field ζ on an open set U is tangent to an analytic set $A \subseteq U$ if $\zeta(a)$ belongs to the tangent space $T_a A$ for every a in the smooth part of A . We say that a germ of an analytic vector field ζ at 0 is tangent to the germ of the postcritical set of F if there is a neighborhood U of 0 such that

- F and ζ are defined and analytic on U ,
- $F : U \rightarrow F(U) \subseteq U$ is proper and
- the vector field ζ is tangent to the critical value set of $F^{\circ n} : U \rightarrow F^{\circ n}(U)$ for all $n \geq 1$.

When F is not postcritically finite, the postcritical set of F is not analytic and the third condition involves tangency to *a priori* infinitely many analytic sets.

1.2. Quasihomogeneous Maps. Let L be a \mathbb{C} -linear space. If $H : L \rightarrow L$ is a homogeneous map of degree $k \geq 1$, then

$$(1) \quad DH \circ \vartheta_{\text{rad}} = k \cdot \vartheta_{\text{rad}} \circ H.$$

This is known as Euler's identity. In fact, the converse is true. If $H : (L, 0) \rightarrow (L, 0)$ is a germ of an analytic map, then H is the germ of a homogeneous map of degree $k \geq 1$ if and only if $DH \circ \vartheta_{\text{rad}} = k \cdot \vartheta_{\text{rad}} \circ H$ near 0. We shall adapt this result to our setting as follows.

Definition 4. A map $H : E \rightarrow E$ is quasihomogeneous of multidegree (k_1, \dots, k_p) if there are homogeneous maps $H_j : E_j \rightarrow E_j$ of degree k_j with $H = H_1 \oplus \dots \oplus H_p$.

The following two lemmas about quasihomogeneous maps will be used later.

Lemma 1. Let $X \subset E$ be such that near 0, the set $H^{-1}X$ coincides with the graph of a function $\varphi : (E_j^\top, 0) \rightarrow (E_j, 0)$. Then $X = H^{-1}X = E_j^\top$ near 0.

Proof. Due to the structure of $H = H_j \oplus H_j^\top : E_j \oplus E_j^\top \rightarrow E_j \oplus E_j^\top$, and due to the homogeneity of H_j , we have that

$$H(e^{2\pi i/k_j} v_j + v_j^\top) = H_j(e^{2\pi i/k_j} v_j) + H_j^\top(v_j^\top) = H_j(v_j) + H_j^\top(v_j^\top) = H(v_j + v_j^\top).$$

So

$$v_j + v_j^\top \in H^{-1}X \iff e^{2\pi i/k_j} v_j + v_j^\top \in H^{-1}X.$$

Since near 0, the set $H^{-1}X$ coincides with the graph of φ , we have

$$v_j = \varphi(v_j^\top) \iff e^{2\pi i/k_j} v_j = \varphi(v_j^\top)$$

for v_j sufficiently close to 0. Thus $\varphi \equiv e^{-2\pi i/k_j} \varphi$ and so φ vanishes identically near 0. This shows that $H^{-1}X = E_j^\top$ near 0, which implies that $X = E_j^\top$ near 0. \square

Lemma 2. Let $H : (E, 0) \rightarrow (E, 0)$ be a germ of an analytic map. Then, H is the germ of a quasihomogeneous map of multidegree (k_1, \dots, k_p) if and only if

$$\forall j \in [1, p], \quad DH \circ \vartheta_j = k_j \cdot \vartheta_j \circ H.$$

Proof. Assume $H = H_1 \oplus \dots \oplus H_p$ with $H_i : E_i \rightarrow E_i$ homogeneous of degree $k_i \geq 1$. Then, $H_i(v_i) = \phi_i(v_i, \dots, v_i)$ with $\phi_i : E_i^{k_i} \rightarrow E_i$ a symmetric k_i -linear map and so, $H_i'(v_i; w_i) = k_i \cdot \phi_i(v_i, \dots, v_i, w_i)$. Thus, for all $v := v_1 + \dots + v_p \in E$,

$$H'(v; v_j) = \sum_{i=1}^p k_i \cdot \phi_i(v_i, \dots, v_i, \pi_i(v_j)) = k_j \cdot \phi_j(v_j, \dots, v_j) = k_j \cdot H_j(v_j).$$

This shows that $DH \circ \vartheta_j(v) = k_j \cdot \vartheta_j \circ H(v)$.

Conversely, assume $DH \circ \vartheta_j = k_j \cdot \vartheta_j \circ H$ for all $j \in [1, p]$. In other words,

$$\forall j \in [1, p], \quad H'(v; v_j) = k_j \cdot \pi_j \circ H(v).$$

Let $(U_j \subseteq E_j)_{j \in [1, p]}$ be neighborhoods of 0 such that

- H is analytic on $U := U_1 + \dots + U_p$ and
- if $v_j \in U_j$ and $|\lambda| \leq 1$, then $\lambda v_j \in U_j$

Let $v = v_1 + \dots + v_p$ be a point in U . Fix $j \in [1, p]$. The map

$$\chi_j : (\lambda_1, \dots, \lambda_p) \mapsto \pi_j \circ H(\lambda_1 v_1 + \dots + \lambda_p v_p)$$

is defined and analytic in a neighborhood of the closed polydisk $\overline{\mathbb{D}}^p$. In addition, for all $i \in [1, p]$,

$$\begin{aligned} \frac{\partial \chi_j}{\partial \lambda_i}(\lambda_1, \dots, \lambda_p) &= \pi_j \circ H'(\lambda_1 v_1 + \dots + \lambda_p v_p; v_i) \\ &= \frac{1}{\lambda_i} \cdot \pi_j \circ H'(\lambda_1 v_1 + \dots + \lambda_p v_p; \lambda_i v_i) \\ &= \frac{k_i}{\lambda_i} \cdot \pi_j \circ \pi_i \circ H(\lambda_1 v_1 + \dots + \lambda_p v_p). \end{aligned}$$

If $i \neq j$, then $\partial \chi_j / \partial \lambda_i = 0$, which shows that χ_j only depends on λ_j . In addition

$$(2) \quad \frac{\partial \chi_j}{\partial \lambda_j} = \frac{k_j}{\lambda_j} \cdot \pi_j \circ \chi_j.$$

Let $H_j : (E_j, 0) \rightarrow (E_j, 0)$ be the restriction of $\pi_j \circ H$ to E_j . The previous discussion shows that

$$\forall (\lambda_1, \dots, \lambda_p) \in \overline{\mathbb{D}}^p, \quad \chi_j(\lambda_1, \dots, \lambda_p) = \chi_j(0, \dots, 0, \lambda_j, 0, \dots, 0) = H_j(\lambda_j v_j).$$

In particular, taking $\lambda_1 = \dots = \lambda_p = 1$, we have that $\pi_j \circ H(v) = H_j(v_j)$, and so $H = H_1 \oplus \dots \oplus H_p$.

Finally, the differential equation (2) implies that

$$\frac{\partial H_j(\lambda v_j)}{\partial \lambda} = \frac{k_j}{\lambda} H_j(\lambda v_j).$$

Consider the map

$$\psi : \lambda \mapsto \lambda^{-k_j} H_j(\lambda v_j)$$

which is defined and analytic in a neighborhood of $\overline{\mathbb{D}}$. Its derivative satisfies

$$\psi'(\lambda) = -k_j \lambda^{-k_j-1} H_j(\lambda v_j) + \lambda^{-k_j} \frac{k_j}{\lambda} H_j(\lambda v_j) = 0.$$

As a consequence, $\psi(\lambda) = \psi(1)$ for all $\lambda \in \overline{\mathbb{D}}$. So $H_j(\lambda v_j) = \lambda^{k_j} H_j(v_j)$ and H_j is the germ of a homogeneous map of degree k_j . \square

1.3. Linearizability of analytic vector fields. The following result is due to Poincaré. We include a proof for completeness.

Lemma 3 (Poincaré). *Any asymptotically radial vector field ξ is linearizable: there is a germ of an analytic map $\Phi : (E, 0) \rightarrow (E, 0)$ such that $\Phi(0) = 0$, $D_0 \Phi = \text{id}$ and $D\Phi \circ \xi = \vartheta_{\text{rad}} \circ \Phi$.*

Proof. Let J be defined in neighborhood of 0 in E by

$$J(v) := \frac{\text{Re} \langle \xi(v) | v \rangle}{\|v\|^2}.$$

Since ξ is asymptotically radial, we have that

$$J(v) \underset{v \rightarrow 0}{=} 1 + O(\|v\|).$$

So, there are constants C and $r > 0$ such that

$$\forall v \in B(0, r), \quad |J(v) - 1| \leq C \|v\| \leq 1/2.$$

In particular, ξ is outward pointing on the boundary of $B(0, r)$. Let $\mathcal{F}_t(v) = \mathcal{F}(t, v)$ be the flow of the vector field ξ . Since ξ is outward pointing on the boundary of $B(0, r)$, the map \mathcal{F}_t is defined and analytic on $B(0, r)$ for all $t \leq 0$.

For $t \leq 0$, set $G_t := \log \|\mathcal{F}_t\|$. Then, for $t \leq 0$,

$$\frac{\partial G_t}{\partial t} = J \circ \mathcal{F}_t \geq 1/2$$

and so, $G_t \leq G_0 + t/2$ and $\|\mathcal{F}_t\| \leq re^{t/2}$. In addition,

$$\left| \frac{\partial(G_t - t)}{\partial t} \right| = |J \circ \mathcal{F}_t - 1| \leq C \|\mathcal{F}_t\| \leq Cre^{t/2}.$$

So,

$$(G_t - t) - (G_0 - 0) \leq \int_t^0 Cre^{u/2} du \leq 2Cr \quad \text{and therefore} \quad e^{-t} \|\mathcal{F}_t\| \leq re^{2Cr}.$$

For $t \leq 0$, let $\Phi_t : B(0, r) \rightarrow E$ be the map defined by

$$\Phi_t := e^{-t} \cdot \mathcal{F}_t.$$

Note that $\Phi_t(0) = 0$, $D_0\Phi_t = \text{id}$. Since $D\mathcal{F}_t = \xi \circ \mathcal{F}_t$, we see that

$$D\Phi_t \circ \xi = e^{-t} \cdot \xi \circ \mathcal{F}_t = e^{-t} \cdot (\vartheta_{\text{rad}} \circ \mathcal{F}_t + o(\|\mathcal{F}_t\|)) = \vartheta_{\text{rad}} \circ \Phi_t + o(\|\Phi_t\|).$$

The previous estimates show that the family $(\Phi_t)_{t \leq 0}$ is uniformly bounded on $B(0, r)$ by re^{2Cr} . Thus, it is normal. Any limit value Φ as $t \rightarrow -\infty$ linearizes ξ :

$$D\Phi \circ \xi = \vartheta_{\text{rad}} \circ \Phi. \quad \square$$

Corollary 1. *Let (ξ_1, \dots, ξ_p) be an admissible p -tuple of germs of vector fields. Then, the ξ_j are simultaneously linearizable, i.e., there is a germ of an analytic map $\Phi : (E, 0) \rightarrow (E, 0)$ such that $\Phi(0) = 0$, $D_0\Phi = \text{id}$ and $D\Phi \circ \xi_j = \vartheta_j \circ \Phi$ for all $j \in [1, p]$.*

Proof. The germ of the vector field $\xi := \xi_1 + \dots + \xi_p$ is asymptotically radial, thus linearizable. Let $\Phi : (E, 0) \rightarrow (E, 0)$ be the linearizer. The vector fields $\Phi_*\xi_j$ commute with $\Phi_*\xi = \vartheta_{\text{rad}}$. It follows that they are linear vector fields and so, $\Phi_*\xi_j = \vartheta_j$. \square

1.4. Lifiable Vector Fields.

Definition 5. *Let $F : U \rightarrow V$ be an analytic map. An analytic vector field ξ on V is liftable if there is an analytic vector field ζ on U which satisfies $DF \circ \zeta = \xi \circ F$. We say that ζ lifts ξ .*

The critical point set of F is the set \mathcal{C}_F of points $x \in U$ for which $D_x F$ is not invertible. The critical value set of F is $\mathcal{V}_F := F(\mathcal{C}_F)$. If ξ is a liftable vector field on V and if ζ lifts ξ , then for all $x \in U - \mathcal{C}_F$, we have

$$\zeta(x) = (D_x F)^{-1} \xi \circ F(x).$$

Thus, when F has discrete fibers, a liftable vector field ξ on V admits a unique lift ζ on U and we shall use the notation

$$F^*\xi := \zeta.$$

Lemma 4 (Arnol'd). *Let $F : U \rightarrow V$ be an analytic map with discrete fibers. Let ξ be a vector field which is analytic on V and tangent to the critical value set \mathcal{V}_F . Then, ξ is liftable and $F^*\xi$ is tangent to \mathcal{C}_F .*

Proof. The vector field $F^*\xi$ is well defined outside the critical set. According to a lemma of Hartogs, it is enough to show that $F^*\xi$ extends analytically outside a subset of U of codimension 2 in order to know that it extends globally.

Since F has discrete fibers, the critical set \mathcal{C}_F is either empty or has codimension 1. Thus, outside a codimension 2 subset of U , the critical set \mathcal{C}_F is smooth. Moreover, it follows from the Constant Rank Theorem that for generic $x \in \mathcal{C}_F$, the kernel of $D_x F$ does not intersect the tangent space to \mathcal{C}_F at x .

Thus, near generic points in \mathcal{C}_F , the map F may be locally expressed as

$$(x_1, \dots, x_{m-1}, x_m) \mapsto (y_1, \dots, y_{m-1}, y_m) = (x_1, \dots, x_{m-1}, x_m^k)$$

for some integer $k \geq 2$. Since the vector field ξ is tangent to the critical value set $\{y_m = 0\}$, it is of the form

$$\xi = \xi_1 \frac{\partial}{\partial y_1} + \dots + \xi_{m-1} \frac{\partial}{\partial y_{m-1}} + y_m \xi_m \frac{\partial}{\partial y_m}$$

and

$$\begin{aligned} F^*\xi &= (\xi_1 \circ F) \frac{\partial}{\partial x_1} + \dots + (\xi_{m-1} \circ F) \frac{\partial}{\partial x_{m-1}} + \left(\frac{x_m^k}{k x_m^{k-1}} \xi_m \circ F \right) \frac{\partial}{\partial x_m} \\ &= (\xi_1 \circ F) \frac{\partial}{\partial x_1} + \dots + (\xi_{m-1} \circ F) \frac{\partial}{\partial x_{m-1}} + \left(\frac{1}{k} x_m \xi_m \circ F \right) \frac{\partial}{\partial x_m} \end{aligned}$$

which clearly extends analytically through the critical set $\{x_m = 0\}$ and is tangent to the critical set. \square

Lemma 5. *Let $F : U \rightarrow V$ be an analytic map with discrete fibers, let ξ be a liftable vector field on V , and let $\zeta := F^*(\xi)$ be its lift to U . Let $\phi(t, z)$ be the flow of ξ , and $\psi(t, z)$ be the flow of ζ .*

(1) *For all $z \in U$ and for t sufficiently small, we have*

$$F(\phi(t, z)) = \psi(t, F(z)).$$

(2) *If $\zeta \circ F(z) = 0$, then $\xi(z) = 0$.*

(3) *If ζ is tangent to an analytic set $A \subseteq U$, then ξ is tangent to $F^{-1}(A)$.*

Proof. If $z \in U - \mathcal{C}_F$, then $F : (U, z) \rightarrow (V, F(z))$ is a local isomorphism. It sends the vector field ξ to the vector field ζ . Thus it conjugates their flows, and the equality in part (1) holds for $z \in U - \mathcal{C}_F$ and t is sufficiently small. If $z \in \mathcal{C}_F$, it holds by analytic continuation (with respect to z).

Parts (2) and (3) follow immediately since when the flow of ζ preserves an analytic set A (which may be reduced to a point if ζ vanishes at this point), then the flow of ξ preserves the analytic set $F^{-1}(A)$. \square

We shall now study how admissible p -tuple of vector fields behave under pullback.

Lemma 6. *Assume $F : (E, 0) \rightarrow (E, 0)$ is an analytic germ having an adapted superattracting fixed point. Let $(\zeta_1, \dots, \zeta_p)$ be an admissible p -tuple of liftable vector fields. Then, $(k_1 \cdot F^*\zeta_1, \dots, k_p \cdot F^*\zeta_p)$ is an admissible p -tuple of vector fields.*

Proof. Fix $j \in [1, p]$, and set $\xi_j := k_j F^*\zeta_j$. Since the vector field ζ_j is tangent to E_j , the vector field ξ_j is tangent to $F^{-1}(E_j)$. According to proposition 1, $F^{-1}(E_j)$ coincides with E_j in a neighborhood of 0. Thus, the vector field ξ_j is tangent to E_j .

Since the vector field ζ_j vanishes on $E_j^\top := \{v_j = 0\}$, the vector field ξ_j vanishes on $F_j^{-1}(E_j^\top)$. According to proposition 1, $F^{-1}(E_j^\top)$ coincides with E_j^\top in a neighborhood of 0. So ξ_j vanishes on E_j^\top .

Since ξ_j vanishes at 0, we may write

$$\xi_j(v) = \tau_j(v) + o(\|v\|) \quad \text{with} \quad \tau_j(v) := (v; A_j(v))$$

for some linear map $A_j : E \rightarrow E$. It remains to prove that $\tau_j = \vartheta_j$ which amounts to showing that $A_j = \pi_j$. Since ξ_j vanishes on E_j^\top , the linear map A_j vanishes on E_j^\top . Thus it suffices to show that the restriction of A to E_j is the identity.

The map F restricts to a self-map $F_j : E_j \rightarrow E_j$. The vectors fields ζ_j and ξ_j restrict to vector fields on E_j (because they are tangent to E_j). It suffices to show that $A_j(v_j) = v_j$. We therefore restrict our analysis to the space E_j , omitting the index j :

- $F(v) = H(v) + o(\|v\|^k)$ with $H : E \rightarrow E$ a nondegenerate homogeneous map of degree k ,
- $\zeta(v) = (v; v) + o(\|v\|)$,
- $\xi(v) = \tau(v) + o(\|v\|)$, and
- $DF \circ \xi = k \cdot \zeta \circ F$.

On the one hand,

$$\begin{aligned} DF \circ \xi(v) &= (F(v); F' \circ \xi(v)) = (F(v); H' \circ \xi(v) + o(\|v\|^{k-1} \cdot \|\xi(v)\|)) \\ &= (F(v); H' \circ \tau(v) + o(\|v\|^k)) \end{aligned}$$

On the other hand,

$$k \cdot \zeta \circ F(v) = (F(v); k \cdot F(v) + o(\|F(v)\|)) = (F(v); k \cdot H(v) + o(\|v\|^k)).$$

It follows that $H' \circ \tau(v) = k \cdot H(v)$, thus

$$DH \circ \tau(v) = k \cdot (H(v); H(v)).$$

According to Euler's identity, we therefore have $\tau(v) = (v; v)$ which implies that $A(v) = v$ as required. \square

1.5. Dynamical Green Functions. We shall use dynamical Green functions introduced by Hubbard and Papadopol [HP]. We will first recall the construction for homogeneous maps, and then explain how this construction may be adapted to our setting.

Let $H : L \rightarrow L$ be a nondegenerate homogeneous map of degree k on a \mathbb{C} -linear space L of dimension m . Then, the function

$$u_H : v \mapsto \frac{1}{k} \log \|H(v)\| - \log \|v\|$$

is defined and bounded on $L - \{0\}$. It follows that the sequence of plurisubharmonic functions

$$\mathcal{G}_H^n := \frac{1}{k^n} \log \|H^{\circ n}\| = \mathcal{G}_H^0 + \sum_{i=0}^{n-1} \frac{u_H \circ F^{\circ i}}{k^i} : L \rightarrow \mathbb{R} \cup \{-\infty\}$$

converges uniformly on L to a plurisubharmonic function $\mathcal{G}_H : L \rightarrow \mathbb{R} \cup \{-\infty\}$ which is continuous on $L - \{0\}$ and satisfies

$$\mathcal{G}_H(v) = \log \|v\| + \mathcal{O}(1) \quad \text{and} \quad \mathcal{G}_H \circ H = k \cdot \mathcal{G}_H.$$

We shall adapt this construction to our setting as follows.

Lemma 7. *Let $F : (E, 0) \rightarrow (E, 0)$ be an analytic germ having an adapted superattracting fixed point. There is a neighborhood U of 0 in E such that for all $j \in [1, p]$, the sequence of functions*

$$\mathcal{G}_j^n := \frac{1}{k_j^n} \log \|\pi_j \circ F^{\circ n}\| : U \rightarrow \mathbb{R} \cup \{-\infty\}$$

converges locally uniformly in U to a plurisubharmonic function $\mathcal{G}_j : U \rightarrow \mathbb{R} \cup \{-\infty\}$ satisfying,

$$\mathcal{G}_j(v) \underset{v \rightarrow 0}{=} \log \|\pi_j(v)\| + O(1) \quad \text{and} \quad \mathcal{G}_j \circ F = k_j \cdot \mathcal{G}_j.$$

Proof. Let $U \subset E$ be a sufficiently small neighborhood of 0 so that F is defined and analytic on U , $F(U) \subseteq U$ and U is contained in the basin of attraction of 0. The functions

$$\mathcal{G}_j^n := \frac{1}{k_j^n} \log \|\pi_j \circ F^{\circ n}\| : U \rightarrow \mathbb{R} \cup \{-\infty\}$$

are then defined and plurisubharmonic on U .

By assumption

$$F_j(v) = H_j(v_j) + o(\|v_j\|^{k_j}).$$

Since H_j is nondegenerate, $\|v_j\|^{k_j} = \mathcal{O}(\|H_j(v_j)\|)$. Thus

$$\|\pi_j \circ F(v)\| = \|F_j(v)\| \underset{v \rightarrow 0}{\sim} \|H_j(v_j)\| = \|H_j \circ \pi_j(v)\|.$$

As a consequence, restricting U if necessary, the function

$$\log \|\pi_j \circ F\| - \log \|H_j \circ \pi_j\|$$

is defined and bounded in $U - E_j^\top$. It follows that the function

$$u_j := \frac{1}{k_j} \log \|\pi_j \circ F\| - \log \|\pi_j\|$$

is defined and bounded in $U - E_j^\top$.

Now, the sequence

$$\mathcal{G}_j^N = \mathcal{G}_j^0 + \sum_{n=0}^{N-1} \frac{u_j \circ F^{\circ n}}{k_j^n} : U \rightarrow \mathbb{R} \cup \{-\infty\}$$

converges uniformly on U to a plurisubharmonic function \mathcal{G}_j whose difference with $\mathcal{G}_j^0 = \log \|\pi_j\|$ is bounded as required. \square

1.6. Cartan's Lemma. We shall use the following lemma of Cartan which is a multidimensional version of the Schwarz Lemma. We include a proof for completeness.

Lemma 8 (Cartan). *Let V be a bounded connected open subset of E containing 0, let $\Phi : V \rightarrow V$ be an analytic map such that $\Phi(0) = 0$ and $D_0\Phi = \text{id}$. Then, $\Phi = \text{id}$.*

Proof. The iterates $\Phi^{\circ n}$ are defined on V for all $n \geq 0$. For $n \geq 1$, let $\Psi_n : V \rightarrow E$ be defined as the average

$$\Psi_n := \frac{1}{n} \sum_{j=0}^{n-1} \Phi^{\circ j}.$$

Then,

$$\Psi_n(0) = 0, \quad D_0\Psi_n = \text{id} \quad \text{and} \quad \Psi_n \circ \Phi = \Psi_n + \frac{\Phi^{\circ n} - \text{id}}{n}.$$

In addition, the sequence $(\Psi_n)_{n \geq 1}$ is normal. Let Ψ be a limit value. Then,

$$\Psi(0) = 0, \quad D_0\Psi = \text{id} \quad \text{and} \quad \Psi \circ \Phi = \Psi.$$

In particular, Ψ is invertible at 0 and Φ is equal to the identity near 0, thus in V by analytic continuation. \square

Corollary 2. *Let V be a bounded connected open subsets of E containing 0, let $\Phi : V \rightarrow W \subset E$ be an isomorphism and $\Psi : V \rightarrow W$ be an analytic map such that $\Psi(0) = \Phi(0)$ and $D_0\Psi = D_0\Phi$. Then, $\Psi = \Phi$.*

Proof. Apply Cartan's lemma to $\Phi^{-1} \circ \Psi : V \rightarrow V$. \square

1.7. Proof of Theorem 1. (1) \Rightarrow (3). For all $n \geq 1$, $H^{\circ n} = H_1^{\circ n} \oplus \dots \oplus H_p^{\circ n}$ and the critical value set of $H^{\circ n}$ is

$$\mathcal{V}_{H^{\circ n}} = \mathcal{V}_{H_1^{\circ n}} + \dots + \mathcal{V}_{H_p^{\circ n}}.$$

For all $j \in [1, p]$ and all $n \geq 1$, the critical value set of $H_j^{\circ n} : E_j \rightarrow E_j$ is homogeneous (*i.e.*, a complex cone with vertex at the origin). Thus, for all $j \in [1, p]$, the vector field ϑ_j is tangent to the critical value set of $H^{\circ n}$ and the vector field $\zeta_j := \Phi^* \vartheta_j$ is tangent to the critical value locus of $F^{\circ n}$.

For all i, j , the vector fields ϑ_i and ϑ_j commute, thus ζ_i and ζ_j commute. Since $D_0\Phi = \text{id}$ the linear part of ζ_j at 0 is ϑ_j . Since the vector field ϑ_j is tangent to E_j and vanishes on E_j^\top , the vector field ζ_j is tangent to $\Phi^{-1}(E_j)$ and vanishes on $\Phi^{-1}(E_j^\top)$.

We claim that for all $j \in [1, p]$, we have that $\Phi(E_j^\top) = E_j^\top$ near 0. Indeed, recall that E_j^\top is locally totally invariant by F (see proposition 1). So $X := \Phi(E_j^\top)$ is locally totally invariant by H . Since $D_0\Phi = \text{id}$, X and thus $H^{-1}X$ is locally the graph of a function $\varphi : E_j^\top \rightarrow E_j$. According to lemma 1, $X = H^{-1}X = E_j^\top$.

It follows that $\Phi^{-1}(E_j^\top) = E_j^\top$ near 0. In particular, $\Phi^{-1}(E_j) = E_j$ near 0 (because E_j is the intersection of all the E_i^\top for $i \neq j$). Consequently, ζ_j is tangent to E_j and vanishes on E_j^\top . Thus the p -tuple of vector fields $(\zeta_1, \dots, \zeta_p)$ is admissible.

(2) \Rightarrow (1). According to corollary 1, since (ξ_1, \dots, ξ_p) is admissible, there exists a germ of an analytic map $\Phi : (E, 0) \rightarrow (E, 0)$ such that $D_0\Phi = \text{id}$ and $D\Phi \circ \xi_j = \vartheta_j \circ \Phi$ for all $j \in [1, p]$. Then, Φ conjugates F to a map \check{F} , defined and analytic near 0 in E , satisfying

$$D\check{F} \circ \vartheta_j = k_j \cdot \vartheta_j \circ \check{F}.$$

According to lemma 2, \check{F} is quasihomogeneous with multidegree (k_1, \dots, k_p) . Since $D_0\Phi = \text{id}$ we have that $\check{F} = H$.

(3) \Rightarrow (2). Let U_0 be a sufficiently small neighborhood of 0 in E so that

- F is defined and analytic on U_0 , $F(U_0) \subseteq U_0$, and U_0 is contained in the attracting basin of 0,
- ζ_j is defined and analytic on U_0 , tangent to the critical value set of $F^{\circ n} : U_0 \rightarrow F^{\circ n}(U_0)$ for all $n \geq 0$ and all $j \in [1, p]$.

Then, ζ_j is liftable by $F^{\circ n}$ and we may define a holomorphic vector field ζ_j^n on U_0 by

$$\zeta_j^n := k_j^n \cdot (F^{\circ n})^* \zeta_j.$$

Then, for all $n \geq 0$, we have that $DF \circ \zeta_j^{n+1} = k_j \cdot \zeta_j^n \circ F$. According to lemma 6, the p -tuple of vector fields $(\zeta_1^n, \dots, \zeta_p^n)$ is admissible. We will show that there is a neighborhood V of 0 in E on which the sequence of vector fields $(\zeta_j^n)_{n \geq 0}$ converges uniformly to a vector field ξ_j for all $j \in [1, p]$. Then, the p -tuple of vector fields (ξ_1, \dots, ξ_p) is admissible and satisfies $DF \circ \xi_j = k_j \cdot \xi_j \circ F$ for all $j \in [1, p]$. The proof will be completed.

According to lemma 7, the sequence of functions

$$\mathcal{G}_{F,j}^n := \frac{1}{k_j^n} \log \|\pi_j \circ F^{o^n}\| : U_0 \rightarrow \mathbb{R} \cup \{-\infty\}$$

converges locally uniformly in U_0 to a function $\mathcal{G}_{F,j} : U_0 \rightarrow \mathbb{R} \cup \{-\infty\}$ which is plurisubharmonic and satisfies

$$\mathcal{G}_{F,j}(v) \underset{v \rightarrow 0}{=} \log \|\pi_j(v)\| + \mathcal{O}(1) \quad \text{and} \quad \mathcal{G}_{F,j} \circ F = k_j \cdot \mathcal{G}_{F,j}.$$

We set

$$\mathcal{G}_F^n := \max_{j \in [1, p]} \mathcal{G}_{F,j}^n \quad \text{and} \quad \mathcal{G}_F := \max_{j \in [1, p]} \mathcal{G}_{F,j}.$$

Note that those functions are plurisubharmonic in U_0 and take the value $-\infty$ only at 0. In addition, the sequence of functions \mathcal{G}_F^n converges locally uniformly to \mathcal{G}_F in U_0 . In particular, if $M > 0$ is sufficiently large, the level sets $\{\mathcal{G}_F^n < -M\}$ are compactly contained in U_0 . From now on, we assume that $M > 0$ is sufficiently large so that the set

$$V_n := \{v \in U_0 : \forall j \in [1, p], \mathcal{G}_{F,j}^n(v) < -M\}$$

are compactly contained in U_0 for all $n \geq 0$.

Similarly, the sequence of plurisubharmonic functions

$$\mathcal{G}_{H,j}^n := \frac{1}{k_j^n} \log \|\pi_j \circ H^{o^n}\| : E \rightarrow \mathbb{R} \cup \{-\infty\}$$

converges locally uniformly in E to a function $\mathcal{G}_{H,j} : E \rightarrow \mathbb{R} \cup \{-\infty\}$ which is plurisubharmonic and satisfies

$$\mathcal{G}_{H,j}(v) \underset{v \rightarrow 0}{=} \log \|\pi_j(v)\| + \mathcal{O}(1) \quad \text{and} \quad \mathcal{G}_{H,j} \circ H = k_j \cdot \mathcal{G}_{H,j}.$$

We set

$$\mathcal{G}_H^n := \max_{j \in [1, p]} \mathcal{G}_{H,j}^n, \quad \mathcal{G}_H := \max_{j \in [1, p]} \mathcal{G}_{H,j} \quad \text{and} \quad W_n := \{v \in E : \mathcal{G}_H^n(v) < -M\}.$$

According to corollary 1, there are germs of analytic maps $\Phi_n : (E, 0) \rightarrow (E, 0)$ such that $D_0 \Phi_n = \text{id}$ and $D \Phi_n \circ \zeta_j^n = \vartheta_j \circ \Phi_n$ for all $n \geq 0$ and all $j \in [1, p]$. According to lemma 2, for all $n \geq 0$, the map $\Phi_n \circ F \circ \Phi_{n+1}^{-1} : (E, 0) \rightarrow (E, 0)$ is quasihomogeneous, thus equal to H . In other words, we have the following commutative diagram:

$$\begin{array}{ccc} (E, 0) & \xrightarrow{\Phi_{n+1}} & (E, 0) \\ F \downarrow & & \downarrow H \\ (E, 0) & \xrightarrow{\Phi_n} & (E, 0). \end{array}$$

Lemma 9. *For all $n \geq 0$, the linearizing map Φ_n is analytic on V_n .*

Proof. Let us first show that the linearizing map Φ_n is defined on V_n . Recall that Φ_n is defined as the linearizing map of the asymptotically radial vector field

$$\zeta^n := \zeta_1^n + \cdots + \zeta_p^n.$$

Note that flowing along ζ_j^0 during time $t < 0$ decreases $\mathcal{G}_{F,j}^0$ by t . It follows that flowing along $(F^{\circ n})^* \zeta_j^0$ during time $t < 0$ decreases $\mathcal{G}_{F,j}^0 \circ F^{\circ n}$ by t . Thus, flowing along ζ_j^n during time $t < 0$ decreases $\mathcal{G}_{F,j}^n$ by t . Finally, since the vector fields $(\zeta_j^n)_{j \in [1,p]}$ commute, flowing along ζ^n during time $t < 0$ decreases \mathcal{G}_F^n by t .

In particular, the flow of ζ^n is defined on V_n for all $t < 0$, every trajectory remains in V_n and converges to 0. If we denote by \mathcal{F}_n this flow, the linearizing map Φ_n may be obtained on V_n as

$$\Phi_n(v) = \lim_{t \rightarrow -\infty} e^{-t} \cdot \mathcal{F}_n(t, x).$$

□

Since $DF \circ \zeta_j^{n+1} = k_j \cdot \zeta_j^n \circ F$, the analytic germ \check{F} defined in a neighborhood of 0 in E by $\check{F} := \Phi_n \circ F \circ \Phi_{n+1}^{-1}$ satisfies $D\check{F} \circ \vartheta_j = k_j \cdot \vartheta_j \circ \check{F}$. Thus, it is the germ of a quasihomogeneous map of multidegree (k_1, \dots, k_p) . Since, $D_0\Phi_n = D_0\Phi_{n+1} = \text{id}$, we have that $\check{F} = H$. So, we have the following commutative diagram:

$$\begin{array}{ccc} V_n & \xrightarrow{\Phi_n} & E \\ F^{\circ n} \downarrow & & \downarrow H^{\circ n} \\ F^{\circ n}(V_n) & \xrightarrow{\Phi_0} & E. \end{array}$$

Lemma 10. *The function*

$$u_j := \log \|\pi_j \circ \Phi_0\| - \log \|\pi_j\|$$

is defined and bounded on $U_0 - E_j^\top$.

Proof. Since $D_0\Phi_0 = \text{id}$, we have the a priori estimate

$$\pi_j \circ \Phi_0(v) \underset{v \rightarrow 0}{=} \pi_j(v) + o(\|v\|).$$

We claim that $X := \Phi_0(E_j^\top)$ coincides with E_j^\top near 0. Indeed, E_j^\top is locally totally invariant by F . So $H^{-1}X = \Phi_1(E_j^\top)$ is a graph of a function $\varphi : (E_j^\top, 0) \rightarrow (E_j, 0)$ near 0. According to lemma 1, $X = H^{-1}X = E_j^\top$.

Consequently, $\pi_j \circ \Phi_0$ vanishes on E_j^\top , and so

$$\pi_j \circ \Phi_0(v) \underset{v \rightarrow 0}{=} \pi_j(v) + o(\|\pi_j(v)\|).$$

The lemma follows immediately. □

Observe that

$$\begin{aligned} \mathcal{G}_{H,j}^n \circ \Phi_n &= \frac{1}{k_j^n} \log \|\pi_j \circ H^{\circ n} \circ \Phi_n\| \\ &= \frac{1}{k_j^n} \log \|\pi_j \circ \Phi_0 \circ F^{\circ n}\| = \frac{1}{k_j^n} u_j \circ F^{\circ n} + \mathcal{G}_{F,j}^n. \end{aligned}$$

Setting

$$V := \{v \in U_0 : \mathcal{G}_F(v) < -M\} \quad \text{and} \quad W := \{v \in U_0 : \mathcal{G}_H(v) < -M\},$$

we deduce that $\mathcal{G}_H^n \circ \Phi_n$ converges uniformly to \mathcal{G}_F^n on every compact subset of V . Therefore the sequence (Φ_n) is uniformly bounded on every compact subset of V . Similarly, any compact subset of W is contained in the image of Φ_n for n large enough, and the sequence (Φ_n^{-1}) is uniformly bounded on every compact subset of W .

Thus, the sequence of maps (Φ_n) is normal on any compact subset of V and the sequence of maps (Φ_n^{-1}) is normal on any compact subset of W . Extracting a subsequence, we see that there is an isomorphism $\Phi : V \rightarrow W$ such that $\Phi(0) = 0$ and $D_0\Phi = \text{id}$. According to Cartan's lemma, any limit value of the sequence (Φ_n) must coincide with Φ . Thus, the whole sequence (Φ_n) converges to Φ locally uniformly in V and the sequence $(\zeta_j^n = \Phi_n^* \vartheta_j)$ converges to $\xi_j = \Phi^* \vartheta_j$.

This completes the proof of theorem 1.

2. THE GLOBAL RESULT

We will now prove Theorem 2. Since $F : \Omega \rightarrow \Omega$ is proper and since $\mathcal{B}_a(F)$ is a connected component of $F^{-1}(\mathcal{B}_a(F))$, the restriction $F : \mathcal{B}_a(F) \rightarrow \mathcal{B}_a(F)$ is proper.

Let $\mathcal{G}_{H,j} : E \rightarrow \mathbb{R} \cup \{-\infty\}$ and $\mathcal{G}_H : E \rightarrow \mathbb{R} \cup \{-\infty\}$ be the dynamical Green functions of H introduced above. Let $\mathcal{G}_{F,j} : \mathcal{B}_a(F) \rightarrow \mathbb{R} \cup \{-\infty\}$ be defined by

$$\mathcal{G}_{F,j}(x) = \frac{1}{k_j^n} \mathcal{G}_{H,j} \circ \Phi \circ F^{on}(x)$$

where $n \geq 0$ is chosen sufficiently large so that $F^{on}(x)$ belongs to a neighborhood of a on which Φ is defined. Let $\mathcal{G}_F : \mathcal{B}_a(F) \rightarrow \mathbb{R} \cup \{-\infty\}$ be defined by

$$\mathcal{G}_F := \max_{j \in [1,p]} \mathcal{G}_{F,j}.$$

Let $M > 0$ be sufficiently large so that $\Phi : (E, 0) \rightarrow (E, 0)$ has an inverse branch defined on

$$W := \{v \in E : \mathcal{G}_F(v) < -M\}.$$

Set $V := \Phi^{-1}(W)$, so that $\Phi : V \rightarrow W$ is an isomorphism. Increasing M if necessary, we see that V is relatively compact in $\mathcal{B}_a(F)$.

For $j \in [1, p]$, set

$$\xi_j := \Phi^* \vartheta_j$$

which is defined and analytic near 0. Then, $\xi_j = k_j \cdot F^* \xi_j$ near a . Since ϑ_j is tangent to the postcritical set of H , and since near a Φ maps the postcritical set of $F : \mathcal{B}_a(F) \rightarrow \mathcal{B}_a(F)$ to the postcritical set of $H : \mathcal{B}_0(H) \rightarrow \mathcal{B}_0(H)$, the vector field ξ_j is tangent to the postcritical set of $F : \mathcal{B}_a(F) \rightarrow \mathcal{B}_a(F)$. In particular, ξ_j is tangent to the critical value set of F^{on} for n large enough. Thus we can extend ξ_j to the whole basin of attraction $\mathcal{B}_a(F)$ using the formula $\xi_j = k_j^n \cdot (F^{on})^* \xi_j$ for n large enough. We therefore have a vector field ξ_j which is defined and analytic on $\mathcal{B}_a(F)$ and satisfies $\xi_j = k_j \cdot F^* \xi_j$ on $\mathcal{B}_a(F)$. Let $(t, x) \mapsto \mathcal{F}_{t,j}(x)$ be the flow of the vector field ξ_j and let $(t, x) \mapsto \mathcal{F}_t(x)$ be the flow of the vector field

$$\xi := \xi_1 + \cdots + \xi_p.$$

Lemma 11. *The map \mathcal{F}_t is defined on $\mathcal{B}_a(F)$ for all $t \leq 0$. For all $x \in \mathcal{B}_a(F)$, we have that $\mathcal{F}_t(x) \rightarrow 0$ as $t \rightarrow -\infty$.*

Proof. We first want to prove that \mathcal{F}_t is defined on $\mathcal{B}_a(F)$ for all $t \leq 0$. The maps $\mathcal{F}_{t,j}$ are defined and analytic on V for all $t \leq 0$. Indeed, $\Phi : V \rightarrow W$ conjugates $\mathcal{F}_{t,j}$ to the linear map

$$\mathcal{H}_{t,j} : v_1 + \cdots + v_p \mapsto v_1 + \cdots + v_{j-1} + e^t v_j + v_{j+1} + \cdots + v_p.$$

The equality $\xi_j = k_j \cdot F^* \xi_j$ yields $F \circ \mathcal{F}_{t,j} = \mathcal{F}_{k_j t, j} \circ F$ and so, $F^{\circ n} \circ \mathcal{F}_{t,j} = \mathcal{F}_{k_j^n t, j} \circ F^{\circ n}$. For all $x \in \mathcal{B}_a(F)$, if n is large enough, $\mathcal{F}_{k_j^n t} \circ F^{\circ n}(x)$ remains in a compact subset of $\mathcal{B}_a(F)$ for all $t \leq 0$. Since $F^{\circ n} : \mathcal{B}_a(F) \rightarrow \mathcal{B}_a(F)$ is proper, $\mathcal{F}_{t,j}(x)$ also remains in a compact subset of $\mathcal{B}_a(F)$, and so, is defined for all $t \leq 0$. Since the vector fields ξ_j commute, we have that

$$\mathcal{F}_t = \mathcal{F}_{t,1} \circ \cdots \circ \mathcal{F}_{t,p}.$$

So, \mathcal{F}_t is defined on $\mathcal{B}_a(F)$ for all $t \leq 0$.

Near 0, we have that $\mathcal{G}_{F,j} = \mathcal{G}_{H,j} \circ \Phi$ and $\Phi \circ \mathcal{F}_{t,j} = \mathcal{H}_{t,j} \circ \Phi$. In addition, $\mathcal{G}_{H,j} \circ \mathcal{H}_{t,j} = \mathcal{G}_{H,j} + t$. It follows that near 0, we have that

$$\mathcal{G}_{F,j} \circ \mathcal{F}_{t,j} = \mathcal{G}_{F,j} + t.$$

The equality $\xi_j = k_j \cdot F^* \xi_j$ yields $F \circ \mathcal{F}_{t,j} = \mathcal{F}_{k_j t, j} \circ F$. So, for $t \leq 0$, the following equality is valid on $\mathcal{B}_a(F)$:

$$\begin{aligned} \mathcal{G}_{F,j} \circ \mathcal{F}_{t,j} &= \frac{1}{k_j^n} \mathcal{G}_{F,j} \circ F^{\circ n} \circ \mathcal{F}_{t,j} = \frac{1}{k_j^n} \mathcal{G}_{F,j} \circ \mathcal{F}_{k_j^n t, j} \circ F^{\circ n} \\ &= \frac{1}{k_j^n} \mathcal{G}_{F,j} \circ F^{\circ n} + t = \mathcal{G}_{F,j} + t. \end{aligned}$$

As a consequence, for $t \leq 0$, the following equality is valid on $\mathcal{B}_a(F)$:

$$\mathcal{G}_F \circ \mathcal{F}_t = \mathcal{G}_F \circ \mathcal{F}_{t,1} \circ \cdots \circ \mathcal{F}_{t,p} = \mathcal{G}_F + t.$$

We now prove that $F^{-1}\{0\} = \{0\}$. As we have seen, each trajectory $(\mathcal{F}_t(x))_{t \leq 0}$ remains in a compact subset of $\mathcal{B}_a(F)$. Since $\mathcal{G}_F \circ \mathcal{F}_t = \mathcal{G}_F + t$, as $t \rightarrow -\infty$, each trajectory $(\mathcal{F}_t(x))_{t \leq 0}$ must converge to a point where \mathcal{G}_F takes the value $-\infty$, i.e., to a point in the backward orbit of 0. We may therefore partition $\mathcal{B}_a(F)$ in the basins of those points (for the flow \mathcal{F}_t). The basins are open and since $\mathcal{B}_a(F)$ is connected, there is only one such basin: the basin of 0. \square

The linearizer Φ of ξ extends to the whole set $\mathcal{B}_a(F)$ by

$$\Phi(x) = e^{-t} \cdot \Phi \circ \mathcal{F}_t(x)$$

where $t \leq 0$ is chosen sufficiently negative so that $\mathcal{F}_t(x) \in V$.

It is injective on $\mathcal{B}_a(F)$. Indeed, assume $\Phi(x_1) = \Phi(x_2)$ with x_1 and x_2 in $\mathcal{B}_a(F)$. Choose $t \leq 0$ sufficiently negative so that $\mathcal{F}_t(x_1)$ and $\mathcal{F}_t(x_2)$ belong to V . Then,

$$\Phi \circ \mathcal{F}_t(x_1) = e^t \Phi(x_1) = e^t \Phi(x_2) = \Phi \circ \mathcal{F}_t(x_2)$$

Since $\Phi : V \rightarrow W$ is an isomorphism, we have that $\mathcal{F}_t(x_1) = \mathcal{F}_t(x_2)$, and so,

$$x_1 = \mathcal{F}_{-t} \circ \mathcal{F}_t(x_1) = \mathcal{F}_{-t} \circ \mathcal{F}_t(x_2) = x_2.$$

The equality $\Phi \circ F = H \circ \Phi$ holds on $\mathcal{B}_a(F)$ by analytic continuation. This shows that $\Phi(\mathcal{B}_a(F))$ is contained in the basin of attraction $\mathcal{B}_0(H)$.

Finally, $\Phi : \mathcal{B}_a(F) \rightarrow \mathcal{B}_0(H)$ is proper, thus an isomorphism. Indeed, let $K \subset \mathcal{B}_0(H)$ be a compact set and let $n \geq 0$ be sufficiently large so that $K_n := H^{\circ n}(K) \subset W$. Since $\Phi : V \rightarrow W$ is an isomorphism, $\Phi^{-1}(K_n) \subset V$ is compact. Since $F^{\circ n} : \mathcal{B}_a(F) \rightarrow \mathcal{B}_a(F)$ is proper, $F^{-n}(\Phi^{-1}(K_n))$ is compact. This compact set

contains $\Phi^{-1}(K)$ which is closed since Φ is continuous. It follows that $\Phi^{-1}(K)$ is compact.

This completes the proof that $\Phi : \mathcal{B}_a(F) \rightarrow \mathcal{B}_0(H)$ is an isomorphism.

3. APPLICATIONS

In this section we apply our theorems to a family of postcritically finite endomorphisms of projective space which arose in [K] and were studied in [BEKP].

Let $h : \mathbb{P}^m \rightarrow \mathbb{P}^m$ be an endomorphism; that is, a holomorphic map with no indeterminacy points. Let \mathcal{C}_h be critical locus of h . We define the *postcritical locus* of h to be

$$\mathcal{P}_h := \bigcup_{n \geq 1} h^{\circ n}(\mathcal{C}_h).$$

The endomorphism h is *postcritically finite* if \mathcal{P}_h is algebraic. Equivalently (via a Baire category argument), each component of \mathcal{C}_h is either periodic, or preperiodic to a periodic cycle of components in \mathcal{P}_h . Postcritically finite endomorphisms of \mathbb{P}^m were first studied by Fornæss and Sibony in [FS], and by Ueda in [Ue2].

3.1. Constructing endomorphisms. The following construction is a particular case of a more general construction in [K]. Let I be a finite set of cardinality $m \geq 1$. Denote by E the \mathbb{C} -vector space of functions $x : I \rightarrow \mathbb{C}$ whose average is 0. For $x \in E$, we use the notation $x_i := x(i)$ and set

$$P_x(t) := \frac{m+1}{m} \sum_{j \in I} \int_{x_j}^t \prod_{i \in I} (w - x_i) dw.$$

The polynomial P_x is the unique monic centered polynomial of degree $m+1$ whose critical points are the points $(x_i)_{i \in I}$, repeated according to their multiplicities, and for which the barycenter of the critical values $(P_x(x_i))_{i \in I}$ is 0.

The function $y := P_x \circ x : I \rightarrow \mathbb{C}$ belongs to E and satisfies

$$\forall i \in I \quad y_i = P_x(x_i).$$

We denote by $H : E \rightarrow E$ the map defined by

$$H(x) := P_x \circ x.$$

Proposition 2. *For $m \geq 2$, the map $H : E \rightarrow E$ is a homogeneous map of degree $m+1$. For $m \geq 3$ it induces an endomorphism $h : \mathbb{P}(E) \rightarrow \mathbb{P}(E)$, where $\mathbb{P}(E)$ is the projective space associated to E (isomorphic to $\mathbb{P}^{m-2}(\mathbb{C})$).*

Proof. The polynomial P_x depends analytically on $x \in E$, therefore H is analytic. Since

$$\begin{aligned} P_{\lambda x}(\lambda t) &= \frac{m+1}{m} \sum_{j \in I} \int_{\lambda x_j}^{\lambda t} \prod_{i \in I} (w - \lambda x_i) dw \\ &= \frac{m+1}{m} \sum_{w=\lambda v} \int_{x_j}^t \prod_{i \in I} (\lambda v - \lambda x_i) d(\lambda v) = \lambda^{m+1} P_x(t), \end{aligned}$$

we have $H(\lambda x) = \lambda^{m+1} H(x)$, so H is homogeneous of degree $m+1$.

Assume $P_x(x_i) = 0$ for all $i \in I$. This means that P_x has only one critical value, namely 0. Therefore P_x has only one critical point. Thus x is constant, and the average of x is 0, we have $x = 0$. This implies that $H^{-1}(0) = \{0\}$, and consequently, $H : E \rightarrow E$ induces an endomorphism $h : \mathbb{P}(E) \rightarrow \mathbb{P}(E)$. \square

The significance of these endomorphisms is that their fixed points correspond to polynomials with fixed critical points. More precisely, we have the following correspondence.

Proposition 3. *If x is a fixed point of $H : E \rightarrow E$, then P_x is a monic centered polynomial of degree $m + 1$ with fixed critical points. If P is a monic centered polynomial of degree $m + 1$ with fixed critical points, then there exists $x \in E$ such that $H(x) = x$, and $P_x = P$.*

Proof. First let x be a fixed point of H . The polynomial P_x is monic, centered, and of degree $m + 1$. The critical points of P_x are the x_i . Since x is a fixed point of H , $x_i = P_x(x_i)$ for all $i \in I$, and thus P_x fixes its critical points.

Conversely, let P be a monic centered polynomial of degree $m + 1$ with fixed critical points. The polynomial P has exactly m critical points (counted with multiplicity). Let x be a bijection from I to the set of critical points of P ; there are $m!$ choices for x . Then $x \in E$, and $P_x = P$. Therefore $P_x(x_i) = P(x_i) = x_i$, which implies that x is a fixed point of H . \square

A set of particular interest is the noninjectivity locus

$$\Delta := \{x \in E : \exists i \neq j \text{ with } x_i = x_j\}.$$

By proposition 3, the fixed points of H correspond to polynomials with fixed critical points: the fixed points in $E - \Delta$ correspond to polynomials with simple critical points, whereas the fixed points in Δ correspond to polynomials with at least one multiple critical point.

The locus Δ is a union of hyperplanes which are invariant by H ; it is a stratified space where each stratum is invariant. More precisely, denote by $\text{Part}(I)$ the set of all partitions of I , and set $\text{Part}^*(I) = \text{Part}(I) - \{I\}$. Let $\mathcal{I} \in \text{Part}^*(I)$ be the singleton partition of I : that is

$$\mathcal{I} := \{\{i\} : i \in I\}.$$

Given $\mathcal{J} \in \text{Part}^*(I)$, let $L_{\mathcal{J}} \subseteq E$ be the linear space defined by

$$L_{\mathcal{J}} := \{x \in E : x \text{ is constant on each element of } \mathcal{J}\}.$$

Note that $E = L_{\mathcal{I}}$ and the dimension of $L_{\mathcal{J}}$ is $|\mathcal{J}| - 1$. We say that $\mathcal{K} \in \text{Part}^*(I)$ is a contraction of $\mathcal{J} \in \text{Part}^*(I)$ if all elements of \mathcal{K} are unions of elements of \mathcal{J} . We denote this as $\mathcal{K} \preceq \mathcal{J}$, and if $\mathcal{K} \neq \mathcal{J}$, we use the notation $\mathcal{K} \prec \mathcal{J}$. When \mathcal{K} is a contraction of \mathcal{J} , the linear space $L_{\mathcal{K}}$ is contained in $L_{\mathcal{J}}$, and the codimension of $L_{\mathcal{K}}$ in $L_{\mathcal{J}}$ is $|\mathcal{J}| - |\mathcal{K}|$. The stratification of Δ is given by

$$\Delta = \bigcup_{\mathcal{J} \in \text{Part}^*(I)} L_{\mathcal{J}}.$$

If $x \in L_{\mathcal{J}}$, then $H(x) = P_x \circ x$ is constant on each element of \mathcal{J} . Thus $H(L_{\mathcal{J}}) \subseteq L_{\mathcal{J}}$. In particular, $h : \mathbb{P}(E) \rightarrow \mathbb{P}(E)$ restricts an endomorphism $h : \mathbb{P}(L_{\mathcal{J}}) \rightarrow \mathbb{P}(L_{\mathcal{J}})$.

Remark 2. Assume x is a fixed point of H in $E - \Delta$; then by proposition 3, P_x has m fixed critical points. The polynomial P_x is of degree $m + 1$, so there is a unique (repelling) fixed point of P_x which is not critical. Moreover, since P_x is centered and $m + 1 \geq 3$, the fixed points of P_x are also centered; this implies that this fixed point is at 0.

Consequently, the rational map $f : w \mapsto 1/P_x(1/w)$ has degree $m + 1$, a repelling fixed point at ∞ , and $m + 1$ superattracting fixed points. This map is therefore the Newton's method of a polynomial Q of degree $m + 1$.

The critical points of f are the zeroes of Q , and the zeroes of Q'' . Since f has a critical point of multiplicity m at 0 , $Q''(w) = aw^{m-1}$ for some $a \in \mathbb{C}^*$ and Q vanishes at 0 . An elementary computation then shows that

$$P_x(z) = \frac{m+1}{m}z + z^{m+1}.$$

This polynomial is indeed one with simple critical points which are fixed. So H has exactly $m!$ fixed points in $E - \Delta$.




FIGURE 2. The polynomial $P(z) = \frac{6}{5}z + z^6$ has five critical points, each of which is a superattracting fixed point of P . There is a repelling fixed point at 0 . The polynomial P is conjugate to a Newton's method for finding roots of some polynomial Q as discussed in remark 2.

3.2. The endomorphisms are postcritically finite. Our goal now is to prove that the endomorphisms $h : \mathbb{P}(L_{\mathcal{J}}) \rightarrow \mathbb{P}(L_{\mathcal{J}})$ are postcritically finite: we will identify the critical locus of $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$, and show that it is invariant. For this we will use the following observation.

Let J be a nonempty proper subset of I . Let E_J be the set of functions $J \rightarrow \mathbb{C}$ whose average is 0. There is a natural projection $\pi_J : E \rightarrow E_J$ given by

$$E \ni x \mapsto x|_J - \text{Average}(x|_J) \in E_J.$$

Let $H_J : E_J \rightarrow E_J$ be the homogeneous map constructed as above with the set J instead of the set I .

Lemma 12. *Assume $\mathcal{J} \in \text{Part}^*(I)$, $x \in L_{\mathcal{J}} - \Delta_{\mathcal{J}}$, and $J \in \mathcal{J}$. Then, as $v \rightarrow 0$ in E , we have the following expansion*

$$\pi_J \circ H(x + v) = C_J \cdot H_J \circ \pi_J(v) + \mathcal{O}(\|v\| \cdot \|\pi_J(v)\|^{|J|+1}) \quad \text{with } C_J \in \mathbb{C} - \{0\}.$$

Proof. Without loss of generality, assume that $m_J := |J| \geq 2$ since otherwise E_J has dimension 0, and the result is vacuous. Set $y := x + v$, $a_y := \text{Average}(y|_J)$ and $z := \pi_J(v) = \pi_J(y) = y|_J - a_y$. Let Q_z be the polynomial defined by

$$Q_z(t) := \frac{m_J + 1}{m_J} \sum_{j \in J} \int_{z_j}^t \prod_{i \in J} (w - z_i) dw.$$

Then

$$P'_y(t) = (m + 1) \prod_{i \in I} (t - y_i) \quad \text{and} \quad Q'_z(t - a_y) = (m_J + 1) \prod_{i \in J} (t - y_i).$$

Thus

$$P'_y(t) = Q'_z(t - a_y) \cdot R_y(t) \quad \text{with} \quad R_y(t) = \frac{m + 1}{m_J + 1} \prod_{i \in I - J} (t - y_i).$$

Since $x \in L_{\mathcal{J}} - \Delta_{\mathcal{J}}$, for all $i \in I - J$ we have $x_i \neq a_x$, thus $R_x(a_x) \neq 0$. For all $i \in J$ and $j \in J$, as $y \rightarrow x$, we have:

- $|y_j - y_i| = |z_j - z_i| = \mathcal{O}(\|z\|)$,
- $\sup_{t \in [y_i, y_j]} |Q'_z(t - a_y)| = \mathcal{O}(\|z\|^{m_J})$, and
- $\sup_{t \in [y_i, y_j]} |R_y(t) - R_x(a_x)| = \mathcal{O}(\|v\|)$.

Thus for all $i \in J$ and $j \in J$,

$$\begin{aligned} P_y(y_j) - P_y(y_i) - (Q_z(z_j) - Q_z(z_i))R_x(a_x) \\ &= \int_{y_i}^{y_j} P'_y(t) - Q'_z(t - a_y)R_x(a_x) dt \\ &= \int_{y_i}^{y_j} Q'_z(t - a_y)(R_y(t) - R_x(a_x)) dt \in \mathcal{O}(\|v\| \cdot \|z\|^{m_J+1}). \end{aligned}$$

Setting $C_J := R_x(a_x)$, we deduce that for all $j \in J$,

$$\begin{aligned} P_y(y_j) - \frac{1}{m_J + 1} \sum_{i \in J} P_y(y_i) &= C_J \cdot \left(Q_z(z_j) - \frac{1}{m_J + 1} \sum_{i \in J} Q_z(z_i) \right) + \mathcal{O}(\|v\| \cdot \|z\|^{m_J+1}) \\ &= C_J \cdot Q_z(z_j) + \mathcal{O}(\|v\| \cdot \|z\|^{m_J+1}). \end{aligned}$$

The result follows since

$$\pi_J \circ H = P_y \circ y|_J - \text{Average}(P_y \circ y|_J) \quad \text{and} \quad H_J \circ \pi_J(y) = H_J(z) = Q_z \circ z. \quad \square$$

Proposition 4. For $\mathcal{J} \in \text{Part}^*(I)$, the critical set of $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$ is

$$\Delta_{\mathcal{J}} := \bigcup_{\substack{\mathcal{K} \in \text{Part}^*(I) \\ \mathcal{K} \prec \mathcal{J}}} L_{\mathcal{K}},$$

$H(\Delta_{\mathcal{J}}) = \Delta_{\mathcal{J}}$, and $h : \mathbb{P}(L_{\mathcal{J}}) \rightarrow \mathbb{P}(L_{\mathcal{J}})$ is postcritically finite.

Proof. It is enough to prove that the critical set of $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$ is $\Delta_{\mathcal{J}}$ as the rest is an immediate consequence. Given $J_1, J_2 \in \mathcal{J}$, set

$$K := J_1 \cup J_2 \quad \text{and} \quad \mathcal{K} = \mathcal{K}(J_1, J_2) := \mathcal{J} - \{J_1, J_2\} \cup \{K\}.$$

Note that $\mathcal{K} \prec \mathcal{J}$ and $|\mathcal{K}| = |\mathcal{J}| - 1$, so that $L_{\mathcal{K}}$ has codimension 1 in $L_{\mathcal{J}}$. Choose a vector $v \in L_{\mathcal{J}} - L_{\mathcal{K}}$ so that $L_{\mathcal{J}} = L_{\mathcal{K}} \oplus \text{Span}(v)$. According to lemma 12, for all $x \in L_{\mathcal{K}}$ as $t \rightarrow 0$ we have

$$\pi_{\mathcal{K}} \circ H(x + tv) = \mathcal{O}(t^{|\mathcal{K}|+1}).$$

It follows that the Jacobian of $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$ vanishes with order at least $|\mathcal{K}|$ along $L_{\mathcal{K}}$.

Now

$$\Delta_{\mathcal{J}} = \bigcup_{\substack{J_1, J_2 \in \mathcal{J} \\ J_1 \neq J_2}} L_{\mathcal{K}(J_1, J_2)}$$

and $L_{\mathcal{K}(J_1, J_2)}$ is a critical component of $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$ with multiplicity $|J_1| + |J_2|$. So $\Delta_{\mathcal{J}}$ is contained in the critical set of $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$ as a hypersurface of total degree

$$\sum_{\substack{J_1, J_2 \in \mathcal{J} \\ J_1 \neq J_2}} |J_1| + |J_2| = (|\mathcal{J}| - 1) \cdot \sum_{J \in \mathcal{J}} |J| = (|\mathcal{J}| - 1) \cdot m.$$

Since $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$ is a homogeneous map of degree $m + 1$, its critical locus is a hypersurface of total degree $\dim(L_{\mathcal{J}}) \cdot m = (|\mathcal{J}| - 1) \cdot m$. And therefore the critical set of $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$ coincides with $\Delta_{\mathcal{J}}$ as required. \square

We now analyze the spectrum of $D_x H : T_x E \rightarrow T_x E$, where x is a fixed point of $H : E \rightarrow E$. It turns out that for the fixed points in $E - \Delta$, we have a complete understanding of this spectrum as outlined in the example below.

Example 2. We now demonstrate by way of example that the eigenvalues of $D_x H$ at a fixed point $x \in E - \Delta$ are precisely $\lambda_k := (m + 1)/k$, $k \in [1, m - 1]$, with corresponding eigenspace $\text{Span}(x^k)$.¹

According to remark 2, there is a unique polynomial

$$P(z) = \frac{m + 1}{m} z + z^{m+1}$$

which is monic and centered, with simple fixed critical points, corresponding to a fixed point of H in $E - \Delta$. Observe that if c is a critical point of P , then $P''(c) = -(m + 1)/c$.

Now if $P_t(z) = P(z) + tQ(z) + o(t)$ with $Q(z) \in \mathbb{C}_{m-1}[z]$, and if $c_t = c + tv + o(t)$ is a critical point of P_t , then

$$0 = P'_t(c_t) = P'(c) + t(P''(c)v + Q'(c)) + o(t)$$

so that

$$v = -\frac{Q'(c)}{P''(c)} = \frac{c}{m + 1} Q'(c),$$

and

$$P_t(c_t) = P(c) + tQ(c) + tP'(c)v + o(t) = c + tQ(c) + o(t).$$

Therefore $v \in T_x E$ is an eigenvector associated to the eigenvalue λ if and only if

$$\forall i \in I, \quad Q(x_i) = \lambda v_i = \frac{\lambda x_i}{m + 1} Q'(x_i).$$

This is clearly true if $Q(z) = z^k$, $\lambda = (m + 1)/k$, and $v_i = x_i^k / \lambda$.

¹Recall that $x : I \rightarrow \mathbb{C}$ is a function and x^k is the function $i \mapsto x_i^k \in \mathbb{C}$.

3.3. Fixed points are super-saddles.

Proposition 5. *Any fixed point of $H : E \rightarrow E$ is a super-saddle. More precisely, if $\mathcal{J} \in \text{Part}^*(I)$ and if $x \in L_{\mathcal{J}} - \Delta_{\mathcal{J}}$ is a fixed point of $H : E \rightarrow E$, then:*

- $T_x E = \text{Ker}(D_x H) \oplus \text{Im}(D_x H)$
- $\text{Im}(D_x H) = T_x L_{\mathcal{J}}$, and
- *the spectrum of $D_x H : T_x L_{\mathcal{J}} \rightarrow T_x L_{\mathcal{J}}$ is contained in $\mathbb{C} - \overline{\mathbb{D}}$ and therefore x is a repelling fixed point of $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$.*

Proof. According to proposition 4, x is not a critical point of $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$, thus the restriction $D_x H : T_x L_{\mathcal{J}} \rightarrow T_x L_{\mathcal{J}}$ is invertible. As a consequence the image of $D_x H$ contains $T_x L_{\mathcal{J}}$ and $\text{Ker}(D_x H) \cap T_x L_{\mathcal{J}} = \{0\}$. The kernel of the projection

$$\pi := \sum_{J \in \mathcal{J}} \pi_J : T_x E \rightarrow \bigoplus_{J \in \mathcal{J}} T_0 E_J$$

is $T_x L_{\mathcal{J}}$. According to lemma 12 $\pi_J \circ D_x H = 0$ for all $J \in \mathcal{J}$. So $\pi \circ D_x H = 0$, and the image of $D_x H$ is contained in $\text{Ker}(\pi) = T_x L_{\mathcal{J}}$. This implies that $\text{Im}(D_x H) = T_x L_{\mathcal{J}}$. In addition the codimension of $\text{Ker}(D_x H)$ is the dimension of $T_x L_{\mathcal{J}}$ and since $\text{Ker}(D_x H) \cap T_x L_{\mathcal{J}} = \{0\}$, we conclude that $T_x E = \text{Ker}(D_x H) \oplus \text{Im}(D_x H)$.

Since $H : L_{\mathcal{J}} \rightarrow L_{\mathcal{J}}$ is homogeneous of degree $m + 1$, there is an obvious eigenvalue $m + 1$ associated to the eigenspace $\text{Span}(x)$. Proposition 6 below asserts that we can endow $T_x L_{\mathcal{J}}$ with an appropriate norm so that the linear map $(D_x H)^{-1} : T_x L_{\mathcal{J}} \rightarrow T_x L_{\mathcal{J}}$ is contracting. As a consequence, the spectrum of $(D_x H)^{-1} : T_x L_{\mathcal{J}} \rightarrow T_x L_{\mathcal{J}}$ is contained in \mathbb{D} and thus, the spectrum of the linear map $D_x H : T_x L_{\mathcal{J}} \rightarrow T_x L_{\mathcal{J}}$ is contained in $\mathbb{C} - \overline{\mathbb{D}}$. \square

By definition E is a subspace of \mathbb{C}^I of codimension 1. It is the kernel of the linear form

$$e^I : E \ni v \mapsto \frac{1}{|I|} \sum_{i \in I} v_i \in \mathbb{C}.$$

As a consequence E^* may be identified with the quotient $(\mathbb{C}^I)^*/\text{Span}(e^I)$. Let $\mathbf{1} : I \rightarrow \mathbb{C}$ be the function which is constant and equal to 1. Since $\mathbb{C}^I = E \oplus \text{Span}(\mathbf{1})$, the dual space E^* may also be identified with the orthogonal space

$$(\text{Span}(\mathbf{1}))^\perp = \{\alpha \in (\mathbb{C}^I)^* : \alpha(\mathbf{1}) = 0\}.$$

More generally, given $J \subseteq I$ let $e^J \in (\mathbb{C}^I)^*$ be the linear form defined by

$$e^J(v) = \frac{1}{|J|} \sum_{j \in J} v_j.$$

Note that for $\mathcal{J} \in \text{Part}^*(I)$,

$$E = L_{\mathcal{J}} \oplus \bigcap_{J \in \mathcal{J}} \text{Ker}(e^J)$$

and therefore $L_{\mathcal{J}}^*$ may be identified with

$$(\text{Span}(\mathbf{1}))^\perp \cap \text{Span}(e^J; J \in \mathcal{J}) = \left\{ \sum_{J \in \mathcal{J}} \lambda_J e^J : \lambda_J \in \mathbb{C} \text{ and } \sum_{J \in \mathcal{J}} \lambda_J = 0 \right\}.$$

If $\alpha = \sum_{J \in \mathcal{J}} \lambda_J e^J \in L_{\mathcal{J}}^*$ then the pairing with $v \in L_{\mathcal{J}}$ is given by

$$\alpha(v) = \sum_{J \in \mathcal{J}} \lambda_J v_J \quad \text{with} \quad v(J) = \{v_J\}.$$

Given $x \in L_{\mathcal{J}} - \Delta_{\mathcal{J}}$ we will now identify $(T_x L_{\mathcal{J}})^*$ with a space of quadratic differentials, equip this space with an appropriate norm then identify the transpose of $(D_x H)^{-1} : T_x L_{\mathcal{J}} \rightarrow T_x L_{\mathcal{J}}$, and finally show that this transpose is strictly contracting. So for $J \in \mathcal{J}$, let q_J be the quadratic differential defined on \mathbb{C} by

$$q_J = \frac{dz^2}{z - x_J}.$$

Set

$$\mathcal{Q}_x := \left\{ \sum_{J \in \mathcal{J}} \lambda_J q_J : \sum_{J \in \mathcal{J}} \lambda_J = 0 \right\}.$$

A quadratic differential $q = \sum \lambda_J q_J \in \mathcal{Q}_x$ may be paired with a tangent vector $v \in T_x L_{\mathcal{J}}$ as follows

$$\langle q, v \rangle := \sum_{J \in \mathcal{J}} \lambda_J v_J = \sum_{J \in \mathcal{J}} \text{Res}_{x_J}(q \cdot \xi_v)$$

where ξ_v is any holomorphic vector field near $x(I)$ which takes the value v_J at x_J . According to the previous discussion, this pairing gives an identification of $(T_x L_{\mathcal{J}})^*$ with \mathcal{Q}_x .

Choose R large enough so that $P_x^{-1}(D_R)$ is compactly contained in D_R , where D_R is the disk centered at 0 of radius R . We equip \mathcal{Q}_x with the L^1 norm

$$\|q\| := \int_{D_R} |q|.$$

Proposition 6. *Assume $x \in L_{\mathcal{J}} - \Delta_{\mathcal{J}}$. The transpose of the linear mapping $(D_x H)^{-1} : T_x L_{\mathcal{J}} \rightarrow T_x L_{\mathcal{J}}$ is identified with the push-forward operator*

$$(P_x)_* : \mathcal{Q}_x \ni q \mapsto \sum g^* q \in \mathcal{Q}_x$$

where g ranges over the inverse branches of P_x . In addition,

$$\forall q \in \mathcal{Q}_x \quad \|(P_x)_* q\| < \|q\|.$$

Proof. The space \mathcal{Q}_x is the set of meromorphic quadratic differentials on $\mathbb{P}^1(\mathbb{C})$ which are holomorphic outside $x(I)$, have at most simple poles along $x(I)$ and at most a double pole at ∞ . Set $P := P_x$. If $q \in \mathcal{Q}_x$ then $P_* q$ is a meromorphic quadratic differential on $\mathbb{P}^1(\mathbb{C})$. Since q has at most a double pole at ∞ , $P_* q$ also has at most a double pole at ∞ . The other poles of $P_* q$ are simple and contained in $P(x(I))$ union the critical value set of P , that is $x(I)$. This shows that P_* maps \mathcal{Q}_x to \mathcal{Q}_x . In addition

$$\|P_* q\| = \int_{D_R} \left| \sum g^* q \right| \leq \int_{D_R} \sum |g^* q| = \int_{P^{-1}(D_R)} |q| < \int_{D_R} |q| = \|q\|.$$

Therefore we only need to prove that for all $v \in T_x L_{\mathcal{J}}$ and all $q \in \mathcal{Q}_x$

$$\langle q, v \rangle = \langle P_* q, D_x H(v) \rangle.$$

Fix $v \in T_x L_{\mathcal{J}}$ and $q \in \mathcal{Q}_x$. Let U be the complement in D_R of pairwise disjoint closed disks centered at the points of $x(I)$, and contained in D_R . If ξ is a C^∞

vector field on \mathbb{C} which is holomorphic outside U , vanishes outside D_R , and satisfies $\xi \circ x = v$, then

$$\langle q, v \rangle = \sum_{J \in \mathcal{J}} \text{Res}_{x_J}(q \cdot \xi) = -\frac{1}{2\pi i} \int_{\partial U} q \cdot \xi \stackrel{\text{Stokes}}{=} \frac{1}{2\pi i} \int_U q \cdot \bar{\partial} \xi = \frac{1}{2\pi i} \int_{\mathbb{C}} q \cdot \bar{\partial} \xi.$$

With an abuse of notation set

$$x_t := x + tv, \quad P_t := P_{x_t}, \quad \text{and} \quad y_t := H(x_t) = P_t \circ x_t.$$

Then,

$$\dot{x} := \frac{\partial x_t}{\partial t} \Big|_{t=0} = v \quad \text{and} \quad \dot{y} := \frac{\partial y_t}{\partial t} \Big|_{t=0} = D_x H(v).$$

In addition, the critical point set of P_t is $x_t(I)$ and the critical value set of P_t is $y_t(I)$. Let $(\varphi_t : \mathbb{C} \rightarrow \mathbb{C})_{t \in (-\epsilon, \epsilon)}$ be an analytic family of C^∞ diffeomorphisms such that

- $\varphi_0 = \text{id}$,
- φ_t is the identity outside D_R ,
- φ_t is holomorphic outside U and
- $y_t = \varphi_t \circ y$.

Note that

$$\dot{\varphi} := \frac{\partial \varphi_t}{\partial t} \Big|_{t=0}$$

is a C^∞ vector field on \mathbb{C} which is holomorphic outside U , vanishes outside D_R , and satisfies $\dot{\varphi} \circ x = D_x H(v)$. Since φ_t follows the critical value set of P_t we can lift the diffeomorphisms $\varphi_t : \mathbb{C} \rightarrow \mathbb{C}$ to diffeomorphisms $\psi_t : \mathbb{C} \rightarrow \mathbb{C}$ so that $\psi_0 = \text{id}$ and the following diagram commutes

$$\begin{array}{ccc} (\mathbb{C}, x(I)) & \xrightarrow{\psi_t} & (\mathbb{C}, x_t(I)) \\ \downarrow P & & \downarrow P_t \\ (\mathbb{C}, y(I)) & \xrightarrow{\varphi_t} & (\mathbb{C}, y_t(I)) \end{array}$$

Then,

$$\dot{\psi} := \frac{\partial \psi_t}{\partial t} \Big|_{t=0}$$

is a C^∞ vector field on \mathbb{C} which is holomorphic outside U , vanishes outside D_R , and satisfies $\dot{\psi} \circ x = v$. In addition, the infinitesimal Beltrami differentials $\bar{\partial} \dot{\varphi}$ and $\bar{\partial} \dot{\psi}$ satisfy the relation

$$\bar{\partial} \dot{\psi} = P^*(\bar{\partial} \dot{\varphi}).$$

Therefore

$$\langle q, v \rangle = \int_{\mathbb{C}} q \cdot \bar{\partial} \dot{\psi} = \int_{\mathbb{C}} q \cdot P^*(\bar{\partial} \dot{\varphi}) = \int_{\mathbb{C}} P_* q \cdot \bar{\partial} \dot{\varphi} = \langle P_* q, D_x H(v) \rangle$$

as required. \square




FIGURE 3. On the left is a real slice of the immediate basin $\mathcal{B}_{a_{\mathcal{J}}}(h)$ in the case $|I| = 4$ and $\mathcal{J} = \{J_1, J_2\}$ with $|J_1| = 3$ and $|J_2| = 1$. The critical set $\mathbb{P}(\Delta)$ contains three lines passing through $a_{\mathcal{J}}$. Almost every orbit in $\mathcal{B}_{a_{\mathcal{J}}}(h)$ converges to $a_{\mathcal{J}}$ tangentially along one of these lines; the immediate basin is colored accordingly. Since $|J_2| = 1$, $E_{J_2} = \{0\}$ and $H_{J_1} \oplus H_{J_2} = H_{J_1}$. According to proposition 7 below, there is a Böttcher coordinate $\Phi : \mathcal{B}_{a_{\mathcal{J}}}(h) \rightarrow \mathcal{B}_0(H_{J_1})$. On the right is a real slice of the immediate basin $\mathcal{B}_0(H_{J_1})$. The isomorphism Φ respects the coloring.

3.4. Superattracting fixed points of $h : \mathbb{P}(E) \rightarrow \mathbb{P}(E)$. If $\mathcal{J} \in \text{Part}^*(I)$ has cardinality 2, then $L_{\mathcal{J}}$ has dimension 1 and its image in $\mathbb{P}(E)$ is a point $a_{\mathcal{J}}$. This is a superattracting fixed point for $h : \mathbb{P}(E) \rightarrow \mathbb{P}(E)$.

We will now show that we can apply theorems 1 and 2. Note that $H_{J_1} \oplus H_{J_2} : E_{J_1} \oplus E_{J_2} \rightarrow E_{J_1} \oplus E_{J_2}$ is a quasihomogeneous map of bidegree $(|J_1|, |J_2|)$.

Proposition 7. *Let $\mathcal{J} := \{J_1, J_2\} \in \text{Part}^*(I)$ and $a_{\mathcal{J}}$ be the image of $L_{\mathcal{J}}$ in $\mathbb{P}(E)$. Then there is an analytic isomorphism*

$$\Phi : \mathcal{B}_{a_{\mathcal{J}}}(h) \longrightarrow \mathcal{B}_0(H_{J_1} \oplus H_{J_2})$$

conjugating h to $H_{J_1} \oplus H_{J_2}$.

Proof. The analytic map $h : \mathbb{P}(E) \rightarrow \mathbb{P}(E)$ is proper, and has a superattracting fixed point at $a_{\mathcal{J}}$. We will prove that there is a local isomorphism

$$\Phi : (\mathbb{P}(E), a_{\mathcal{J}}) \rightarrow (E_{J_1} \oplus E_{J_2}, 0)$$

conjugating h to $H_{J_1} \oplus H_{J_2}$. Such a Φ automatically maps the germ of the postcritical set of h at $a_{\mathcal{J}}$, i.e. the germ of $\mathbb{P}(\Delta)$ at $a_{\mathcal{J}}$, to the germ of the postcritical set of $H_{J_1} \oplus H_{J_2}$ at 0. So near $a_{\mathcal{J}}$, Φ maps the postcritical set of $h : \mathcal{B}_{a_{\mathcal{J}}}(h) \rightarrow \mathcal{B}_{a_{\mathcal{J}}}(h)$, that is $\mathcal{B}_{a_{\mathcal{J}}}(h) \cap \mathbb{P}(\Delta)$, to the postcritical set of $H_{J_1} \oplus H_{J_2}$. The result then follows from theorem 2.

To prove that there is a local conjugacy, we use theorem 1. Choose $x \in L_{\mathcal{J}}$ and let $L_{\mathcal{J}}^{\perp} \subset E$ be a subspace of codimension 1 such that $E = L_{\mathcal{J}} \oplus L_{\mathcal{J}}^{\perp}$. Let $\sigma : (\mathbb{P}(E), a_{\mathcal{J}}) \rightarrow (E, x)$ be the local section with image in $x + L_{\mathcal{J}}^{\perp}$. Then the composition $\pi \circ \sigma : (\mathbb{P}(E), a_{\mathcal{J}}) \rightarrow (E_{J_1} \oplus E_{J_2}, 0)$ is a local isomorphism. It conjugates

$h : (\mathbb{P}(E), a_{\mathcal{J}}) \rightarrow (\mathbb{P}(E), a_{\mathcal{J}})$ to an analytic germ $F : (E_{J_1} \oplus E_{J_2}, 0) \rightarrow (E_{J_1} \oplus E_{J_2}, 0)$. It follows from lemma 12 that F has an adapted superattracting fixed point with quasihomogeneous part $c_1 H_{J_1} \oplus c_2 H_{J_2}$ for some constants c_1 and c_2 in \mathbb{C}^* .

In addition $v := v_1 + v_2$ with $v_1 \in E_{J_1}$ and $v_2 \in E_{J_2}$ is contained in the postcritical set of F near 0 if and only if there is a $w \in L_{\mathcal{J}}^{\perp}$ with $\pi(w) = v$ and $x + w \in \Delta$. Since $x(J_1) \neq x(J_2)$, if v is sufficiently close to 0, the sets $(x + w)(J_1)$ and $(x + w)(J_2)$ are disjoint. In that case, $x + w \in \Delta$ if and only if $v_1 : J_1 \rightarrow \mathbb{C}$ or $v_2 : J_2 \rightarrow \mathbb{C}$ is not injective. As a consequence $\lambda_1 v_1 + \lambda_2 v_2$ is contained in the postcritical set of F for all $(\lambda_1, \lambda_2) \in \mathbb{C}^2$ and the vector fields ϑ_{J_1} and ϑ_{J_2} are tangent to the postcritical set of F near 0.

According to theorem 1, there is a local isomorphism conjugating F near 0 to $H_{J_1} \oplus H_{J_2}$ near 0. The existence of a Böttcher coordinate

$$\Phi : (\mathbb{P}(E), a_{\mathcal{J}}) \rightarrow (E_{J_1} \oplus E_{J_2}, 0)$$

conjugating the map $F : (\mathbb{P}(E), a_{\mathcal{J}}) \rightarrow (\mathbb{P}(E), a_{\mathcal{J}})$ to the quasihomogeneous map $H_{J_1} \oplus H_{J_2} : (E_{J_1} \oplus E_{J_2}, 0) \rightarrow (E_{J_1} \oplus E_{J_2}, 0)$ follows immediately. \square

4. QUESTIONS FOR FURTHER STUDY

Theorem 2 asserts that when $a_{\mathcal{J}} \in \mathbb{P}(E)$ is a superattracting fixed point of $h : \mathbb{P}(E) \rightarrow \mathbb{P}(E)$ then there is a Böttcher coordinate $\Phi : \mathcal{B}_a(h) \rightarrow \mathcal{B}_0(H_{\mathcal{J}})$ with $H_{\mathcal{J}} := H_{J_1} \oplus H_{J_2} : E_{J_1} \oplus E_{J_2} \rightarrow E_{J_1} \oplus E_{J_2}$. The boundary of $\mathcal{B}_0(H_{\mathcal{J}})$ is a topological sphere of real dimension $2m - 5$. Does the inverse $\Phi^{-1} : \mathcal{B}_0(H_{\mathcal{J}}) \rightarrow \mathcal{B}_a(h)$ extend continuously to the boundary of $\mathcal{B}_0(H_{\mathcal{J}})$? Is the boundary of $\mathcal{B}_a(h)$ topologically the quotient of a sphere by an equivalence relation? How could such an equivalence relation be described?

In dimension one, a global Böttcher coordinate gives rise to a dynamical foliation of the immediate basin by rays. What is the higher dimensional analog of these rays?

In dimension one, if two germs with a superattracting fixed point are topologically conjugate, then this topological conjugacy can be promoted to an analytic conjugacy via a pullback argument. Can one give a topological and/or analytic classification of germs having a superattracting fixed point in higher dimensions? Do these classifications coincide?

In section 3, we applied theorems 1 and 2 to a superattracting fixed point $a_{\mathcal{J}}$, which was the image of $L_{\mathcal{J}}$ in $\mathbb{P}(E)$, where $|\mathcal{J}| = 2$. What happens for $|\mathcal{J}| > 2$? In this case, $\dim(L_{\mathcal{J}}) = |\mathcal{J}| - 1$, and the image in $\mathbb{P}(E)$ will be a projective space of dimension $|\mathcal{J}| - 2$. Is this projective space an attractor (in the sense of [M])?

If $|\mathcal{J}| > 2$ and if $a \in \mathbb{P}(L_{\mathcal{J}})$ is a fixed point, then according to proposition 5 the spectrum of $D_a h : T_a \mathbb{P}(L_{\mathcal{J}}) \rightarrow T_a \mathbb{P}(L_{\mathcal{J}})$ belongs to $\mathbb{C} - \overline{\mathbb{D}}$, and $\mathbb{P}(L_{\mathcal{J}})$ is the unstable manifold of a . The unique additional eigenvalue of $D_a h : T_a \mathbb{P}(E) \rightarrow T_a \mathbb{P}(E)$ is 0. What is the structure of the associated superstable manifold

$$W^s(a) := \left\{ b \in \mathbb{P}(E) : h^{\circ n}(b) \xrightarrow{n \rightarrow \infty} a \right\}?$$

Is it a smooth analytic submanifold of $\mathbb{P}(E)$? Is it dynamically parameterized by the attracting basin of the quasihomogeneous map

$$\bigoplus_{J \in \mathcal{J}} H_J : \bigoplus_{J \in \mathcal{J}} E_J \rightarrow \bigoplus_{J \in \mathcal{J}} E_J?$$

The maps to which we applied our theorems in section 3 were postcritically finite. Are there examples (apart from the quasihomogeneous maps themselves) which are algebraic, and not postcritically finite but admit a Böttcher coordinate? The converse is false: consider the map $F : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ given by $F : (x, y) \mapsto (x^2 - y^3, y^2)$. One can verify that F is postcritically finite. The derivative D_0F is nilpotent so that $F^{\circ 2}$ has a superattracting fixed point at 0. In addition

$$F^{\circ 2}(x, y) = (y^2 + o(y^2), y^4 + o(y^4)).$$

The map $F^{\circ 2}$ cannot be locally conjugate to the map $(x, y) \mapsto (y^2, y^4)$ as this map is not open.

REFERENCES

- [BEKP] X. BUFF, A. EPSTEIN, S. KOCH & K. PILGRIM *On Thurston's Pullback Map* Complex Dynamics, Families and Friends, D. Schleicher, AK Peters, 2009.
- [DH] A. DOUADY & J.H. HUBBARD *Etude dynamique des polynômes complexes* I & II, Publ. Math. d'Orsay, 1984-85.
- [F] C. FAVRE *Classification of 2-dimensional contracting rigid germs and Kato surfaces: I*, J. Math. Pures Appl. (9) 79 (5): 2000, 475-514.
- [FJ] C. FAVRE & M. JONSSON *Eigenvaluations*, Ann. Sci. École Norm. Sup. (4) 40: 2007, no. 2, 309-349.
- [FS] J. FORNÆSS & N. SIBONY *Critically Finite Rational Maps on \mathbb{P}^2* , Proceedings of the Madison Symposium honouring Walter Rudin, AMS series in Contemporary Mathematics: (1992) 245-260.
- [HP] J.H. HUBBARD & P. PAPADOPOL *Superattractive fixed points in \mathbb{C}^n* , Indiana Univ. Math. J. (43): 1994, no. 1, 321-365.
- [K] S. KOCH *A new link between Teichmüller theory and complex dynamics*, PhD thesis, Cornell University (2008).
- [M] J. MILNOR *On the concept of attractor*, Comm. Math. Phys. 99 (1985), no. 2, 177-195.
- [Ue1] T. UEDA *Complex dynamical systems on projective space*, Sūrikaiseikikenkyūsho Kōkyūroku (814): 1992 169-186.
- [Ue2] T. UEDA *Critically finite maps on projective spaces*, Sūrikaiseikikenkyūsho Kōkyūroku (1097): 1999 132-138.
- [Us] S. USHIKI *Böttcher's theorem and super-stable manifolds for multidimensional complex dynamical systems*, preprint.

E-mail address: `xavier.buff@math.univ-toulouse.fr`

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UNIVERSITÉ PAUL SABATIER, 118, ROUTE DE NARBONNE, 31062 TOULOUSE CEDEX, FRANCE

E-mail address: `adame@maths.warwick.ac.uk`

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UNITED KINGDOM

E-mail address: `kochs@math.harvard.edu`

DEPARTMENT OF MATHEMATICS, SCIENCE CENTER, HARVARD UNIVERSITY, CAMBRIDGE MA 02138, UNITED STATES