

Convex concentration inequalities and forward-backward stochastic calculus

Thierry KLEIN ^{a)} Yutao MA ^{b) c)} Nicolas PRIVAULT ^{c)}

January 10, 2006

Abstract

Given $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ respectively a forward and a backward martingale with jumps and continuous parts, we prove that $E[\phi(M_t + M_t^*)]$ is non-decreasing in t when ϕ is a convex function, provided the local characteristics of $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ satisfy some comparison inequalities. We deduce convex concentration inequalities and deviation bounds for random variables admitting a predictable representation in terms of a Brownian motion and a non-necessarily independent jump component.

Keywords: Convex concentration inequalities, forward-backward stochastic calculus, deviation inequalities, Clark formula, Brownian motion, jump processes.

2000 MR Subject Classification : 60F99, 60F10, 60H07, 39B62.

1 Introduction

Two random variables F and G satisfy a convex concentration inequality if

$$\mathbb{E}[\phi(F)] \leq \mathbb{E}[\phi(G)] \quad (1.1)$$

for all convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$. By a classical argument, the application of (1.1) to $\phi(x) = \exp(\lambda x)$, $\lambda > 0$, entails the deviation bound

$$P(F \geq x) \leq \inf_{\lambda > 0} \mathbb{E}[e^{\lambda(F-x)} 1_{\{F \geq x\}}] \leq \inf_{\lambda > 0} \mathbb{E}[e^{\lambda(F-x)}] \leq \inf_{\lambda > 0} \mathbb{E}[e^{\lambda(G-x)}], \quad (1.2)$$

a) Institut de Mathématiques - L.S.P., Université Paul Sabatier, 31062 Toulouse, France.
tklein@cict.fr

b) School of Mathematics and Statistics, Wuhan University, 430072 Hubei, P.R. China.
yutao.ma@univ-lr.fr

c) Département de Mathématiques, Université de La Rochelle, 17042 La Rochelle, France.
nicolas.privault@univ-lr.fr

$x > 0$, hence the deviation probabilities for F can be estimated via the Laplace transform of G , see [2], [13] for more results on this topic.

On the other hand, if F is more convex concentrated than G then $E[F] = E[G]$ as follows from taking successively $\phi(x) = x$ and $\phi(x) = -x$, and applying the convex concentration inequality to $\phi(x) = x \log x$ we get

$$\begin{aligned} \text{Ent}[F] &= \mathbb{E}[F \log F] - \mathbb{E}[F] \log \mathbb{E}[F] \\ &= \mathbb{E}[F \log F] - \mathbb{E}[G] \log \mathbb{E}[G] \\ &\leq \mathbb{E}[G \log G] - \mathbb{E}[G] \log \mathbb{E}[G] \\ &= \text{Ent}[G], \end{aligned}$$

hence a logarithmic Sobolev inequality of the form $\text{Ent}[G] \leq \mathcal{E}(G, G)$ implies

$$\text{Ent}[F] \leq \mathcal{E}(G, G).$$

In this paper we obtain convex concentration inequalities for the sum $M_t + M_t^*$, $t \in \mathbb{R}_+$, of a forward and a backward martingale with jumps and continuous parts. Namely we prove that $M_t + M_t^*$ is more concentrated than $M_s + M_s^*$ if $t \geq s \geq 0$, i.e.

$$\mathbb{E}[\phi(M_t + M_t^*)] \leq \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t,$$

for all convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$, provided the local characteristics of $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ satisfy the comparison inequalities assumed in Theorem 3.2 below. If further $\mathbb{E}[M_t^* | \mathcal{F}_t^M] = 0$, $t \in \mathbb{R}_+$, where $(\mathcal{F}_t^M)_{t \in \mathbb{R}_+}$ denotes the filtration generated by $(M_t)_{t \in \mathbb{R}_+}$, then Jensen's inequality yields

$$\mathbb{E}[\phi(M_t)] \leq \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t,$$

and if in addition we have $M_0 = 0$, then

$$\mathbb{E}[\phi(M_T)] \leq \mathbb{E}[\phi(M_0^*)], \quad T \geq 0. \tag{1.3}$$

In other terms, we will show that a random variable F is more concentrated than M_0^* :

$$\mathbb{E}[\phi(F - E[F])] \leq \mathbb{E}[\phi(M_0^*)],$$

provided certain assumptions are made on the processes appearing in the predictable representation of $F - \mathbb{E}[F] = M_T$ in terms of a point process and a Brownian motion. Consider for example a random variable F represented as

$$F = \mathbb{E}[F] + \int_0^{+\infty} H_t dW_t + \int_0^{+\infty} J_t (dZ_t - \lambda_t dt),$$

where $(Z_t)_{t \in \mathbb{R}_+}$ is a point process with compensator $(\lambda_t)_{t \in \mathbb{R}_+}$, $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion, and $(H_t)_{t \in \mathbb{R}_+}$, $(J_t)_{t \in \mathbb{R}_+}$ are predictable square-integrable processes satisfying $J_t \leq k$, $dPdt$ -a.e., and

$$\int_0^{+\infty} |H_t|^2 dt \leq \beta^2, \quad \text{and} \quad \int_0^{+\infty} |J_t|^2 \lambda_t dt \leq \alpha^2, \quad P - a.s.$$

By applying (1.3) or Theorem 4.1-*ii*) below to forward and backward martingales of the form

$$M_t = \mathbb{E}[F] + \int_0^t H_u dW_u + \int_0^t J_u (dZ_u - \lambda_u du), \quad t \in \mathbb{R}_+,$$

and

$$M_t^* = \hat{W}_{\beta^2} - \hat{W}_{V^2(t)} + k(\hat{N}_{\alpha^2/k^2} - \hat{N}_{U^2(t)/k^2}) - (\alpha^2 - U^2(t))/k, \quad t \in \mathbb{R}_+,$$

where $(\hat{W}_t)_{t \in \mathbb{R}_+}$, $(\hat{N}_t)_{t \in \mathbb{R}_+}$, are a Brownian motion and a left-continuous standard Poisson process, $\beta \geq 0$, $\alpha \geq 0$, $k > 0$, and $(V(t))_{t \in \mathbb{R}_+}$, $(U(t))_{t \in \mathbb{R}_+}$ are suitable random time changes, it will follow in particular that F is more concentrated than

$$M_0^* = \hat{W}_{\beta^2} + k\hat{N}_{\alpha^2/k^2} - \alpha^2/k,$$

i.e.

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi(\hat{W}_{\beta^2} + k\hat{N}_{\alpha^2/k^2} - \alpha^2/k) \right] \quad (1.4)$$

for all \mathcal{C}^2 convex function ϕ such that ϕ'' is non-decreasing.

From (1.2) and (1.4) we get

$$P(F - \mathbb{E}[F] \geq x) \leq \inf_{\lambda > 0} \exp \left(\frac{\alpha^2}{k^2} (e^{\lambda k} - \lambda k - 1) + \frac{\beta^2 \lambda^2}{2} - \lambda x \right),$$

i.e.

$$P(F - \mathbb{E}[F] \geq x) \leq \exp \left(\frac{x}{k} - \frac{\beta^2 \lambda_0(x)}{2k} (2 - k\lambda_0(x)) - (x + \alpha^2/k) \lambda_0(x) \right),$$

where $\lambda_0(x) > 0$ is the unique solution of

$$e^{k\lambda_0(x)} + \frac{k\lambda_0(x)\beta^2}{\alpha^2} - 1 = \frac{kx}{\alpha^2}.$$

When $H_t = 0$, $t \in \mathbb{R}_+$, we can take $\beta = 0$, then $\lambda_0(x) = k^{-1} \log(1 + xk/\alpha^2)$ and this implies the Poisson tail estimate

$$P(F - \mathbb{E}[F] \geq y) \leq \exp\left(\frac{y}{k} - \left(\frac{y}{k} + \frac{\alpha^2}{k^2}\right) \log\left(1 + \frac{ky}{\alpha^2}\right)\right), \quad y > 0. \quad (1.5)$$

Such an inequality has been proved in [1], [17], using (modified) logarithmic Sobolev inequalities and the Herbst method when $Z_t = N_t$, $t \in \mathbb{R}_+$, is a Poisson process, under different hypotheses on the predictable representation of F via the Clark formula, cf. Section (6). When $J_t = \lambda_t = 0$, $t \in \mathbb{R}_+$, we recover classical Gaussian estimates which can be independently obtained from the expression of continuous martingales as time-changed Brownian motions.

We proceed as follows. In Section 3 we present convex concentration inequalities for martingales. In Sections 4 and 5 these results are applied to derive convex concentration inequalities with respect to Gaussian and Poisson distributions. In Section 6 we consider the case of predictable representations obtained from the Clark formula. The proofs of the main results are formulated using forward/backward stochastic calculus and arguments of [8]. Section 7 deals with an application to normal martingales, and in the appendix (Section 8) we prove the forward-backward Itô type change of variable formula which is used in the proof of our convex concentration inequalities.

2 Notation

Let (Ω, \mathcal{F}, P) be a probability space equipped with an increasing filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ and a decreasing filtration $(\mathcal{F}_t^*)_{t \in \mathbb{R}_+}$. Consider $(M_t)_{t \in \mathbb{R}_+}$ an \mathcal{F}_t -forward martingale and $(M_t^*)_{t \in \mathbb{R}_+}$ an \mathcal{F}_t^* -backward martingale. We assume that $(M_t)_{t \in \mathbb{R}_+}$ has right-continuous paths with left limits, and that $(M_t^*)_{t \in \mathbb{R}_+}$ has left-continuous paths with right limits. Denote respectively by $(M_t^c)_{t \in \mathbb{R}_+}$ and $(M_t^{*c})_{t \in \mathbb{R}_+}$ the continuous parts of $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$, and by

$$\Delta M_t = M_t - M_{t-}, \quad \Delta^* M_t^* = M_t^* - M_{t+}^*,$$

their forward and backward jumps. The processes $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ have jump measures

$$\mu(dt, dx) = \sum_{s>0} 1_{\{\Delta M_s \neq 0\}} \delta_{(s, \Delta M_s)}(dt, dx),$$

and

$$\mu^*(dt, dx) = \sum_{s>0} 1_{\{\Delta^* M_s^* \neq 0\}} \delta_{(s, \Delta^* M_s^*)}(dt, dx),$$

where $\delta_{(s,x)}$ denotes the Dirac measure at $(s, x) \in \mathbb{R}_+ \times \mathbb{R}$. Denote by $\nu(dt, dx)$ and $\nu^*(dt, dx)$ the $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ and $(\mathcal{F}_t^*)_{t \in \mathbb{R}_+}$ -dual predictable projections of $\mu(dt, dx)$ and $\mu^*(dt, dx)$, i.e.

$$\int_0^t \int_{-\infty}^{\infty} f(s, x)(\mu(ds, dx) - \nu(ds, dx)) \quad \text{and} \quad \int_t^{\infty} \int_{-\infty}^{\infty} g(s, x)(\mu^*(ds, dx) - \nu^*(ds, dx))$$

are respectively \mathcal{F}_t -forward and \mathcal{F}_t^* -backward local martingales for all sufficiently integrable \mathcal{F}_t -predictable, resp. \mathcal{F}_t^* -predictable, process f , resp. g . The quadratic variations $([M, M])_{t \in \mathbb{R}_+}$, $([M^*, M^*])_{t \in \mathbb{R}_+}$ are defined as the limits in uniform convergence in probability

$$[M, M]_t = \lim_{n \rightarrow \infty} \sum_{i=1}^n |M_{t_i^n} - M_{t_{i-1}^n}|^2,$$

and

$$[M^*, M^*]_t = \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} |M_{t_i^n}^* - M_{t_{i+1}^n}^*|^2,$$

for all refining sequences $\{0 = t_0^n \leq t_1^n \leq \dots \leq t_{k_n}^n = t\}$, $n \geq 1$, of partitions of $[0, t]$ tending to the identity. We then let $M_t^d = M_t - M_t^c$, $M_t^{*d} = M_t^* - M_t^{*c}$,

$$[M^d, M^d]_t = \sum_{0 < s \leq t} |\Delta M_s|^2, \quad [M^{*d}, M^{*d}]_t = \sum_{0 \leq s < t} |\Delta^* M_s^*|^2,$$

and

$$\langle M^c, M^c \rangle_t = [M, M]_t - [M^d, M^d]_t, \quad \langle M^{*c}, M^{*c} \rangle_t = [M^*, M^*]_t - [M^{*d}, M^{*d}]_t,$$

$t \in \mathbb{R}_+$. Note that $([M, M]_t)_{t \in \mathbb{R}_+}$, $(\langle M, M \rangle_t)_{t \in \mathbb{R}_+}$, $([M^*, M^*]_t)_{t \in \mathbb{R}_+}$ and $(\langle M^*, M^* \rangle_t)_{t \in \mathbb{R}_+}$ are \mathcal{F}_t -adapted, but $([M^*, M^*]_t)_{t \in \mathbb{R}_+}$ and $(\langle M^*, M^* \rangle_t)_{t \in \mathbb{R}_+}$ are not \mathcal{F}_t^* -adapted. The pairs

$$(\nu(dt, dx), \langle M^c, M^c \rangle) \quad \text{and} \quad (\nu^*(dt, dx), \langle M^{*c}, M^{*c} \rangle)$$

are called the local characteristics of $(M_t)_{t \in \mathbb{R}_+}$, cf. [6] in the forward case. Denote by $(\langle M^d, M^d \rangle_t)_{t \in \mathbb{R}_+}$, $(\langle M^{*d}, M^{*d} \rangle_t)_{t \in \mathbb{R}_+}$ the quadratic variations of $(M_t^d)_{t \in \mathbb{R}_+}$, $(M_t^{*d})_{t \in \mathbb{R}_+}$, with

$$d\langle M^d, M^d \rangle_t = \int_{\mathbb{R}} |x|^2 \nu(dt, dx) \quad \text{and} \quad d\langle M^{*d}, M^{*d} \rangle_t = \int_{\mathbb{R}} |x|^2 \nu^*(dt, dx).$$

The quadratic variations $(\langle M, M \rangle_t)_{t \in \mathbb{R}_+}$, $(\langle M^*, M^* \rangle_t)_{t \in \mathbb{R}_+}$ of $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ satisfy

$$\langle M, M \rangle_t = \langle M^c, M^c \rangle_t + \langle M^d, M^d \rangle_t, \quad \text{and} \quad \langle M^*, M^* \rangle_t = \langle M^{*c}, M^{*c} \rangle_t + \langle M^{*d}, M^{*d} \rangle_t,$$

$t \in \mathbb{R}_+$.

3 Convex concentration inequalities for martingales

In the sequel, $(M_t)_{t \in \mathbb{R}_+}$ is an \mathcal{F}_t^* -adapted, \mathcal{F}_t -forward martingale, and $(M_t^*)_{t \in \mathbb{R}_+}$ is an \mathcal{F}_t -adapted, \mathcal{F}_t^* -backward martingale, whose characteristics have the form

$$\nu(du, dx) = \nu_u(dx)du \quad \text{and} \quad \nu^*(du, dx) = \nu_u^*(dx)du, \quad (3.1)$$

and

$$d\langle M^c, M^c \rangle_t = |H_t|^2 dt, \quad \text{and} \quad d\langle M^{*c}, M^{*c} \rangle_t = |H_t^*|^2 dt, \quad (3.2)$$

where $(H_t)_{t \in \mathbb{R}_+}$, $(H_t^*)_{t \in \mathbb{R}_+}$, are respectively predictable with respect to $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ and $(\mathcal{F}_t^*)_{t \in \mathbb{R}_+}$. Recall the following Lemma.

Lemma 3.1. *Let m_1, m_2 be two measures on \mathbb{R} such that $m_1([x, \infty)) \leq m_2([x, \infty)) < \infty$, $x \in \mathbb{R}$. Then for all non-decreasing and m_1, m_2 -integrable function f on \mathbb{R} we have*

$$\int_{-\infty}^{\infty} f(x) m_1(dx) \leq \int_{-\infty}^{\infty} f(x) m_2(dx).$$

If m_1, m_2 are probability measures then the above property corresponds to stochastic domination for random variables of respective laws m_1, m_2 .

Theorem 3.2. *Let*

$$\bar{\nu}_u(dx) = x\nu_u(dx), \quad \bar{\nu}_u^*(dx) = x\nu_u^*(dx), \quad u \in \mathbb{R}_+,$$

and assume that:

i) ν_u and ν_u^* are supported by \mathbb{R}_+ ,

ii) $\bar{\nu}_u([x, \infty)) \leq \bar{\nu}_u^*([x, \infty)) < \infty$, $x, u \in \mathbb{R}_+$, and

iii) $|H_u| \leq |H_u^*|$, $dPdu - a.e.$

Then we have:

$$\mathbb{E}[\phi(M_t + M_t^*)] \leq \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t, \quad (3.3)$$

for all convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$.

Next is a different version of the same result, under L^2 hypotheses.

Theorem 3.3. *Let*

$$\tilde{\nu}_u(dx) = |x|^2 \nu_u(dx) + |H_u|^2 \delta_0(dx), \quad \tilde{\nu}_u^*(dx) = |x|^2 \nu_u^*(dx) + |H_u^*|^2 \delta_0(dx),$$

$u \in \mathbb{R}_+$, and assume that:

i) ν_u and ν_u^* are supported by \mathbb{R} , and

ii) $\tilde{\nu}_u([x, \infty)) \leq \tilde{\nu}_u^*([x, \infty)) < \infty$, $x \in \mathbb{R}$, $u \in \mathbb{R}_+$.

Then we have:

$$\mathbb{E}[\phi(M_t + M_t^*)] \leq \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t, \quad (3.4)$$

for all convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ such that ϕ'' is non-decreasing.

In the proof we may assume that ϕ is \mathcal{C}^2 since a convex ϕ can be approximated by an increasing sequence of \mathcal{C}^2 convex Lipschitz functions, and the results can then be extended to the general case by an application of the monotone convergence theorem. In order to prove Theorem 3.2 and Theorem 3.3, we apply Itô's formula for forward/backward martingales (Theorem 8.1 in the Appendix Section 8), to $f(x_1, x_2) = \phi(x_1 + x_2)$:

$$\phi(M_t + M_t^*) = \phi(M_s + M_s^*)$$

$$\begin{aligned}
& + \int_{s^+}^t \phi'(M_{u^-} + M_u^*) dM_u + \frac{1}{2} \int_s^t \phi''(M_u + M_u^*) d\langle M^c, M^c \rangle_u \\
& + \sum_{s < u \leq t} (\phi(M_{u^-} + M_u^* + \Delta M_u) - \phi(M_{u^-} + M_u^*) - \Delta M_u \phi'(M_{u^-} + M_u^*)) \\
& - \int_s^{t^-} \phi'(M_u + M_{u^+}^*) d^* M_u - \frac{1}{2} \int_s^t \phi''(M_u + M_u^*) d\langle M^{*c}, M^{*c} \rangle_u \\
& - \sum_{s \leq u < t} (\phi(M_u + M_{u^+}^* + \Delta^* M_u^*) - \phi(M_u + M_{u^+}^*) - \Delta^* M_u^* \phi'(M_u + M_{u^+}^*)),
\end{aligned}$$

$0 \leq s \leq t$, where d and d^* denote the forward and backward Itô differential, respectively defined as the limits of the Riemann sums

$$\sum_{i=1}^n (M_{t_i^n} - M_{t_{i-1}^n}) \phi'(M_{t_{i-1}^n} + M_{t_{i-1}^n}^*)$$

and

$$\sum_{i=0}^{n-1} (M_{t_i^n}^* - M_{t_{i+1}^n}^*) \phi'(M_{t_{i+1}^n} + M_{t_{i+1}^n}^*)$$

for all refining sequences $\{s = t_0^n \leq t_1^n \leq \dots \leq t_{k_n}^n = t\}$, $n \geq 1$, of partitions of $[s, t]$ tending to the identity.

Proof of Theorem 3.2. Taking expectations on both sides of the above Itô formula we get

$$\begin{aligned}
\mathbb{E}[\phi(M_t + M_t^*)] &= \mathbb{E}[\phi(M_s + M_s^*)] + \frac{1}{2} \mathbb{E} \left[\int_s^t \phi''(M_u + M_u^*) d(\langle M^c, M^c \rangle_u - \langle M^{*c}, M^{*c} \rangle_u) \right] \\
&+ \mathbb{E} \left[\int_s^t \int_{-\infty}^{+\infty} (\phi(M_u + M_u^* + x) - \phi(M_u + M_u^*) - x \phi'(M_u + M_u^*)) \nu_u(dx) du \right] \\
&- \mathbb{E} \left[\int_s^t \int_{-\infty}^{+\infty} (\phi(M_u + M_u^* + x) - \phi(M_u + M_u^*) - x \phi'(M_u + M_u^*)) \nu_u^*(dx) du \right] \\
&= \mathbb{E}[\phi(M_s + M_s^*)] + \frac{1}{2} \mathbb{E} \left[\int_s^t \phi''(M_u + M_u^*) (|H_u|^2 - |H_u^*|^2) du \right] \\
&+ \mathbb{E} \left[\int_s^t \int_{-\infty}^{+\infty} \varphi(x, M_u + M_u^*) (\bar{\nu}_u(dx) - \bar{\nu}_u^*(dx)) du \right],
\end{aligned}$$

where

$$\varphi(x, y) = \frac{\phi(x + y) - \phi(y) - x \phi'(y)}{x}, \quad x, y \in \mathbb{R}.$$

The conclusion follows from the hypotheses and the fact that since ϕ is convex, the function $x \mapsto \varphi(x, y)$ is increasing in $x \in \mathbb{R}$ for all $y \in \mathbb{R}$. \square

Proof of Theorem 3.3. Using the following version of Taylor's formula

$$\phi(y+x) = \phi(y) + x\phi'(y) + |x|^2 \int_0^1 (1-\tau)\phi''(y+\tau x)d\tau, \quad x, y \in \mathbb{R},$$

which is valid for all \mathcal{C}^2 function ϕ , we get

$$\begin{aligned} \mathbb{E}[\phi(M_t + M_t^*)] &= \mathbb{E}[\phi(M_s + M_s^*)] \\ &+ \frac{1}{2}\mathbb{E} \left[\int_s^t \phi''(M_u + M_u^*)(|H_u|^2 - |H_u^*|^2)du \right] \\ &+ \mathbb{E} \left[\int_s^t \int_{-\infty}^{+\infty} |x|^2 \int_0^1 (1-\tau)\phi''(M_u + M_u^* + \tau x)d\tau \nu_u(dx)du \right] \\ &- \mathbb{E} \left[\int_s^t \int_{-\infty}^{+\infty} |x|^2 \int_0^1 (1-\tau)\phi''(M_u + M_u^* + \tau x)d\tau \nu_u^*(dx)du \right] \\ &= \mathbb{E}[\phi(M_s + M_s^*)] \\ &+ \mathbb{E} \left[\int_0^1 (1-\tau) \int_s^t \int_{-\infty}^{+\infty} \phi''(M_u + M_u^* + \tau x)(\tilde{\nu}_u(dx) - \tilde{\nu}_u^*(dx))dud\tau \right], \end{aligned}$$

and the conclusion follows from the hypothesis and the fact that ϕ'' is non-decreasing. \square

Note that if ϕ'' is also convex, then it suffices to assume that $\tilde{\nu}_u$ is more convex concentrated than $\tilde{\nu}_u^*$ instead of hypothesis (ii) in Theorem 3.3.

Remark 3.4. In case $|H_t| = |H_t^*|$ and $\nu_t = \nu_t^*$, $dPdt$ -a.e., from the proof of Theorem 3.2 and Theorem 3.3 we get the identity

$$\mathbb{E}[\phi(M_t + M_t^*)] = \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t, \quad (3.5)$$

for all sufficiently integrable function $\phi: \mathbb{R} \rightarrow \mathbb{R}$.

In particular, Relation (3.5) extends its natural counterpart in the independent increment case: given $(Z_s)_{s \in [0,t]}$, $(\tilde{Z}_s)_{s \in [0,t]}$ two independent copies of a Lévy process without drift, define the backward martingale $(Z_s^*)_{s \in [0,t]}$ as $Z_s^* = \tilde{Z}_{t-s}$, $s \in [0, t]$, then by convolution $\mathbb{E}[\phi(Z_s + Z_s^*)] = \mathbb{E}[\phi(Z_t)]$ does clearly not depend on $s \in [0, t]$.

Remark 3.5. If ϕ is non-decreasing, the proofs and statements of Theorem 3.2, Theorem 3.3, Corollary 3.8 and Corollary 3.7 extend to semi-martingales $(\hat{M}_t)_{t \in \mathbb{R}_+}$, $(\hat{M}_t^*)_{t \in \mathbb{R}_+}$ represented as

$$\hat{M}_t = M_t + \int_0^t \alpha_s ds \quad \text{and} \quad \hat{M}_t^* = M_t^* + \int_t^{+\infty} \beta_s ds, \quad (3.6)$$

provided $(\alpha_t)_{t \in \mathbb{R}_+}$, $(\beta_t)_{t \in \mathbb{R}_+}$, are respectively \mathcal{F}_t and \mathcal{F}_t^* -adapted with $\alpha_t \leq \beta_t$, $dPdt$ -a.e.

Let now $(\mathcal{F}_t^M)_{t \in \mathbb{R}_+}$, resp. $(\mathcal{F}_t^{M^*})_{t \in \mathbb{R}_+}$, denote the forward, resp. backward, filtration generated by $(M_t)_{t \in \mathbb{R}_+}$, resp. $(M_t^*)_{t \in \mathbb{R}_+}$.

Corollary 3.6. *Under the hypothesis of Theorem 3.2, if further $\mathbb{E}[M_t^* | \mathcal{F}_t^M] = 0$, $t \in \mathbb{R}_+$, then*

$$\mathbb{E}[\phi(M_t)] \leq \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t. \quad (3.7)$$

Proof. From (3.16) we get

$$\begin{aligned} \mathbb{E}[\phi(M_s + M_s^*)] &\geq \mathbb{E}[\phi(M_t + M_t^*)] \\ &= \mathbb{E}[\mathbb{E}[\phi(M_t + M_t^*) | \mathcal{F}_t^M]] \\ &\geq \mathbb{E}[\phi(M_t + \mathbb{E}[M_t^* | \mathcal{F}_t^M])] \\ &= \mathbb{E}[\phi(M_t)], \end{aligned}$$

$0 \leq s \leq t$, where we used Jensen's inequality. \square

In particular, if $M_0 = E[M_t]$ is deterministic (or \mathcal{F}_0^M is the trivial σ -field), Corollary 3.6 shows that $M_t - E[M_t]$ is more concentrated than M_0^* :

$$\mathbb{E}[\phi(M_t - E[M_t])] \leq \mathbb{E}[\phi(M_0^*)], \quad t \geq 0.$$

The filtrations $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ and $(\mathcal{F}_t^*)_{t \in \mathbb{R}_+}$ considered in Theorem 3.2 can be taken as $\mathcal{F}_t = \mathcal{F}_0^{M^*} \vee \mathcal{F}_t^M$, $\mathcal{F}_t^* = \mathcal{F}_\infty^M \vee \mathcal{F}_t^{M^*}$, $t \in \mathbb{R}_+$, provided $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ are independent. In this case, if additionally we have $M_T^* = 0$, then $\mathbb{E}[M_t^* | \mathcal{F}_t^M] = \mathbb{E}[M_t^*] = \mathbb{E}[M_T^*] = 0$, $0 \leq t \leq T$, hence the hypothesis of Corollary 3.6 is also satisfied. However the independence of $\langle M, M \rangle_t$ with $\langle M^*, M^* \rangle_t$, $t \in \mathbb{R}_+$, is not compatible (except in particular situations) with the assumptions imposed in Theorem 3.2.

In applications to convex concentration inequalities between random variables (admitting a predictable representation) and Poisson or Gaussian random variables, the independence of $(M_t)_{t \in \mathbb{R}_+}$ with $(M_t^*)_{t \in \mathbb{R}_+}$ will not be required, see Sections 4 and 5.

The case of bounded jumps

Assume now that $\nu^*(dt, dx)$ has the form

$$\nu^*(dt, dx) = \lambda_t^* \delta_k dt, \quad (3.8)$$

where $k \in \mathbb{R}_+$ and $(\lambda_t^*)_{t \in \mathbb{R}_+}$ is a positive \mathcal{F}_t^* -predictable process. Let

$$\lambda_{1,t} = \int_{-\infty}^{+\infty} x \nu_t(dx), \quad \lambda_{2,t}^2 = \int_{-\infty}^{+\infty} |x|^2 \nu_t(dx), \quad t \in \mathbb{R}_+,$$

denote respectively the compensator and quadratic variation of the jump part of $(M_t)_{t \in \mathbb{R}_+}$, under the respective assumptions

$$\int_{-\infty}^{+\infty} |x| \nu_t(dx) < \infty, \quad \text{and} \quad \int_{-\infty}^{+\infty} |x|^2 \nu_t(dx) < \infty, \quad (3.9)$$

$t \in \mathbb{R}_+$, P -a.s.

Corollary 3.7. *Assume that $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ have jump characteristics satisfying (3.8) and (3.9), that $(M_t)_{t \in \mathbb{R}_+}$ is \mathcal{F}_t^* -adapted, and that $(M_t^*)_{t \in \mathbb{R}_+}$ is \mathcal{F}_t -adapted. Then we have:*

$$\mathbb{E}[\phi(M_t + M_t^*)] \leq \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t, \quad (3.10)$$

for all \mathcal{C}^2 convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$, provided any of the three following conditions is satisfied:

i) $0 \leq \Delta M_t \leq k$, $dPdt$ - a.e., and

$$|H_t| \leq |H_t^*|, \quad \lambda_{1,t} \leq k \lambda_t^*, \quad dPdt - a.e.,$$

ii) $\Delta M_t \leq k$, $dPdt$ - a.e., and

$$|H_t| \leq |H_t^*|, \quad \lambda_{2,t}^2 \leq k^2 \lambda_t^*, \quad dPdt - a.e.,$$

iii) $\Delta M_t \leq 0$, $dPdt$ - a.e., and

$$|H_t|^2 + \lambda_{2,t}^2 \leq |H_t^*|^2 + k^2 \lambda_t^*, \quad dPdt - a.e.,$$

with moreover ϕ'' non-decreasing in cases *ii*) and *iii*).

Proof. The conditions $0 \leq \Delta M_t \leq k$, $\Delta M_t \leq k$, $\Delta M_t \leq 0$, are respectively equivalent to $\nu_t([0, k]^c) = 0$, $\nu_t((k, \infty)) = 0$, $\nu_t((0, \infty)) = 0$, hence under condition (*i*), the result follows from Theorem 3.2-*i*), and under conditions (*ii*) – (*iii*) it is an application of Theorem 3.2-*ii*). \square

For example we may take $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ of the form

$$M_t = M_0 + \int_0^t H_s dW_s + \int_0^t \int_{-\infty}^{+\infty} x(\mu(ds, dx) - \nu_s(dx)ds), \quad t \geq 0, \quad (3.11)$$

where $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion, and

$$M_t^* = \int_t^{+\infty} H_s^* d^*W_s^* + k \left(Z_t^* - \int_t^{+\infty} \lambda_s^* ds \right), \quad (3.12)$$

where $(W_t^*)_{t \in \mathbb{R}_+}$ is a backward Brownian motion and $(Z_t^*)_{t \in \mathbb{R}_+}$ is a backward point process with intensity $(\lambda_t^*)_{t \in \mathbb{R}_+}$. However in Section 5 we will consider an example for which the decomposition (3.12) does not hold.

The case of point processes

In particular, $(M_t)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ can be taken as

$$M_t = M_0 + \int_0^t H_s dW_s + \int_0^t J_s(dZ_s - \lambda_s ds), \quad t \in \mathbb{R}_+, \quad (3.13)$$

and

$$M_t^* = \int_t^{+\infty} H_s^* d^*W_s^* + \int_t^{+\infty} J_s^*(d^*Z_s^* - \lambda_s^* ds), \quad t \in \mathbb{R}_+, \quad (3.14)$$

where $(W_t)_{t \in \mathbb{R}_+}$ is a standard Brownian motion, $(Z_t)_{t \in \mathbb{R}_+}$ is a point process with intensity $(\lambda_t)_{t \in \mathbb{R}_+}$, $(W_t^*)_{t \in \mathbb{R}_+}$ is a backward standard Brownian motion, and $(Z_t^*)_{t \in \mathbb{R}_+}$ is a backward point process with intensity $(\lambda_t^*)_{t \in \mathbb{R}_+}$, and $(H_t)_{t \in \mathbb{R}_+}$, $(J_t)_{t \in \mathbb{R}_+}$, resp. $(H_t^*)_{t \in \mathbb{R}_+}$, $(J_t^*)_{t \in \mathbb{R}_+}$ are predictable with respect to $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$, resp. $(\mathcal{F}_t^*)_{t \in \mathbb{R}_+}$.

In this case we have

$$\nu(dt, dx) = \lambda_t \delta_{J_t}(dx) dt \quad \text{and} \quad \nu^*(dt, dx) = \lambda_t^* \delta_{J_t^*}(dx) dt. \quad (3.15)$$

Corollary 3.8. *Let $(M_t)_{t \in \mathbb{R}_+}$, $(M_t^*)_{t \in \mathbb{R}_+}$ have the jump characteristics (3.15) and assume that $(M_t)_{t \in \mathbb{R}_+}$ is \mathcal{F}_t^* -adapted and $(M_t^*)_{t \in \mathbb{R}_+}$ is \mathcal{F}_t -adapted. Then we have:*

$$\mathbb{E}[\phi(M_t + M_t^*)] \leq \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t, \quad (3.16)$$

for all \mathcal{C}^2 convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$, provided any of the three following conditions are satisfied:

i) $0 \leq J_t \leq J_t^*$, $\lambda_t dPdt - a.e.$ and

$$|H_t| \leq |H_t^*|, \quad \lambda_t J_t \leq \lambda_t^* J_t^*, \quad dPdt - a.e.,$$

ii) $J_t \leq J_t^*$, $\lambda_t dPdt - a.e.$, and

$$|H_t| \leq |H_t^*|, \quad \lambda_t |J_t|^2 \leq \lambda_t^* |J_t^*|^2, \quad dPdt - a.e..$$

iii) $J_t \leq 0 \leq J_t^*$, $\lambda_t dPdt - a.e.$, and

$$|H_t|^2 + \lambda_t |J_t|^2 \leq |H_t^*|^2 + \lambda_t^* |J_t^*|^2, \quad dPdt - a.e..$$

with moreover ϕ'' non-decreasing in cases ii) and iii).

Proof. Under condition (i), the result follows from Theorem 3.3-i), and under conditions (ii) – (iii) it is an application of Theorem 3.3-ii). \square

Note that condition i) in Corollary 3.8 can be replaced with:

i') $0 \leq J_t \leq J_t^*$, $\lambda_t dPdt - a.e.$ and

$$|H_t| \leq |H_t^*|, \quad \lambda_t \leq \lambda_t^*, \quad dPdt - a.e.$$

4 Application to point processes

Let $(W_t)_{t \in \mathbb{R}_+}$ and $(Z_t)_{t \in \mathbb{R}_+}$ be a standard Brownian motion and a point process with intensity $(\lambda_t)_{t \in \mathbb{R}_+}$, which generate a filtration $(\mathcal{F}_t^M)_{t \in \mathbb{R}_+}$. Consider F a random variable with the representation

$$F = \mathbb{E}[F] + \int_0^{+\infty} H_t dW_t + \int_0^{+\infty} J_t (dZ_t - \lambda_t dt), \quad (4.1)$$

where $(H_u)_{u \in \mathbb{R}_+}$ is a square-integrable predictable process and $(J_t)_{t \in \mathbb{R}_+}$ is a predictable process which is either square-integrable or positive and integrable. Let $\tilde{N}(c)$ denote a compensated Poisson random variable with intensity $c > 0$, and let $W(\beta^2)$ denote a centered Gaussian random variable with variance $\beta^2 \geq 0$. Theorem 4.1 is a consequence of Corollary 3.8 below. Note that the inequality (4.2) below is weaker than (4.3) but it holds for a wider class of functions, i.e. for all convex functions instead of convex functions with non-decreasing second derivative.

Theorem 4.1. *Let F have the representation (4.1):*

$$F = \mathbb{E}[F] + \int_0^{+\infty} H_t dW_t + \int_0^{+\infty} J_t (dZ_t - \lambda_t dt).$$

i) *Assume that $0 \leq J_t \leq k$, $dPdt$ -a.e., for some $k > 0$, and let*

$$\beta_1^2 = \left\| \int_0^{+\infty} |H_t|^2 dt \right\|_{\infty} \quad \text{and} \quad \alpha_1 = \left\| \int_0^{+\infty} J_t \lambda_t dt \right\|_{\infty}.$$

Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(W(\beta_1^2) + k\tilde{N}(\alpha_1/k) \right) \right], \quad (4.2)$$

for all convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$.

ii) *Assume that $J_t \leq k$, $dPdt$ -a.e., for some $k > 0$, and let*

$$\beta_2^2 = \left\| \int_0^{+\infty} |H_t|^2 dt \right\|_{\infty} \quad \text{and} \quad \alpha_2^2 = \left\| \int_0^{+\infty} |J_t|^2 \lambda_t dt \right\|_{\infty}.$$

Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(W(\beta_2^2) + k\tilde{N}(\alpha_2^2/k^2) \right) \right], \quad (4.3)$$

for all \mathcal{C}^2 convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ such that ϕ'' is non-decreasing.

iii) *Assume that $J_t \leq 0$, $dPdt$ -a.e., and let*

$$\beta_3^2 = \left\| \int_0^{+\infty} |H_t|^2 dt + \int_0^{+\infty} |J_t|^2 \lambda_t dt \right\|_{\infty}.$$

Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi(W(\beta_3^2)) \right], \quad (4.4)$$

for all \mathcal{C}^2 convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ such that ϕ'' is non-decreasing.

Proof. Consider the martingale

$$M_t = \mathbb{E}[F|\mathcal{F}_t] - \mathbb{E}[F] = \int_0^t H_s dW_s + \int_0^t J_s (dZ_s - \lambda_s ds), \quad t \geq 0,$$

generating the filtration $(\mathcal{F}_t^M)_{t \in \mathbb{R}_+}$, and let $(\hat{N}_s)_{s \in \mathbb{R}_+}$, $(\hat{W}_s)_{s \in \mathbb{R}_+}$ denote a left-continuous standard Poisson process and a standard Brownian motion independent of $(\mathcal{F}_s^M)_{s \in \mathbb{R}_+}$.

i) – ii) For $p = 1, 2$, let the filtrations $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ and $(\mathcal{F}_t^*)_{t \in \mathbb{R}_+}$ be defined by

$$\mathcal{F}_t^* = \mathcal{F}_\infty^M \vee \sigma(\hat{W}_{\beta_p^2} - \hat{W}_{V_p^2(s)}, \hat{N}_{\alpha_p^p/k^p} - \hat{N}_{U_p^p(s)/k^p} : s \geq t),$$

and $\mathcal{F}_t = \sigma(\hat{W}_s, \hat{N}_s : s \geq 0) \vee \mathcal{F}_t^M$, $t \in \mathbb{R}_+$, and let

$$M_t^* = \hat{W}_{\beta_p^2} - \hat{W}_{V_p^2(t)} + k(\hat{N}_{\alpha_p^p/k^p} - \hat{N}_{U_p^p(t)/k^p}) - (\alpha_p^p - U_p^p(t))/k^{p-1}, \quad (4.5)$$

where

$$V_p^2(t) = \int_0^t |H_s|^2 ds \quad \text{and} \quad U_p^p(t) = \int_0^t J_s^p \lambda_s ds, \quad P - a.s., \quad s \geq 0.$$

Then $(M_t^*)_{t \in \mathbb{R}_+}$ satisfies the hypothesis of Corollary 3.8–*i) – ii)*, as well as the condition $E[M_t^* | \mathcal{F}_t^M] = 0$, $t \in \mathbb{R}_+$, with $H_s^* = H_s$, $J_s^* = k$, $\lambda_s^* = J_s^p \lambda_s / k^p$, $dPds$ -a.e., hence

$$\mathbb{E}[\phi(M_t)] \leq \mathbb{E}[\phi(M_0^*)],$$

and letting t go to infinity we obtain (4.2) and (4.3), respectively for $p = 1$ and $p = 2$.

iii) Let

$$M_s^* = W_{\beta_3^2} - W_{U_3^2(s)}, \quad (4.6)$$

where

$$U_3^2(s) = \int_0^s |H_u|^2 du + \int_0^s |J_u|^2 \lambda_u du, \quad P - a.s.$$

Then $(M_t^*)_{t \in \mathbb{R}_+}$ satisfies the hypothesis of Corollary 3.8–*iii)* with $|H_s^*|^2 = |H_s|^2 + |J_s|^2 \lambda_s$ and $\lambda_s^* = J_s^* = 0$, $dPds$ -a.e., hence

$$\mathbb{E}[\phi(M_t)] \leq \mathbb{E}[\phi(M_0^*)],$$

and letting t go to infinity we obtain (4.4). □

Remark 4.2. *The proof of Theorem 4.1 can also be obtained from Corollary 3.7.*

Proof. Let

$$\mu(dt, dx) = \sum_{\Delta Z_s \neq 0} \delta_{(s, J_s)}(dt, dx), \quad \nu_t(dx) = \lambda_t \delta_{J_t}(dx).$$

i) – ii) In both cases $p = 1, 2$, let $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$, $(\mathcal{F}_t^*)_{t \in \mathbb{R}_+}$ and $(M_t^*)_{t \in \mathbb{R}_+}$ be defined in (4.5), with $V_p^2(t) = \int_0^t |H_s|^2 ds$ and $U_p^p(t) = \int_0^t |J_s|^p ds$, P -a.s., $t \geq 0$. Then $(M_t^*)_{t \in \mathbb{R}_+}$ satisfies the hypothesis of Corollary 3.7-*i) – ii)*, with $H_s^* = H_s$, $\nu_s^* = |J_s|^p/k^p$, $dPds$ -a.e.

iii) Let $(M_s^*)_{s \in \mathbb{R}_+}$ be defined as in (4.6), and let $U_3^2(s) = \int_0^s |H_u|^2 du + \int_0^s |J_u|^2 du$, $|H_s^*|^2 = |H_s|^2 + |J_s|^2$ and $\nu_s^* = 0$, $dPds$ -a.e. \square

In the pure jump case, Theorem 4.1-*ii)* yields

$$P(M_T \geq x) \leq \exp \left(\frac{y}{k} - \left(\frac{y}{k} + \frac{\alpha_2^2}{k^2} \right) \log \left(1 + \frac{ky}{\alpha_2^2} \right) \right) \leq \exp \left(-\frac{y}{2k} \log \left(1 + \frac{ky}{\alpha_2^2} \right) \right),$$

$y > 0$, with $\alpha_2^2 = \|\langle M, M \rangle_T\|_\infty$, cf. Theorem 23.17 of [7], although some differences in the hypotheses make the results not directly comparable: here no lower bound is assumed on jump sizes, and the presence of a continuous component is treated in a different way.

The results of this section and the next one apply directly to solutions of stochastic differential equations such as

$$dX_t = a(t, X_t)dW_t + b(t, X_t)(dZ_t - \lambda_t dt),$$

with $H_t = a(t, X_t)$, $J_t = b(t, X_t)$, $t \in \mathbb{R}_+$, for which the hypotheses can be formulated directly on the coefficients $a(\cdot, \cdot)$, $b(\cdot, \cdot)$ without explicit knowledge of the solution.

5 Application to Poisson random measures

Since a large family of point processes can be represented as stochastic integrals with respect to Poisson random measures (see e.g. [5], Section 4, Ch. XIV), it is natural to investigate the consequences of Theorem 3.2 in the setting of Poisson random

measures. Let σ be a Radon measure on \mathbb{R}^d , diffuse on $\mathbb{R}^d \setminus \{0\}$, such that $\sigma(\{0\}) = 1$, and

$$\int_{\mathbb{R}^d \setminus \{0\}} (|x|^2 \wedge 1) \sigma(dx) < \infty,$$

and consider a random measure $\omega(dt, dx)$ of the form

$$\omega(dt, dx) = \sum_{i \in \mathbb{N}} \delta_{(t_i, x_i)}(dt, dx)$$

identified to its (locally finite) support $\{(t_i, x_i)\}_{i \in \mathbb{N}}$. We assume that $\omega(dt, dx)$ is Poisson distributed with intensity $dt\sigma(dx)$ on $\mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}$, and consider a standard Brownian motion $(W_t)_{t \in \mathbb{R}_+}$, independent of $\omega(dt, dx)$, under a probability P on Ω . Let

$$\mathcal{F}_t = \sigma(W_s, \omega([0, s] \times A)) : 0 \leq s \leq t, A \in \mathcal{B}_b(\mathbb{R}^d \setminus \{0\}), \quad t \in \mathbb{R}_+,$$

where $\mathcal{B}_b(\mathbb{R}^d \setminus \{0\}) = \{A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}) : \sigma(A) < \infty\}$. The stochastic integral of a square-integrable \mathcal{F}_t -predictable process $u \in L^2(\Omega \times \mathbb{R}_+ \times \mathbb{R}^d, dP \times dt \times d\sigma)$ is written as

$$\int_0^{+\infty} u(t, 0) dW_t + \int_{\mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}} u(t, x) (\omega(dt, dx) - \sigma(dx)dt), \quad (5.1)$$

and satisfies the Itô isometry

$$\begin{aligned} E \left[\left(\int_0^{+\infty} u(t, 0) dW_t + \int_0^{+\infty} \int_{\mathbb{R}^d \setminus \{0\}} u(t, x) (\omega(dt, dx) - \sigma(dx)dt) \right)^2 \right] \\ = E \left[\int_0^{+\infty} u^2(t, 0) dt \right] + E \left[\int_{\mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}} u^2(t, x) \sigma(dx) dt \right] \end{aligned} \quad (5.2)$$

$$= E \left[\int_{\mathbb{R}_+ \times \mathbb{R}^d} u^2(t, x) \sigma(dx) dt \right]. \quad (5.3)$$

Recall that due to the Itô isometry, the predictable and adapted version of u can be used indifferently in the stochastic integral (5.1), cf. p. 199 of [3] for details. When $u \in L^2(\mathbb{R}_+ \times \mathbb{R}^d, dt \times d\sigma)$, the characteristic function of

$$I_1(u) := \int_0^{+\infty} u(t, 0) dW_t + \int_{\mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}} u(t, x) (\omega(dt, dx) - \sigma(dx)dt),$$

is given by the Lévy-Khintchine formula

$$E [e^{iI_1(u)}] = \exp \left(-\frac{1}{2} \int_0^{+\infty} u^2(t, 0) dt + \int_{\mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}} (e^{iu(t, x)} - 1 - iu(t, x)) \sigma(dx) dt \right).$$

Theorem 5.1. *Let F with the representation*

$$F = E[F] + \int_0^{+\infty} H_s dW_s + \int_0^{+\infty} \int_{\mathbb{R}^d \setminus \{0\}} J_{u,x} (\omega(du, dx) - \sigma(dx)du),$$

where $(H_t)_{t \in \mathbb{R}_+} \in L^2(\Omega \times \mathbb{R}_+)$, and $(J_{t,x})_{(t,x) \in \mathbb{R}_+ \times \mathbb{R}^d}$ are \mathcal{F}_t -predictable with $(J_{t,x})_{(t,x) \in \mathbb{R}_+ \times \mathbb{R}^d} \in L^1(\Omega \times \mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}, dP \times dt \times d\sigma)$ and $(J_{t,x})_{(t,x) \in \mathbb{R}_+ \times \mathbb{R}^d} \in L^2(\Omega \times \mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}, dP \times dt \times d\sigma)$ respectively in (i) and in (ii – iii) below.

i) Assume that $0 \leq J_{u,x} \leq k$, $dP\sigma(dx)du$ -a.e., for some $k > 0$, and let

$$\beta_1^2 = \left\| \int_0^{+\infty} |H_u|^2 du \right\|_{\infty}, \quad \text{and} \quad \alpha_1(x) = \left\| \int_0^{+\infty} J_{u,x} du \right\|_{\infty}, \quad \sigma(dx) - \text{a.e.}$$

Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(W(\beta_1^2) + k\tilde{N} \left(\int_{\mathbb{R}^d \setminus \{0\}} \frac{\alpha_1(x)}{k} \sigma(dx) \right) \right) \right],$$

for all convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$.

ii) Assume that $J_{u,x} \leq k$, $dP\sigma(dx)du$ -a.e., for some $k > 0$, and let

$$\beta_2^2 = \left\| \int_0^{+\infty} |H_u|^2 du \right\|_{\infty}, \quad \text{and} \quad \alpha_2^2(x) = \left\| \int_0^{+\infty} |J_{u,x}|^2 du \right\|_{\infty}, \quad \sigma(dx) - \text{a.e.}$$

Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(W(\beta_2^2) + k\tilde{N} \left(\int_{\mathbb{R}^d \setminus \{0\}} \frac{\alpha_2^2(x)}{k^2} \sigma(dx) \right) \right) \right],$$

for all \mathcal{C}^2 convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ such that ϕ'' is non-decreasing.

iii) Assume that $J_{u,x} \leq 0$, $dP\sigma(dx)du$ -a.e., and let

$$\beta_3^2 = \left\| \int_0^{+\infty} |H_u|^2 du + \int_0^{+\infty} \int_{\mathbb{R}^d \setminus \{0\}} |J_{u,x}|^2 du \sigma(dx) \right\|_{\infty}.$$

Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} [\phi(W(\beta_3^2))],$$

for all \mathcal{C}^2 convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ such that ϕ'' is non-decreasing.

Proof. The proof is similar to that of Theorem 4.1, replacing the use of Corollary 3.8 by that of Corollary 3.7. Let

$$M_t = M_0 + \int_0^t H_u dW_u + \int_{\mathbb{R}^d \setminus \{0\}} \int_0^t J_{u,x} (\omega(du, dx) - \sigma(dx)du),$$

generating the filtration $(\mathcal{F}_t^M)_{t \in \mathbb{R}_+}$. Here, $\nu_t(dx)$ denotes the image measure of $\sigma(dx)$ by the mapping $x \mapsto J_{t,x}$, $t \geq 0$, and $\mu(dt, dx)$ denotes the image measure of $\omega(dt, dx)$ by $(s, y) \mapsto (s, J_{s,y})$, i.e.

$$\mu(dt, dx) = \sum_{\omega(\{(s,y)\})=1} \delta_{(s, J_{s,y})}(dt, dx).$$

i) – ii) For $p = 1, 2$, let the filtrations $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ and $(\mathcal{F}_t^*)_{t \in \mathbb{R}_+}$ be defined by

$$\mathcal{F}_t^* = \mathcal{F}_\infty^M \vee \sigma(\hat{W}_{\beta_p^2} - \hat{W}_{V_p^2(s)} + \hat{N}_{\alpha_p^p/k^p} - \hat{N}_{U_p^p(s)/k^p} : s \geq t),$$

and

$$\mathcal{F}_t = \mathcal{F}_t^M \vee \sigma(\hat{W}_s, \hat{N}_s : s \geq 0), \quad t \in \mathbb{R}_+,$$

and let

$$M_t^* = \hat{W}_{\beta_p^2} - \hat{W}_{V_p^2(t)} + k(\hat{N}_{\alpha_p^p/k^p} - \hat{N}_{U_p^p(t)/k^p}) - (\alpha_p^p - U_p^p(t))/k^{p-1},$$

where

$$V_p^2(t) = \int_0^t |H_s|^2 ds \quad \text{and} \quad U_p^p(t) = \int_0^t \int_{-\infty}^{+\infty} x^p \nu_s(dx) ds, \quad P - a.s., \quad t \geq 0.$$

Then $(M_t^*)_{t \in \mathbb{R}_+}$ satisfies the hypothesis of Theorem 3.2–*i) – ii)*, and also the condition $E[M_t^* | \mathcal{F}_t^M] = 0$, $t \in \mathbb{R}_+$, with $H_s^* = H_s$, $\nu_s^* = \int_{-\infty}^{+\infty} x^p \nu_s(dx)$, $dP ds$ -a.e., hence

$$\mathbb{E}[\phi(M_t)] \leq \mathbb{E}[\phi(M_0^*)].$$

Letting t go to infinity we obtain (4.2) and (4.3), respectively for $p = 1, 2$.

iii) Let

$$M_s^* = W_{\beta_3^2} - W_{U_3^2(s)},$$

where

$$U_3^2(s) = \int_0^s |H_u|^2 du + \int_0^s \int_{-\infty}^{+\infty} |x|^2 \nu_u(dx) du, \quad P - a.s., \quad s \geq 0.$$

Then $(M_t^*)_{t \in \mathbb{R}_+}$ satisfies the hypotheses of Theorem 3.2–iii) with

$$|H_s^*|^2 = |H_s|^2 + \int_{-\infty}^{+\infty} |x|^2 \nu_s(dx)$$

and $\nu_s^* = 0$, $dPds$ -a.e., hence

$$\mathbb{E}[\phi(M_t)] \leq \mathbb{E}[\phi(M_0^*)],$$

and letting t go to infinity we obtain (4.4). \square

In Theorem 4.1, $(Z_t)_{t \in \mathbb{R}_+}$ can be taken equal to the standard Poisson process $(N_t)_{t \in \mathbb{R}_+}$, which also satisfies the hypotheses of Theorem 5.1 since it can be defined with $d = 1$ and $\sigma(dx) = 1_{[0,1]}(x)dx$ as

$$N_t = \omega([0, t] \times [0, 1]), \quad t \geq 0.$$

In other terms, being a point process, $(N_t)_{t \in \mathbb{R}_+}$ is at the intersection of Corollary 3.7 and Corollary 3.8, as already noted in Remark 4.2.

6 Clark formula

In this section we examine the consequence of results of Section 5 when the predictable representation of random variables is obtained via the Clark formula. The multiple stochastic integral $I_n(h_n)$ of $h_n \in L^2(\mathbb{R}_+ \times \mathbb{R}^d)^{\otimes n}$ can be defined by induction with

$$I_n(h_n) = n \int_{\mathbb{R}_+ \times \mathbb{R}^d} I_{n-1}(\pi_{t,x}^n h_n)(\omega(dt, dx) - \sigma(dx)dt),$$

where

$$(\pi_{t,x}^n h_n)(t_1, x_1, \dots, t_{n-1}, x_{n-1}) := h_n(t_1, x_1, \dots, t_{n-1}, x_{n-1}, t, x) 1_{[0,t]}(t_1) \cdots 1_{[0,t]}(t_{n-1}),$$

$t_1, \dots, t_{n-1}, t \in \mathbb{R}_+$, $x_1, \dots, x_{n-1}, x \in \mathbb{R}^d$. The isometry property

$$E [I_n(h_n)^2] = n! \|h_n\|_{L^2(\mathbb{R}_+ \times \mathbb{R}^d)^{\otimes n}}^2$$

follows by induction from (5.2). Let the linear, closable, finite difference operator

$$D : L^2(\Omega, P) \longrightarrow L^2(\Omega \times \mathbb{R}_+ \times \mathbb{R}^d, dP \times dt \times d\sigma)$$

be defined as

$$D_{t,x}I_n(f_n) = nI_{n-1}(f_n(*, t, x)), \quad \sigma(dx)dt - a.e.,$$

cf. e.g. [10], [15], and

$$D_{t,0}I_n(f_n) = nI_{n-1}(f_n(*, t, x)), \quad \sigma(dx)dt - a.e.,$$

Recall that for $\sigma(dx)dt$ -a.e. every $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$ we have

$$D_{t,x}F(\omega) = F(\omega \cup \{(t, x)\}) - F(\omega), \quad P(d\omega) - a.e.,$$

cf. e.g. [10], [12], and that $D_{t,0}$ has the derivation property, i.e.

$$D_{t,0}f(I_1(f_1^{(1)}), \dots, I_1(f_1^{(n)})) = \sum_{k=1}^n f_1^{(i)}(t, 0) \partial_k f(I_1(f_1^{(1)}), \dots, I_1(f_1^{(n)})),$$

$P(d\omega) - a.e.$, cf. e.g. [14].

The Clark formula for Lévy processes, cf. [11], [14], states that every $F \in L^2(\Omega)$ has the representation

$$F = \mathbb{E}[F] + \int_0^{+\infty} \mathbb{E}[D_{s,0}F | \mathcal{F}_s] dW_s + \int_0^{+\infty} \int_{\mathbb{R}^d \setminus \{0\}} \mathbb{E}[D_{s,x}F | \mathcal{F}_s] (\omega(ds, dx) - \sigma(dx)ds). \quad (6.1)$$

(The formula originally holds for F in the domain of D but its extension to $L^2(\Omega)$ is straightforward, cf. [14], Proposition 12). Theorem 5.1 immediately yields the following corollary when applied to any $F \in L^2(\Omega)$ represented as in (6.1).

Corollary 6.1. *Let $F \in L^2(\Omega)$ have the representation (6.1), and assume additionally that $\int_0^{+\infty} \int_{\mathbb{R}^d \setminus \{0\}} |\mathbb{E}[D_{s,x}F | \mathcal{F}_s]| \sigma(dx)ds < \infty$ a.s. in (i) below.*

i) *Assume that $0 \leq \mathbb{E}[D_{u,x}F | \mathcal{F}_u] \leq k$, $dP\sigma(dx)du$ -a.e., for some $k > 0$, and let*

$$\beta_1^2 = \left\| \int_0^{+\infty} (\mathbb{E}[D_{u,0}F | \mathcal{F}_u])^2 du \right\|_{\infty}, \quad \text{and} \quad \alpha_1(x) = \left\| \int_0^{+\infty} \mathbb{E}[D_{u,x}F | \mathcal{F}_u] du \right\|_{\infty},$$

$\sigma(dx)$ -a.e. Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(W(\beta_1^2) + k\tilde{N} \left(\int_{\mathbb{R}^d \setminus \{0\}} \frac{\alpha_1(x)}{k} \sigma(dx) \right) \right) \right], \quad (6.2)$$

for all convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$.

ii) Assume that $\mathbb{E}[D_{u,x}F|\mathcal{F}_u] \leq k$, $dP\sigma(dx)du$ -a.e., for some $k > 0$, and let

$$\beta_2^2 = \left\| \int_0^{+\infty} (\mathbb{E}[D_{u,0}F|\mathcal{F}_u])^2 du \right\|_\infty, \quad \text{and} \quad \alpha_2^2(x) = \left\| \int_0^{+\infty} (\mathbb{E}[D_{u,x}F|\mathcal{F}_u])^2 du \right\|_\infty,$$

$\sigma(dx)$ -a.e. Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(W(\beta_2^2) + k\tilde{N} \left(\int_{\mathbb{R}^d \setminus \{0\}} \frac{\alpha_2^2(x)}{k^2} \sigma(dx) \right) \right) \right], \quad (6.3)$$

for all \mathcal{C}^2 convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ such that ϕ'' is non-decreasing.

iii) Assume that $\mathbb{E}[D_{u,x}F|\mathcal{F}_u] \leq 0$, $dP\sigma(dx)du$ -a.e., and let

$$\beta_3^2 = \left\| \int_0^{+\infty} (\mathbb{E}[D_{u,0}F|\mathcal{F}_u])^2 du + \int_0^{+\infty} \int_{\mathbb{R}^d \setminus \{0\}} (\mathbb{E}[D_{u,x}F|\mathcal{F}_u])^2 du \sigma(dx) \right\|_\infty.$$

Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} [\phi(W(\beta_3^2))], \quad (6.4)$$

for all \mathcal{C}^2 convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$ such that ϕ'' is non-decreasing.

As mentioned in the introduction, from (6.4) we deduce the deviation inequality

$$P(E - \mathbb{E}[F] \geq y) \leq \exp \left(-\frac{y^2}{2\beta_3^2} \right), \quad y > 0,$$

provided $\mathbb{E}[D_{u,x}F|\mathcal{F}_u] \leq 0$, $dP\sigma(dx)du$ -a.e., and

$$\int_0^{+\infty} (\mathbb{E}[D_{u,0}F|\mathcal{F}_u])^2 du + \int_0^{+\infty} \int_{\mathbb{R}^d \setminus \{0\}} (\mathbb{E}[D_{u,x}F|\mathcal{F}_u])^2 du \sigma(dx) \leq \beta_3^2, \quad P - a.s.$$

Similarly from (6.3) we get

$$P(E - \mathbb{E}[F] \geq y) \leq \exp \left(\frac{y}{k} - \left(\frac{y}{k} + \frac{\alpha_2^2}{k^2} \right) \log \left(1 + \frac{ky}{\alpha_2^2} \right) \right), \quad y > 0, \quad (6.5)$$

provided

$$E[D_{t,x}F|\mathcal{F}_t] \leq k, \quad dP\sigma(dx)dt - a.e., \quad (6.6)$$

and

$$\int_{\mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}} (E[D_{t,x}F|\mathcal{F}_t])^2 \sigma(dx)dt \leq \alpha_2^2, \quad P - a.s.,$$

for some $k > 0$ and $\alpha_2^2 > 0$. In [1] this latter estimate has been proved using (modified) logarithmic Sobolev inequalities and the Herbst method under the stronger condition

$$|D_{t,x}F| \leq k, \quad dP\sigma(dx)dt\text{-a.e.}, \quad (6.7)$$

and

$$\int_{\mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}} |D_{t,x}F|^2 \sigma(dx)dt \leq \alpha_2^2, \quad P - a.s., \quad (6.8)$$

for some $k > 0$ and $\alpha_2^2 > 0$. In [17] it has been shown, using sharp logarithmic Sobolev inequalities, that the condition $|D_{t,x}F| \leq k$ can be relaxed to

$$D_{t,x}F \leq k, \quad dP\sigma(dx)dt\text{-a.e.}, \quad (6.9)$$

which is nevertheless stronger than (6.6).

In the next result, which however imposes uniform almost sure bounds on DF , we consider Poisson random measures on $\mathbb{R}^d \setminus \{0\}$ instead of $\mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}$.

Corollary 6.2. *i) Assume that $0 \leq D_x F \leq \beta(x) \leq k$, $dP\sigma(dx) - a.e.$, where $\beta(\cdot) : \mathbb{R}^d \setminus \{0\} \rightarrow [0, k]$ is deterministic and $k > 0$. Then for all convex function ϕ we have*

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(k\tilde{N} \left(\int_{\mathbb{R}^d \setminus \{0\}} \frac{\beta(x)}{k} \sigma(dx) \right) \right) \right].$$

ii) Assume that $|D_x F| \leq \beta(x) \leq k$, $dP\sigma(dx)\text{-a.e.}$, where $\beta(\cdot) : \mathbb{R} \rightarrow [0, k]$ and $k > 0$ are deterministic. Then for all \mathcal{C}^2 convex function ϕ with non-decreasing second derivative we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(k\tilde{N} \left(\int_{\mathbb{R}^d \setminus \{0\}} \frac{\beta^2(x)}{k^2} \sigma(dx) \right) \right) \right].$$

iii) Assume that $-\beta(x) \leq D_x F \leq 0$, $dP\sigma(dx)$ -a.e., where $\beta(\cdot) : \mathbb{R} \rightarrow [0, \infty)$ is deterministic. Then for all \mathcal{C}^2 convex function ϕ with non-decreasing second derivative we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(W \left(\int_{\mathbb{R}^d \setminus \{0\}} \beta^2(x) \sigma(dx) \right) \right) \right].$$

Proof. Assume that $\omega(dt, dx)$ has intensity $1_{[0,1]}(s)\sigma(dx)ds$ on $\mathbb{R}_+ \times \mathbb{R}^d \setminus \{0\}$, we define the random measure $\hat{\omega}$ on $\mathbb{R}^d \setminus \{0\}$ with intensity $\sigma(dx)$ as

$$\hat{\omega}(A) = \omega([0, 1] \times A), \quad A \in \mathcal{B}_b(\mathbb{R}^d \setminus \{0\}),$$

and let $\hat{F}(\omega) = F(\hat{\omega})$. Then it remains to apply Corollary 6.1 with the relation $D_{s,x}\hat{F}(\omega) = D_x F(\hat{\omega})$, for $(s, x) \in [0, 1] \times \mathbb{R}^d \setminus \{0\}$. \square

In Corollary 6.2, $\mathbb{R}^d \setminus \{0\}$ can be replaced by \mathbb{R}^d without additional difficulty.

7 Normal martingales

In this section we interpret the above results in the framework of normal martingales. Let $(Z_t)_{t \in \mathbb{R}_+}$ be a normal martingale, i.e. $(Z_t)_{t \in \mathbb{R}_+}$ is a martingale such that $d\langle Z, Z \rangle_t = dt$. If $(Z_t)_{t \in \mathbb{R}_+}$ is in L^4 and has the chaotic representation property it satisfies the structure equation

$$d[Z, Z]_t = dt + \gamma_t dZ_t, \quad t \in \mathbb{R}^+,$$

where $(\gamma_t)_{t \in \mathbb{R}_+}$ is a predictable square-integrable process, cf. [4]. Recall that the cases $\gamma_s = 0$, $\gamma_s = c \in \mathbb{R} \setminus \{0\}$, $\gamma_s = \beta Z_s$, $\beta \in (-2, 0)$, correspond respectively to Brownian motion, the compensated Poisson process with jump size c and intensity $1/c^2$, and to the Azéma martingales. Consider the martingale

$$M_t = M_0 + \int_0^t R_u dZ_u, \tag{7.1}$$

where $(R_u)_{u \in \mathbb{R}_+} \in L^2(\Omega \times \mathbb{R}_+)$ is predictable. We have

$$d\langle M^c, M^c \rangle_t = 1_{\{\gamma_t=0\}} |R_t|^2 dt$$

and

$$\mu(dt, dx) = \sum_{\Delta Z_s \neq 0} \delta_{(s, R_s \gamma_s)}(dt, dx), \quad \nu(dt, dx) = \frac{1}{\gamma_t^2} \sum_{\Delta Z_s \neq 0} \delta_{R_s \gamma_s}(dx) dt,$$

and the Itô formula, cf. [4]:

$$\begin{aligned}\phi(M_t) &= \phi(M_s) + \int_s^t 1_{\{\gamma_u=0\}} R_u \phi'(M_u) dZ_u + \int_s^t 1_{\{\gamma_u \neq 0\}} \frac{\phi(M_{u-} + \gamma_u R_u) - \phi(M_{u-})}{\gamma_u} dZ_u \\ &\quad + \frac{1}{2} \int_s^t 1_{\{\gamma_u=0\}} |R_u|^2 \phi''(M_u) du + \int_s^t 1_{\{\gamma_u \neq 0\}} \frac{\phi(M_u + \gamma_u R_u) - \phi(M_u) - \gamma_u R_u \phi'(M_u)}{|\gamma_u|^2} du,\end{aligned}$$

$\phi \in \mathcal{C}^2(\mathbb{R})$. The multiple stochastic integrals with respect to $(M_t)_{t \in \mathbb{R}_+}$ are defined as

$$I_n(f_n) = n! \int_0^{+\infty} \int_0^{t_n} \cdots \int_0^{t_2} f_n(t_1, \dots, t_n) dM_{t_1} \cdots dM_{t_n},$$

for f_n a symmetric function in $L^2(\mathbb{R}_+^n)$. As an application of Corollary 3.8 we have the following result.

Theorem 7.1. *Let $(M_t)_{t \in \mathbb{R}_+}$ have the representation (7.1), let $(M_t^*)_{t \in \mathbb{R}_+}$ be represented as*

$$M_t^* = \int_t^{+\infty} H_s^* d^* W_s^* + \int_t^{+\infty} J_s^* (d^* Z_s^* - \lambda_s^* ds),$$

assume that $(M_t)_{t \in \mathbb{R}_+}$ is an \mathcal{F}_t^ -adapted \mathcal{F}_t -martingale and that $(M_t^*)_{t \in \mathbb{R}_+}$ is an \mathcal{F}_t -adapted \mathcal{F}_t^* -martingale. Then we have*

$$\mathbb{E}[\phi(M_t + M_t^*)] \leq \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t,$$

for all \mathcal{C}^2 convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$, provided any of the three following conditions is satisfied:

i) $0 \leq \gamma_t R_t \leq J_t^$, $1_{\{\gamma_t=0\}} |R_t|^2 \leq |H_t^*|^2$, and*

$$1_{\{\gamma_t \neq 0\}} \frac{R_t}{\gamma_t} \leq \lambda_t^* J_t^*, \quad dPdt - a.e.,$$

ii) $\gamma_t R_t \leq J_t^$, $1_{\{\gamma_t=0\}} |R_t|^2 \leq |H_t^*|^2$, and*

$$1_{\{\gamma_t \neq 0\}} |R_t|^2 \leq \lambda_t^* |J_t^*|^2, \quad dPdt - a.e.,$$

and ϕ'' is non-decreasing,

iii) $\gamma_t R_t \leq 0$, $|R_t|^2 \leq |H_t^|^2$, $J_t^* = 0$, $dPdt - a.e.$, and ϕ'' is non-decreasing.*

As above, if further $E[M_t^* | \mathcal{F}_t^M] = 0$, $t \in \mathbb{R}_+$, we obtain

$$\mathbb{E}[\phi(M_t)] \leq \mathbb{E}[\phi(M_s + M_s^*)], \quad 0 \leq s \leq t.$$

As a consequence we have the following result which admits the same proof as Theorem 4.1.

Theorem 7.2. *Let $F \in L^2(\Omega, \mathcal{F}, P)$ have the predictable representation*

$$F = \mathbb{E}[F] + \int_0^{+\infty} R_t dZ_t.$$

i) *Assume that $0 \leq \gamma_t R_t \leq k$, $dPdt$ -a.e., for some $k > 0$, and let*

$$\beta_1^2 = \left\| \int_0^{+\infty} 1_{\{\gamma_s=0\}} |R_s|^2 ds \right\|_{\infty} \quad \text{and} \quad \alpha_1 = \left\| \int_0^{+\infty} 1_{\{\gamma_s \neq 0\}} \frac{R_s}{\gamma_s} ds \right\|_{\infty}.$$

Then we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(W(\beta_1^2) + k\tilde{N}(\alpha_1/k) \right) \right],$$

for all convex function $\phi : \mathbb{R} \rightarrow \mathbb{R}$.

ii) *Assume that $\gamma_u R_u \leq k$, $dPdt$ -a.e., for some $k > 0$ and*

$$\beta_2^2 = \left\| \int_0^{+\infty} 1_{\{\gamma_s=0\}} |R_s|^2 ds \right\|_{\infty} \quad \text{and} \quad \alpha_2^2 = \left\| \int_0^{+\infty} 1_{\{\gamma_s \neq 0\}} |R_s|^2 ds \right\|_{\infty}.$$

Then for all \mathcal{C}^2 convex function with non-decreasing second derivative ϕ , we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E} \left[\phi \left(W(\beta_2^2) + k\tilde{N}(\alpha_2^2/k^2) \right) \right].$$

iii) *Assume that $\gamma_u R_u \leq 0$ and let*

$$\beta_3^2 = \left\| \int_0^{+\infty} |R_s|^2 ds \right\|_{\infty}.$$

Then for all \mathcal{C}^2 convex function with non-decreasing second derivative ϕ , we have

$$\mathbb{E}[\phi(F - \mathbb{E}[F])] \leq \mathbb{E}[\phi(\hat{W}(\beta_3^2))].$$

Let now

$$D : L^2(\Omega, \mathcal{F}, P) \mapsto L^2(\Omega \times [0, T], dP \times dt)$$

denote the annihilation operator on multiple stochastic integrals defined as $D_t = I_n(f_n) = nI_n(f_n(*, t))$, $t \in \mathbb{R}_+$. The Clark formula for normal martingales [9] provides a predictable representation for $F \in \text{Dom}(D) \subset L^2(\Omega, \mathcal{F}, P)$, which can be used in Theorem 7.2:

$$F = \mathbb{E}[F] + \int_0^{+\infty} \mathbb{E}[D_t F | \mathcal{F}_t] dZ_t,$$

where $\mathcal{F}_t = \sigma(Z_s, 0 \leq s \leq t)$.

8 Appendix

In this section we prove the Itô type change of variable formula for forward/backward martingales which has been used in the proofs of Theorem 3.2 and Theorem 3.3. Assume that (Ω, \mathcal{F}, P) is equipped with an increasing filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ and a decreasing filtration $(\mathcal{F}_t^*)_{t \in \mathbb{R}_+}$.

Theorem 8.1. *Consider $(M_t)_{t \in \mathbb{R}_+}$ an \mathcal{F}_t^* -adapted, \mathcal{F}_t -forward martingale with right-continuous paths and left limits, and $(M_t^*)_{t \in \mathbb{R}_+}$ an \mathcal{F}_t -adapted, \mathcal{F}_t^* -backward martingale with left-continuous paths and right limits, whose characteristics have the form (3.1) and (3.2). For all $f \in C^2(\mathbb{R}^2, \mathbb{R})$ we have*

$$\begin{aligned} & f(M_t, M_t^*) - f(M_0, M_0^*) \\ &= \int_{0+}^t \frac{\partial f}{\partial x_1}(M_{u-}, M_u^*) dM_u + \frac{1}{2} \int_0^t \frac{\partial^2 f}{\partial x_1^2}(M_u, M_u^*) d\langle M^c, M^c \rangle_u \\ &+ \sum_{0 < u \leq t} \left(f(M_u, M_u^*) - f(M_{u-}, M_u^*) - \Delta M_u \frac{\partial f}{\partial x_1}(M_{u-}, M_u^*) \right) \\ &- \int_0^{t-} \frac{\partial f}{\partial x_2}(M_u, M_{u+}^*) d^* M_u^* - \frac{1}{2} \int_0^t \frac{\partial^2 f}{\partial x_2^2}(M_u, M_u^*) d\langle M^{*c}, M^{*c} \rangle_u \\ &- \sum_{0 \leq u < t} \left(f(M_u, M_u^*) - f(M_u, M_{u+}^*) - \Delta M_u^* \frac{\partial f}{\partial x_2}(M_u, M_{u+}^*) \right), \end{aligned}$$

where d^* denotes the backward Itô differential and $(M_t^c)_{t \in \mathbb{R}_+}$, $(M_t^{*c})_{t \in \mathbb{R}_+}$ respectively denote the continuous parts of $(M_t)_{t \in \mathbb{R}_+}$, $(M_t^*)_{t \in \mathbb{R}_+}$.

Proof. We adapt the arguments of Theorem 32 in [16], using here the following version of Taylor's formula:

$$\begin{aligned}
f(y_1, y_2) - f(x_1, x_2) &= f(y_1, y_2) - f(y_1, x_2) + f(y_1, x_2) - f(x_1, x_2) \quad (8.1) \\
&= (y_1 - x_1) \frac{\partial f}{\partial x_1}(x_1, x_2) + \frac{1}{2}(y_1 - x_1)^2 \frac{\partial^2 f}{\partial x_1^2}(x_1, x_2) \\
&\quad + (y_2 - x_2) \frac{\partial f}{\partial x_2}(y_1, y_2) - \frac{1}{2}(y_2 - x_2)^2 \frac{\partial^2 f}{\partial x_2^2}(y_1, y_2) \\
&\quad + R(x, y),
\end{aligned}$$

where $R(x, y) \leq |y - x|^2 o(|y - x|)$. Assume first that $(M_s)_{s \in [0, t]}$ and $(M_s^*)_{s \in [0, t]}$ take their values in a bounded interval, and let $\{0 = t_0^n \leq t_1^n \leq \dots \leq t_{k_n}^n = t\}$, $n \geq 1$, be a refining sequence of partitions of $[0, t]$ tending to the identity. As in [16], for any $\varepsilon > 0$, consider $A_{\varepsilon, t}$, $B_{\varepsilon, t}$ two random subsets of $[0, t]$ such that

- i) $A_{\varepsilon, t}$ is finite, a.s.,
- ii) $A_{\varepsilon, t} \cup B_{\varepsilon, t}$ exhausts the jumps of $(M_s)_{s \in [0, t]}$ and $(M_s^*)_{s \in [0, t]}$,
- iii) $\sum_{s \in B_{\varepsilon, t}} |\Delta M_s|^2 + |\Delta^* M_s^*|^2 \leq \varepsilon^2$,
- iv) for all $1 \leq i \leq n$ we have either $A_{\varepsilon, t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset$ or $B_{\varepsilon, t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset$, P -a.s.

We have

$$\begin{aligned}
f(M_t, M_t^*) - f(M_0, M_0^*) &= \sum_{A_{\varepsilon, t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} f(M_{t_i^n}, M_{t_i^n}^*) - f(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) \\
&\quad + \sum_{B_{\varepsilon, t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} f(M_{t_i^n}, M_{t_i^n}^*) - f(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*),
\end{aligned}$$

and from Taylor's formula (8.1) we get

$$\begin{aligned}
&f(M_t, M_t^*) - f(M_0, M_0^*) \\
&= \sum_{A_{\varepsilon, t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} f(M_{t_i^n}, M_{t_i^n}^*) - f(M_{t_i^n}, M_{t_{i-1}^n}^*) + f(M_{t_i^n}, M_{t_{i-1}^n}^*) - f(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) \\
&\quad + \sum_{B_{\varepsilon, t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} f(M_{t_i^n}, M_{t_i^n}^*) - f(M_{t_i^n}, M_{t_{i-1}^n}^*) + f(M_{t_i^n}, M_{t_{i-1}^n}^*) - f(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{A_{\varepsilon,t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} f(M_{t_i^n}, M_{t_i^n}^*) - f(M_{t_i^n}, M_{t_{i-1}^n}^*) + f(M_{t_i^n}, M_{t_{i-1}^n}^*) - f(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) \\
&+ \sum_{B_{\varepsilon,t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} (M_{t_i^n} - M_{t_{i-1}^n}) \frac{\partial f}{\partial x_1}(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) + \frac{1}{2} |M_{t_i^n} - M_{t_{i-1}^n}|^2 \frac{\partial^2 f}{\partial x_1^2}(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) \\
&+ \sum_{B_{\varepsilon,t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} (M_{t_i^n}^* - M_{t_{i-1}^n}^*) \frac{\partial f}{\partial x_2}(M_{t_i^n}, M_{t_i^n}^*) - \frac{1}{2} |M_{t_i^n}^* - M_{t_{i-1}^n}^*|^2 \frac{\partial^2 f}{\partial x_2^2}(M_{t_i^n}, M_{t_i^n}^*) \\
&+ \sum_{B_{\varepsilon,t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} R(M_{t_i^n}, M_{t_i^n}^*, M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) \\
&= \sum_{A_{\varepsilon,t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} f(M_{t_i^n}, M_{t_i^n}^*) - f(M_{t_i^n}, M_{t_{i-1}^n}^*) + f(M_{t_i^n}, M_{t_{i-1}^n}^*) - f(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) \\
&+ \sum_{i=1}^n (M_{t_i^n} - M_{t_{i-1}^n}) \frac{\partial f}{\partial x_1}(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) + \frac{1}{2} |M_{t_i^n} - M_{t_{i-1}^n}|^2 \frac{\partial^2 f}{\partial x_1^2}(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) \\
&- \sum_{A_{\varepsilon,t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} (M_{t_i^n} - M_{t_{i-1}^n}) \frac{\partial f}{\partial x_1}(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) + \frac{1}{2} |M_{t_i^n} - M_{t_{i-1}^n}|^2 \frac{\partial^2 f}{\partial x_1^2}(M_{t_{i-1}^n}, M_{t_{i-1}^n}^*) \\
&+ \sum_{i=1}^n (M_{t_i^n}^* - M_{t_{i-1}^n}^*) \frac{\partial f}{\partial x_2}(M_{t_i^n}, M_{t_i^n}^*) - \frac{1}{2} |M_{t_i^n}^* - M_{t_{i-1}^n}^*|^2 \frac{\partial^2 f}{\partial x_2^2}(M_{t_i^n}, M_{t_i^n}^*) \\
&- \sum_{A_{\varepsilon,t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} (M_{t_i^n}^* - M_{t_{i-1}^n}^*) \frac{\partial f}{\partial x_2}(M_{t_i^n}, M_{t_i^n}^*) - \frac{1}{2} |M_{t_i^n}^* - M_{t_{i-1}^n}^*|^2 \frac{\partial^2 f}{\partial x_2^2}(M_{t_i^n}, M_{t_i^n}^*) \\
&+ \sum_{B_{\varepsilon,t} \cap (t_{i-1}^n, t_i^n] \neq \emptyset} R(M_{t_i^n}, M_{t_i^n}^*, M_{t_{i-1}^n}, M_{t_{i-1}^n}^*).
\end{aligned}$$

By the same arguments as in [16], letting n tend to infinity we get

$$\begin{aligned}
&f(M_t, M_t^*) - f(M_0, M_0^*) \\
&= \sum_{u \in A_{\varepsilon,t}} \left(f(M_u, M_u^*) - f(M_{u-}, M_u^*) - \Delta M_u \frac{\partial f}{\partial x_1}(M_{u-}, M_u^*) - \frac{1}{2} |\Delta M_u|^2 \frac{\partial^2 f}{\partial x_1^2}(M_{u-}, M_u^*) \right) \\
&- \sum_{u \in A_{\varepsilon,t}} \left(f(M_u, M_{u+}^*) - f(M_u, M_u^*) - \Delta^* M_u^* \frac{\partial f}{\partial x_2}(M_u, M_{u+}^*) + \frac{1}{2} |\Delta^* M_u^*|^2 \frac{\partial^2 f}{\partial x_2^2}(M_u, M_{u+}^*) \right) \\
&+ \int_0^t \frac{\partial f}{\partial x_1}(M_{u-}, M_u^*) dM_u + \frac{1}{2} \int_0^t \frac{\partial^2 f}{\partial x_1^2}(M_{u-}, M_u^*) d[M, M]_u \\
&- \int_0^t \frac{\partial f}{\partial x_2}(M_u, M_{u+}^*) d^* M_u^* - \frac{1}{2} \int_0^t \frac{\partial^2 f}{\partial x_2^2}(M_u, M_{u+}^*) d[M^*, M^*]_u.
\end{aligned}$$

Then letting ε tend to 0, the above sum converges to

$$f(M_t, M_t^*) - f(M_0, M_0^*)$$

$$\begin{aligned}
&= \int_{0+}^t \frac{\partial f}{\partial x_1}(M_{u-}, M_u^*) dM_u + \frac{1}{2} \int_{0+}^t \frac{\partial^2 f}{\partial x_1^2}(M_{u-}, M_u^*) d[M, M]_u \\
&+ \sum_{0 < u \leq t} \left(f(M_u, M_u^*) - f(M_{u-}, M_u^*) - \Delta M_u \frac{\partial f}{\partial x_1}(M_{u-}, M_u^*) - \frac{1}{2} |\Delta M_u|^2 \frac{\partial^2 f}{\partial x_1^2}(M_{u-}, M_u^*) \right) \\
&- \int_0^{t-} \frac{\partial f}{\partial x_2}(M_u, M_{u+}^*) d^* M_u^* - \frac{1}{2} \int_0^{t-} \frac{\partial^2 f}{\partial x_2^2}(M_u, M_{u+}^*) d[M^*, M^*]_u \\
&- \sum_{0 \leq u < t} \left(f(M_u, M_u^*) - f(M_u, M_{u+}^*) - \Delta^* M_u^* \frac{\partial f}{\partial x_2}(M_u, M_{u+}^*) + \frac{1}{2} |\Delta^* M_u^*|^2 \frac{\partial^2 f}{\partial x_2^2}(M_u, M_{u+}^*) \right),
\end{aligned}$$

which yields

$$\begin{aligned}
f(M_t, M_t^*) - f(M_0, M_0^*) &= \int_{0+}^t \frac{\partial f}{\partial x_1}(M_{u-}, M_u^*) dM_u + \frac{1}{2} \int_0^t \frac{\partial^2 f}{\partial x_1^2}(M_u, M_u^*) d\langle M^c, M^c \rangle_u \\
&+ \sum_{0 < u \leq t} \left(f(M_u, M_u^*) - f(M_{u-}, M_u^*) - \Delta M_u \frac{\partial f}{\partial x_1}(M_{u-}, M_u^*) \right) \\
&- \int_0^{t-} \frac{\partial f}{\partial x_2}(M_u, M_{u+}^*) d^* M_u^* - \frac{1}{2} \int_0^t \frac{\partial^2 f}{\partial x_2^2}(M_u, M_u^*) d\langle M^{*c}, M^{*c} \rangle_u \\
&- \sum_{0 \leq u < t} \left(f(M_u, M_u^*) - f(M_u, M_{u+}^*) - \Delta^* M_u^* \frac{\partial f}{\partial x_2}(M_u, M_{u+}^*) \right),
\end{aligned}$$

where the integral with respect to $(\langle M^*, M^* \rangle_t)_{t \in \mathbb{R}_+}$ is defined as a Stieltjes integral with respect to a (not \mathcal{F}_t^* -adapted) increasing process. In the general case, define the stopping times

$$R_m = \inf\{u \in [0, t] : |M_u| \geq m\}, \quad \text{and} \quad R_m^* = \sup\{u \in [0, t] : |M_u^*| \geq m\}.$$

The stopped process $(M_{u \wedge R_m}, M_{u \vee R_m}^*)_{u \in [0, t]}$ is bounded by $2m$ and since Itô's formula is valid for $(X_u^{R_m})_{u \in [0, t]}$ for each m , it is also valid for $(X_u)_{u \in \mathbb{R}_+}$. \square

Note that the cross partial derivative $\frac{\partial^2 f}{\partial x_1 \partial x_2}(M_u, M_u^*)$ does not appear in the formula and there is no need to consider or define a bracket of the form $d\langle M, M^* \rangle_t$.

References

- [1] C. Ané and M. Ledoux. On logarithmic Sobolev inequalities for continuous time random walks on graphs. *Probab. Theory Related Fields*, 116(4):573–602, 2000.
- [2] V. Bentkus. On Hoeffding's inequalities. *Ann. Probab.*, 32(2):1650–1673, 2004.

- [3] C. Dellacherie, B. Maisonneuve, and P.A. Meyer. *Probabilités et Potentiel*, volume 5. Hermann, 1992.
- [4] M. Émery. On the Azéma martingales. In *Séminaire de Probabilités XXIII*, volume 1372 of *Lecture Notes in Mathematics*, pages 66–87. Springer Verlag, 1990.
- [5] J. Jacod. *Calcul stochastique et problèmes de martingales*, volume 714 of *Lecture Notes in Mathematics*. Springer Verlag, 1979.
- [6] J. Jacod and J. Mémin. Caractéristiques locales et conditions de continuité absolue pour les semi-martingales. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 35(1):1–37, 1976.
- [7] O. Kallenberg. *Foundations of modern probability*. Probability and its Applications. Springer-Verlag, New York, 2002.
- [8] Th. Klein. Inégalités de concentration, martingales et arbres aléatoires. Thèse, Université de Versailles-Saint-Quentin, 2003.
- [9] J. Ma, Ph. Protter, and J. San Martin. Anticipating integrals for a class of martingales. *Bernoulli*, 4:81–114, 1998.
- [10] D. Nualart and J. Vives. Anticipative calculus for the Poisson process based on the Fock space. In *Séminaire de Probabilités XXIV*, volume 1426 of *Lecture Notes in Math.*, pages 154–165. Springer, Berlin, 1990.
- [11] B. Øksendal and F. Proske. White noise of Poisson random measures. *Potential Anal.*, 21(4):375–403, 2004.
- [12] J. Picard. Formules de dualité sur l’espace de Poisson. *Ann. Inst. H. Poincaré Probab. Statist.*, 32(4):509–548, 1996.
- [13] I. Pinelis. Optimal tail comparison based on comparison of moments. In *High dimensional probability (Oberwolfach, 1996)*, volume 43 of *Progr. Probab.*, pages 297–314. Birkhäuser, Basel, 1998.
- [14] N. Privault. An extension of stochastic calculus to certain non-Markovian processes. Prépublication 49, Université d’Evry, 1997.
- [15] N. Privault. Equivalence of gradients on configuration spaces. *Random Operators and Stochastic Equations*, 7(3):241–262, 1999.
- [16] Ph. Protter. *Stochastic integration and differential equations. A new approach*. Springer-Verlag, Berlin, 1990.
- [17] L. Wu. A new modified logarithmic Sobolev inequality for Poisson point processes and several applications. *Probab. Theory Related Fields*, 118(3):427–438, 2000.