

# Three dimensional modeling and simulations of ionospheric plasma instabilities

Christophe Besse, Pierre Degond, Fabrice Deluzet

Laboratoire MIP, Toulouse, France


Summer school on Multiscale Modeling and Applications,  
7–12 August 2006, Cargèse (Corsica), France

# Outline

- 1 Physical context, motivations
  - The earth environment
  - Motivations of the study
- 2 Modeling
- 3 The Striation model
- 4 The 3D-Dynamo model



# The earth environment



- the atmosphere is a stratified and ionized medium,
- the ionosphere is characterized by
  - heights ranging from 90 to 1500 kilometers,
  - a neutral density much larger than the plasma density,
  - a maximum in the plasma density  $10^6 \text{ cm}^{-3}$  in the  $\sim 300 \text{ km}$  heights area (the F-region).

Fig. 1: Stratification of the earth atmosphere.

# Properties of the ionosphere





Fig. 2: Radio waves transmissions and interaction with the ionosphere.

The ionospheric plasma :


- reflects low frequency waves,
- transmits high frequency waves,
- is subjected to instabilities (solar eruptions, striations, ...).

**Striations** : ionospheric plasma instability observed along the earth magnetic field lines.

# Striations formations



Time evolution of a plasma bubble located at a 300 km altitude.



The plasma bubble is progressively tired up along the magnetic field lines. In a plane perpendicular to the magnetic field the plasma bubble frontier is submitted to an instability referred to as " $E \times B$  instability".



# Outline

- 1 Physical context, motivations
- 2 Modeling
  - The dynamo effect
  - The Euler-Maxwell system
  - A model hierarchy
- 3 The Striation model
- 4 The 3D-Dynamo model








Fig. 4: The Dynamo effect.

The model :

- Maxwell equations,
- Euler equations with collisions against neutrals.

Simplifications

- only one ion species :  $O^+$ ,
- no chemical reaction,
- electron-ions collisions and gravity effect are neglected.

# Notations

$n_e, n_i$	electronic, ionic density
$u_e, u_i$	electronic, ionic velocity
$u_n$	neutral velocity
$P_e, P_i$	electronic, ionic pressure
$\nu_e, \nu_i$	electronic-neutral, ionic-neutral collision frequency
$m_e, m_i$	electronic, ionic mass
$j = e(n_i u_i - n_e u_e)$	current density
$\rho_c = e(n_i - n_e)$	charge density

# The Euler-Maxwell system

## Euler equations

$$\mathcal{L}_{u_e}(n_e) = 0,$$

$$m_e \mathcal{L}_{u_e}(n_e u_{ek}) = -\partial_{x_k} P_e - \overbrace{en_e(E_k + (u_e \times B)_k)}^{\text{Lorentz}} + n_e F_{ek},$$

$$\mathcal{L}_{u_i}(n_i) = 0,$$

$$m_i \mathcal{L}_{u_i}(n_i u_{ik}) = -\partial_{x_k} P_i + qn_i(E_k + (u_i \times B)_k) + n_i F_{ik},$$

## The Maxwell system

$$\frac{1}{c^2} \partial_t E - \nabla \times B = -\mu_0 j,$$

$$\partial_t B + \nabla \times E = 0,$$

$$\nabla \cdot E = \rho_c / \epsilon_0,$$

$$\nabla \cdot B = 0,$$

$$\rho_c = q(n_i - n_e), \quad j = q(n_i u_i - n_e u_e).$$

With the following definition for the transport operator :

$$\mathcal{L}_u(\rho) = \partial_t \rho + \nabla \cdot (\rho u),$$

and the friction forces :

$$F_e = -\nu_e m_e (u_e - u_n),$$

$$F_i = -\nu_i m_i (u_i - u_n).$$

# Scaling relations

Physical parameter	Typical scale	Dimensionless parameter	Typical value
Time	$\bar{t}$	$t' = t/\bar{t}$	$10^3$ s
Length	$\bar{x}$	$x' = x/\bar{x}$	$10^5$ m
Speed	$\bar{u} = \bar{x}/\bar{t}$	$u'_{e,i,n} = u_{e,i,n}/\bar{u}$	$10^2$ ms <sup>-1</sup>
Density	$\bar{n}$	$n'_{e,i} = n_{e,i}/\bar{n}$	$10^{12} \mid 10^{15}$ m <sup>-3</sup>
Magnetic field	$\bar{B}$	$B' = B/\bar{B}$	$10^{-5}$ T
Electric field	$\bar{E} = \bar{u}\bar{B}$	$E' = E/\bar{E}$	$10^{-3}$ Vm <sup>-1</sup>
e-n collision freq.	$\bar{\nu}_e$	$\nu'_e = \nu_e/\bar{\nu}_e$	$10^2$ s <sup>-1</sup>
i-n collision freq.	$\bar{\nu}_i = \frac{m_e}{m_i} \bar{\nu}_e$	$\nu'_i = \nu_i/\bar{\nu}_i$	$10^{-2}$ s <sup>-1</sup>

## The rescaled Euler-Maxwell system

# Some dimensionless parameters

- $\beta$  intensity of the induced magnetic field compared to the ambient earth magnetic field,
- $\kappa$  number of e-n or i-n collisions during a cyclotron period,
- $\varepsilon$  ratio of the electronic and ionic masses,
- $\alpha$  ratio of the typical velocity and the speed of light,
- $\tau$  ratio of the typical time and the time between two ions-neutral collisions,
- $\eta$  measure of the thermic energy.

# The rescaled Euler-Maxwell system

## Euler equations

$$\partial_t n_e + \nabla \cdot (n_e u_e) = 0,$$

$$\tau \varepsilon (\mathcal{L}_{u_e}(n_e u_{ek})) = -\eta \partial_{x_k} P_e - \kappa^{-1} \overbrace{n_e (E_k + (u_e \times B)_k)}^{\text{Lorentz}} - \overbrace{\nu_e n_e (u_{ek} - u_{nk})}^{\text{friction}},$$

$$\partial_t n_i + \nabla \cdot (n_i u_i) = 0,$$

$$\tau (\mathcal{L}_{u_i}(n_i u_{ik})) = -\eta \partial_{x_k} P_i + \kappa^{-1} n_i (E_k + (u_i \times B)_k) - \nu_i n_i (u_{ik} - u_{nk}),$$

## Maxwell system

$$\alpha \partial_t E - \nabla \times B = -\beta j,$$

$$\partial_t B + \nabla \times E = 0,$$

$$\frac{\kappa \alpha}{\beta} \nabla \cdot E = \rho_c,$$

$$\nabla \cdot B = 0,$$

$$\rho_c = n_i - n_e, \quad \kappa j = n_i u_i - n_e u_e.$$

Typical values of the plasma density  
 $n_{i,e} = 10^{12} | 10^{15} m^{-3}$

$$\begin{aligned} \varepsilon &= 10^{-4}, & \tau &= 10^{-1}, \\ \eta &= 10^1, & \kappa &= 10^{-4} \\ \alpha &= 10^{-12}, & \beta &= 10^{-5} | 10^{-2}. \end{aligned}$$

# The rescaled Euler-Maxwell system

## Euler equations

$$\partial_t n_e + \nabla \cdot (n_e u_e) = 0,$$

$$\tau \varepsilon (\mathcal{L}_{u_e}(n_e u_{ek})) = -\eta \partial_{x_k} P_e - \kappa^{-1} n_e (E_k + (u_e \times B)_k) - \nu_e n_e (u_{ek} - u_{nk}),$$

$$\partial_t n_i + \nabla \cdot (n_i u_i) = 0,$$

$$\tau (\mathcal{L}_{u_i}(n_i u_{ik})) = -\eta \partial_{x_k} P_i + \kappa^{-1} n_i (E_k + (u_i \times B)_k) - \nu_i n_i (u_{ik} - u_{nk}),$$

## Maxwell system

$$\alpha \partial_t E - \nabla \times B = -\beta j,$$

$$\partial_t B + \nabla \times E = 0,$$

$$\frac{\kappa \alpha}{\beta} \nabla \cdot E = \rho_c,$$

$$\nabla \cdot B = 0,$$

$$\rho_c = n_i - n_e, \quad \kappa j = n_i u_i - n_e u_e.$$

Typical values of the plasma density

$$n_{i,e} = 10^{12} | 10^{15} m^{-3}$$

$$\varepsilon = 10^{-4}, \quad \tau = 10^{-1},$$

$$\eta = 10^1, \quad \kappa = 10^{-4}$$

$$\alpha = 10^{-12}, \quad \beta = 10^{-5} | 10^{-2}.$$

# The rescaled Euler-Maxwell system

## Euler equations

$$\partial_t n_e + \nabla \cdot (n_e u_e) = 0,$$

$$0 = -\eta \partial_{x_k} P_e - \kappa^{-1} n_e (E_k + (u_e \times B)_k) - \nu_e n_e (u_{ek} - u_{nk}),$$

$$\partial_t n_i + \nabla \cdot (n_i u_i) = 0,$$

$$\tau (\mathcal{L}_{u_i} (n_i u_{ik})) = -\eta \partial_{x_k} P_i + \kappa^{-1} n_i (E_k + (u_i \times B)_k) - \nu_i n_i (u_{ik} - u_{nk}),$$

## Maxwell system

$$\alpha \partial_t E - \nabla \times B = -\beta j,$$

$$\partial_t B + \nabla \times E = 0,$$

$$\frac{\kappa \alpha}{\beta} \nabla \cdot E = \rho_c,$$

$$\nabla \cdot B = 0,$$

$$\rho_c = n_i - n_e, \quad \kappa j = n_i u_i - n_e u_e.$$

Typical values of the plasma density

$$n_{i,e} = 10^{12} | 10^{15} m^{-3}$$

$$\varepsilon = 10^{-4}, \quad \tau = 10^{-1},$$

$$\eta = 10^1, \quad \kappa = 10^{-4}$$

$$\alpha = 10^{-12}, \quad \beta = 10^{-5} | 10^{-2}.$$



# The rescaled Euler-Maxwell system

## Euler equations

$$\partial_t n_e + \nabla \cdot (n_e u_e) = 0,$$

$$0 = -\eta \partial_{x_k} P_e - \kappa^{-1} n_e (E_k + (u_e \times B)_k) - \nu_e n_e (u_{ek} - u_{nk}),$$

$$\partial_t n_i + \nabla \cdot (n_i u_i) = 0,$$

$$\tau (\mathcal{L}_{u_i} (n_i u_{ik})) = -\eta \partial_{x_k} P_i + \kappa^{-1} n_i (E_k + (u_i \times B)_k) - \nu_i n_i (u_{ik} - u_{nk}),$$

## Maxwell system

$$-\nabla \times B = -\beta j,$$

$$\partial_t B + \nabla \times E = 0,$$

$$0 = \rho_c, \Rightarrow \text{quasineutrality}$$

$$\nabla \cdot B = 0,$$

$$\rho_c = n_i - n_e, \quad \kappa j = n_i u_i - n_e u_e.$$

Typical values of the plasma density

$$n_{i,e} = 10^{12} | 10^{15} m^{-3}$$

$$\varepsilon = 10^{-4}, \quad \tau = 10^{-1},$$

$$\eta = 10^1, \quad \kappa = 10^{-4}$$

$$\alpha = 10^{-12}, \quad \beta = 10^{-5} | 10^{-2}.$$




Fig. 5: Model hierarchy.

- 👉 Dynamo hierarchy : standard plasma density,
- 👉 MHD hierarchy : high plasma density.

# Outline

- 1 Physical context, motivations
- 2 Modeling
- 3 The Striation model
  - Derivation of the model
  - Linear stability analysis
  - Formulation in a non uniform magnetic field
  - More accurate physic
- 4 The 3D-Dynamo model

# Main assumptions and properties

## Assumptions :

- 👉 quasi-neutral plasma :  $n_e = n_i = n$ ,
- 👉 the magnetic field is reduced to the earth magnetic field and  $\partial B / \partial t = 0$ . We choose  $B$  to be aligned with  $x_3$  (or  $x_{||}$ ) coordinate,
- 👉 the aligned conductivity is infinite :  $\kappa \rightarrow 0$ ,
- 👉 no incoming or outgoing current flowing through the boundary with normal parallel to  $x_3$  (reconnection to the neutral atmosphere).

## The “Striation model” main properties :

- the electric potential depends only on the two first coordinate (plane perpendicular to the magnetic field  $x_{\perp}$ ),
- by integrating the elliptic equation along the magnetic field line the problem is reduced to a bi-dimensional one.

# The multi-layer “Striation model” (rescaled variables)

$$\begin{aligned}\frac{\partial n}{\partial t} + \nabla \cdot (nu) &= 0 \\ u &= \frac{E \times B}{|B|^2} + \left( \left( u_n - \eta \frac{\nabla p}{n} \right) \cdot \frac{B}{|B|} \right) \frac{B}{|B|}; \quad E = -\nabla_{\perp} \phi(x_{\perp}) \\ \nabla_{\perp} \cdot J_{\perp} &= 0, \\ J_{\perp} &= \frac{1}{|B|^2} (-\sigma(x_{\perp}) \nabla_{\perp} \phi + U_n \times B - \eta \nabla_{\perp} P_{\perp} \times B), \\ \sigma(x_{\perp}) &= \int n \nu dx_3, \quad U_n = \int n \nu u_n dx_3, \quad P_{\perp} = \int P(n) dx_3.\end{aligned}$$

- $\sigma/|B|^2$  : Pedersen conductivity integrated along a magnetic field line,
- 3D transport equation, 2D elliptic equation.

# The 2D “Striations model” (rescaled variables)

## Assumptions :

- 👉 All quantities only depend on  $x_{\perp}$ ,
- 👉  $u_{n3} = 0$  and  $\nabla P = 0$ ,

$$\begin{aligned}\frac{\partial n}{\partial t} + \nabla \cdot (nu) &= 0 \\ -\nabla \cdot \left( \frac{n}{|B|^2} \nabla \phi \right) &= -\nabla \cdot \left( n \frac{u_n \times B}{|B|^2} \right), \\ u &= -\frac{E \times B}{|B|^2}, \\ E &= -\nabla \phi.\end{aligned}$$

# The “ $E \times B$ ” instability

$$\begin{aligned}\partial_t n + \nabla \cdot (nu) &= 0, \\ \nabla \cdot (nh) &= 0, \\ h &= -\nabla\phi - u_n^\perp.\end{aligned}$$

Recall that  $u = (\nabla\phi)^\perp$  and  
 $h = -\nabla\phi - u_n^\perp = u_i - u_e$ .




Fig. 6: Stable configuration




Fig. 7: Unstable configuration

The Striations model is widely used by physicists e.g.

- Ronchi, Similon, Farley (1989),
  - Zalesak, Ossakov, Chaturvedi (1982).
- ☞ Mainly 2D model,
- ☞ poor description of the variations along the third coordinate,
- ☞ Cartesian geometry.




Fig. 8: Magnetic field tube in Cartesian geometry.



# A curvilinear coordinates for non uniform magnetic field.




Fig. 9: Magnetic field tube with Cartesian coordinates.

## Dipolar model for the earth magnetic field :

- Axisymmetric field  
 $B = (B_r(r, z), 0, B_z(r, z))$ ,  
 $(r, \theta, z)$  being the cylindrical coordinates ;

- use of Euler Potentials :  
 $\nabla \cdot B = \nabla \times B = 0 \Rightarrow \exists$   
 $\beta(r, z), \gamma(r, z)$  such that

$$B = -\nabla\gamma = (\nabla^\perp\beta),$$

- magnetic field lines are described by  
 $(\alpha, \beta) = \text{constant}$ .

# A curvilinear coordinate set




Fig. 10: Curvilinear coordinate system associated to the dipolar magnetic field.




Fig. 11: Magnetic field tube using a dipolar model.

# The Striation model in a non-uniform magnetic field (physical variables)

Assumptions : no transport in the direction aligned with the magnetic fields and no pressure term.

$$\frac{\partial n}{\partial t} + \frac{\partial}{\partial \alpha} \left( \frac{v_{i\alpha}}{r} n \right) + |B|^2 \frac{\partial}{\partial \beta} \left( \frac{r v_{i\beta}}{|B|} n \right) = 0,$$

$$v_{i\alpha} = r \frac{\partial \phi}{\partial \beta}, \quad v_{i\beta} = -\frac{1}{r|B|} \frac{\partial \phi}{\partial \alpha},$$

$$\frac{\partial}{\partial \alpha} \left( A_\alpha \frac{\partial \phi}{\partial \alpha} \right) + \frac{\partial}{\partial \beta} \left( A_\beta \frac{\partial \phi}{\partial \beta} \right) = \frac{\partial J_{n\alpha}}{\partial \alpha} + \frac{\partial J_{n\beta}}{\partial \beta},$$

$$A_\alpha = \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta \frac{d\gamma}{r^2 |B|^4}, \quad A_\beta = \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta \frac{r^2 d\gamma}{|B|^2},$$

$$J_{n\alpha} = - \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta u_{n\beta} \frac{d\gamma}{r |B|^3}, \quad J_{n\beta} = \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta u_{n\alpha} \frac{r d\gamma}{|B|^2},$$

$$\zeta = \frac{m_i}{e} \nu_i + \frac{m_e}{e} \nu_e.$$


# Simulations with non-uniform magnetic

**Simulation domain :** One magnetic field tube,

- ☞ The domain is discretized thanks to several layers,
- ☞ each layer being discretized with a Cartesian mesh,


**Initial condition :**

- ☞ earth ionospheric model : IRI (International Reference Ionosphere),
- ☞ earth neutral atmosphere model : MSISE-1990 (Extended version of the Mass Spectrometer Incoherent Scatter Model).



**Fig. 12:** Discretized magnetic field tube.

# Simulations with non-uniform magnetic field



**Fig. 13:** Plasma bubble evolution.  
(The wind blows from the left to the right)

Some assumptions may be relaxed :

- 1 quantities may vary along the magnetic field line,
- 2 a constant ions-neutrals collision frequency is not acceptable : (see Fig. 14),




Fig. 14: Ions-neutrals collision frequency ( $\text{s}^{-1}$ ) versus altitude (km).

# Transport in the aligned direction and pressure term

Assumptions :  $T = T_e = T_i$ , hence  $P = nk_B 2T$ .

Striation model with pressure term and transport in the aligned direction

$$\frac{\partial n}{\partial t} + \frac{\partial}{\partial \alpha} \left( \frac{v_{i\alpha}}{r} n \right) + |B|^2 \left( \frac{\partial}{\partial \beta} \left( \frac{r v_{i\beta}}{|B|} n \right) + \frac{\partial}{\partial \gamma} \left( \frac{v_{i\gamma}}{|B|} n \right) \right) = 0,$$

$$v_{i\alpha} = r \frac{\partial \phi}{\partial \beta}, \quad v_{i\beta} = -\frac{1}{r|B|} \frac{\partial \phi}{\partial \alpha}, \quad v_{i\gamma} = u_{n\gamma} - \frac{|B|}{e n \zeta} \frac{\partial P}{\partial \gamma},$$


$$\frac{\partial}{\partial \alpha} \left( A_\alpha \frac{\partial \phi}{\partial \alpha} \right) + \frac{\partial}{\partial \beta} \left( A_\beta \frac{\partial \phi}{\partial \beta} \right) = \frac{\partial}{\partial \alpha} (J_{n\alpha} + J_{P\alpha}) + \frac{\partial}{\partial \beta} (J_{n\beta} + J_{P\beta}),$$

$$A_\alpha = \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta \frac{d\gamma}{r^2 |B|^4}, \quad A_\beta = \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta \frac{r^2 d\gamma}{|B|^2},$$


$$J_{n\alpha} = - \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta u_{n\beta} \frac{d\gamma}{r |B|^3}, \quad J_{n\beta} = \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta u_{n\alpha} \frac{r d\gamma}{|B|^2},$$

$$J_{P\alpha} = \int_{\gamma_{\min}}^{\gamma_{\max}} \frac{1}{e} \frac{\partial P}{\partial \beta} \frac{d\gamma}{|B|^2}, \quad J_{P\beta} = - \int_{\gamma_{\min}}^{\gamma_{\max}} \frac{1}{e} \frac{\partial P}{\partial \alpha} \frac{d\gamma}{|B|^2}.$$


# Plasma density evolution (“Aligned transport”)




$T=0$



$T = 30 \text{ min}$



$T = 1 \text{ h } 00 \text{ min}$



$T = 1 \text{ h } 30 \text{ min}$



# Non constant i-n collision frequency

## The Striation model

$$\frac{\partial n}{\partial t} + \frac{\partial}{\partial \alpha} \left( \frac{v_{i\alpha}}{r} n \right) + |B|^2 \frac{\partial}{\partial \beta} \left( \frac{r v_{i\beta}}{|B|} n \right) = 0,$$

$$v_{i\alpha} = r \frac{\partial \phi}{\partial \beta}, \quad v_{i\beta} = -\frac{1}{r|B|} \frac{\partial \phi}{\partial \alpha},$$

$$\frac{\partial}{\partial \alpha} \left( A_\alpha \frac{\partial \phi}{\partial \alpha} \right) + \frac{\partial}{\partial \beta} \left( A_\beta \frac{\partial \phi}{\partial \beta} \right) = \frac{\partial J_{n\alpha}}{\partial \alpha} + \frac{\partial J_{n\beta}}{\partial \beta},$$

$$A_\alpha = \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta \frac{d\gamma}{r^2 |B|^4}, \quad A_\beta = \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta \frac{r^2 d\gamma}{|B|^2},$$

$$J_{n\alpha} = - \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta u_{n\beta} \frac{d\gamma}{r |B|^3}, \quad J_{n\beta} = \int_{\gamma_{\min}}^{\gamma_{\max}} n \zeta u_{n\alpha} \frac{r d\gamma}{|B|^2},$$

$$\zeta = \frac{m_i}{e} \nu_i.$$







Fig. 15: Simulation carried out with a **constant** i-n collision frequency.



(a) Simulation domain.



(b) Central layer plasma density ( $\text{m}^{-3}$ ) at  $t=2$  h.




(c)  $n\zeta$  integrated along magnetic field lines.

Fig. 16: Plasma density evolution with a **non constant** i-n collision frequency.


Collisions at low altitudes are very frequent : the plasma is set into motion with the neutral wind without any instability development.

## Solution :


Use the Striation model in a small domain where it is valid.



(a) Simulation Domain.



(b) Central layer plasma density ( $\text{m}^{-3}$ ) at  $t=2$  h.



(c)  $n\kappa$  integrated along magnetic field lines.

Fig. 17: Plasma density with a non-constant i-n collision frequency.

## Conclusion :

Relax the infinite aligned mobility assumption : the Dynamo model.

# Outline

- 1 Physical context, motivations
- 2 Modeling
- 3 The Striation model
- 4 The 3D-Dynamo model
  - Presentation
  - Properties of the Dynamo model
  - Perspectives




Fig. 18: Model hierarchy.

# The 3D Dynamo model

- quasi-neutral plasma :  $n = n_e = n_i$ ,
- the magnetic field reduces to the earth magnetic field,
- on the typical time scale, we assume,  $\partial B / \partial t = 0$ ,

$$\begin{aligned}\partial_t n + \nabla \cdot (n u_i) &= 0, \\ u_e &= \mathbb{M}_e (-E + \kappa \nu_e u_n), \\ u_i &= \mathbb{M}_i (E + \kappa \nu_i u_n), \\ E &= -\nabla \phi, \\ \nabla \cdot j &= 0, \\ \kappa j &= n(u_i - u_e),\end{aligned}$$

$$\mathbb{M}_e = \begin{pmatrix} \mu_e^P & -\mu_e^H & 0 \\ \mu_e^H & \mu_e^P & 0 \\ 0 & 0 & \mu_e^{\parallel} \end{pmatrix},$$

$$\mathbb{M}_i = \begin{pmatrix} \mu_i^P & \mu_i^H & 0 \\ -\mu_i^H & \mu_i^P & 0 \\ 0 & 0 & \mu_i^{\parallel} \end{pmatrix},$$

with

$$\mu_{e,i}^P = \frac{\kappa \nu_{e,i}}{(\kappa \nu_{e,i})^2 + |B|^2}, \quad \mu_{e,i}^H = \frac{|B|}{(\kappa \nu_{e,i})^2 + |B|^2}, \quad \mu_{e,i}^{\parallel} = \frac{1}{\kappa \nu_{e,i}}.$$

# The 3D-Dynamo model

Expression in Cartesian geometry.


$$\begin{aligned} \frac{\partial n}{\partial t} + \nabla \cdot (nu_i) &= 0, \\ -\nabla \cdot \left[ n \left( \mathbb{M}_i + \mathbb{M}_e \right) \nabla \phi \right] &= -\nabla \cdot \left[ n \mathbb{M}_i \left( \kappa_i u_n - \frac{\nabla P_i}{e n} \right) - n \mathbb{M}_e \left( \kappa_e u_n - \frac{\nabla P_e}{e n} \right) \right], \\ u_i &= \mathbb{M}_i \left( E + \kappa_i u_n - \frac{\nabla P_i}{e n} \right), \quad E = -\nabla \phi, \\ \mu_{e,i}^P &= \frac{\kappa_{e,i}}{\kappa_{e,i}^2 + |B|^2}, \quad \mu_{e,i}^H = \frac{|B|}{\kappa_{e,i}^2 + |B|^2}, \quad \mu_{e,i}^{\parallel} = \frac{1}{\kappa_{e,i}}, \quad \kappa_{e,i} = \frac{m_{e,i} \nu_{e,i}}{e}, \\ \mathbb{M}_e &= \begin{pmatrix} \mu_e^P & -\mu_e^H & 0 \\ \mu_e^H & \mu_e^P & 0 \\ 0 & 0 & \mu_e^{\parallel} \end{pmatrix}, \quad \mathbb{M}_i = \begin{pmatrix} \mu_i^P & \mu_i^H & 0 \\ -\mu_i^H & \mu_i^P & 0 \\ 0 & 0 & \mu_i^{\parallel} \end{pmatrix} \end{aligned}$$



# The 3D-Dynamo model

- 👉 The Dynamo model does not assume on infinite aligned conductivity.
  - 👉 The Dynamo model is valid on the whole range of altitudes.
- 
- The 3D-Dynamo model is a full 3D model : the electric potential is the solution of a three dimensional elliptic equation.
  - Furthermore, the discretized equation is ill-conditioned due to the large value of aligned mobilities compared to Pedersen and Hall mobilities.

## III-conditionness of the discretized 3D-Dynamo elliptic equation



(a) Perdersen and aligned mobilities ratio :  $\mu^P / \mu^\parallel$ .

(b) Ratio of the Hall and aligned mobilities  $\mu^H / \mu^\parallel$ .

3D-Dynamo elliptic equation :

$$-\nabla \cdot (\mathbb{M} \nabla \phi) = -\nabla \cdot \mathbf{J},$$

with

$$\mathbb{M} = \mathbb{M}_e + \mathbb{M}_i$$

$$= \begin{pmatrix} \mu^P & -\mu^H & 0 \\ \mu^H & \mu^P & 0 \\ 0 & 0 & \mu^\parallel \end{pmatrix},$$

and

$$\mu^P = \mu_e^P + \mu_i^P,$$

$$\mu^H = \mu_e^H - \mu_i^H,$$

$$\mu^\parallel = \mu_e^\parallel + \mu_i^\parallel.$$

Fig. 19: Mobilities ratios in decimal log scale.

## III-conditionness of the discretized 3D-Dynamo elliptic equation

Solve :  $\mathcal{A}_h \phi_h = b_h$ , a discretization of  $-\nabla \cdot (\mathbb{M} \nabla \phi) = b$  (+ Boundary cond.).  
Then compute the residual  $R = (\mathcal{A}_h \phi_h - b_h)/b_h$  for  $b_h = 1$  ( $R_1$ ) and  
 $b_h \approx -\nabla \cdot J$  ( $R_J$ ).




Fig. 20: Residual norms versus aligned mobility multiplier. Linear system solver : ILU preconditioned conjugate gradient method, with  $50 \times 50 \times 50$  nodes.

# Models coupling




Fig. 21: Domain decomposition.

## Strategy :

- Striation model for high altitudes :  $\Omega_S \rightarrow \phi^S$ ,
- 3D-Dynamo model for lower altitudes  $\Omega_i \rightarrow \phi^i$ ,  $i = 1, 2$ ,
- Matching conditions on the boundaries  $\Sigma_i$ ,  $i = 1, 2$   

$$\phi^S|_{\Sigma_i} = \phi^i|_{\Sigma_i} .$$