

ON LOCAL CONVERGENCE OF THE METHOD OF ALTERNATING PROJECTIONS

DOMINIKUS NOLL*, AUDE RONDEPIERRE†

ABSTRACT. The method of alternating projections is a classical tool to solve feasibility problems. Here we prove local convergence of alternating projections between subanalytic sets A, B under a mild regularity hypothesis on one of the sets. We show that the speed of convergence is $\mathcal{O}(k^{-\rho})$ for some $\rho \in (0, \infty)$.

Keywords. Alternating projections · local convergence · subanalytic set · separable intersection · tangential intersection · Hölder regularity · Gerchberg-Saxton error reduction

Mathematics Subject Classification (2000). Primary: 65K10. Secondary: 90C30, 32B20, 47H04, 49J52

1. INTRODUCTION

The method of alternating projections is a classical tool to solve the following feasibility problem: Given closed sets A, B in \mathbb{R}^n , find a point $x^* \in A \cap B$. Alternating projections can be traced back to the work of Schwarz [26] in 1869, and were popularized in lecture notes of von Neumann [23] since the 1930s. The method generates sequences $a_k \in P_A(b_{k-1})$, $b_k \in P_B(a_k)$, where P_A, P_B are the set-valued orthogonal projection operators on A and B . If the alternating sequence a_k, b_k is bounded and satisfies $a_k - b_k \rightarrow 0$, then each of its accumulation points is a solution of the feasibility problem. The fundamental question is when such a sequence converges to a single limit point $x^* \in A \cap B$.

For convex sets alternating projections are globally convergent as soon as $A \cap B \neq \emptyset$, and the survey [2] gives an excellent state-of-art of the convex theory. In one of the earliest contributions to the nonconvex case, Combettes and Trussell [11] proved in 1990 that the set of accumulation points of a bounded sequence of alternating projections with $a_k - b_k \rightarrow 0$ is either a singleton or a nontrivial compact continuum. In 2013 it was shown in [6] by way of an example that the continuum case may indeed occur. This shows that without convexity a sequence of alternating projections a_k, b_k may fail to converge even when it is bounded and satisfies $a_k - b_k \rightarrow 0$.

In 2008 Lewis and Malick [20] proved that a sequence a_k, b_k of alternating projections converges locally linearly if A, B are C^2 -manifolds intersecting transversally. Expanding on this in 2009, Lewis *et al.* [21] proved local linear convergence for general A, B intersecting non-tangentially in the sense of linear regularity, where one of the sets is superregular. In 2013 Bauschke *et al.* [4, 5] investigate the case of non-tangential intersection further and prove linear convergence under weaker regularity and transversality hypotheses.

Here we prove local convergence under less restrictive conditions, where A, B may also intersect tangentially. We propose a new geometric concept, called *separable intersection*, which gives local convergence of alternating projections when combined with *Hölder regularity*, a mild hypothesis less restrictive than prox-regularity.

Communicated by Michael Todd.

*Université de Toulouse, Institut de Mathématiques, 118, route de Narbonne, 31062 Toulouse, France, Tel. 0033 5 61 55 86 22, Fax. 0033 5 61 55 83 85. E-mail: noll@math.univ-toulouse.fr. Corresponding author.

†Université de Toulouse, Institut de Mathématiques, 118 route de Narbonne, 31062 Toulouse, France. Tel. 0033 5 61 55 76 48. Fax. 0033 5 61 55 75 99. E-mail: aude.rondepierre@insa-toulouse.fr.

Separable intersection has wide scope for applications, as it not only includes non-tangential intersection, but goes beyond and allows also a large variety of cases where A, B intersect tangentially. In particular, we prove that closed subanalytic sets A, B *always* intersect separably. This leads to the central result that alternating projections between subanalytic sets converge locally with rate $\mathcal{O}(k^{-\rho})$ for some $\rho \in (0, \infty)$ if one of the sets is Hölder regular with respect to the other. As these hypotheses are satisfied in practical situations, we obtain a theoretical explanation for the fact, observed in practice, that even without convexity alternating projections converge well in the neighborhood of $A \cap B$. As an application, we obtain a local convergence proof for the classical Gerchberg-Saxton error reduction algorithm in phase retrieval.

The structure of the paper is as follows. Section 3 introduces the concept of separable intersection of two closed sets. Then 0-separability is related to existing transversality concepts. In section 4 we discuss Hölder regularity and compare it to older regularity concepts like prox-regularity, Clarke regularity, and superregularity. The central section 5 gives the convergence proof with rate for sets intersecting separably. In section 6 we show that subanalytic sets intersect separably and then deduce the convergence result for subanalytic sets. Section 6 gives also some applications indicating the versatility of our convergence test. In particular, we prove local convergence of an averaged projection method related to in [1, Corollary 12], where the authors use the Kurdyka-Łojasiewicz inequality. The final section 7 gives limiting examples.

After the initial version of this article appeared, a concept related to our notion of 0-separability, called intrinsic transversality, was announced in [12]. We compare this to our own transversality and regularity concepts in sections 3 and 4.

2. PREPARATION

Given a nonempty closed subset A of \mathbb{R}^n , the projection onto A is the set-valued mapping P_A associating with $x \in \mathbb{R}^n$ the nonempty set

$$P_A(x) = \{a \in A : \|x - a\| = d_A(x)\},$$

where $\|\cdot\|$ is the Euclidean norm, induced by the scalar product $\langle \cdot, \cdot \rangle$, and where $d_A(x) = \min\{\|x - a\| : a \in A\}$. The closed Euclidean ball with center x and radius r is denoted $\mathcal{B}(x, r)$. We write $a \in P_A(b)$ if the projection is potentially set-valued, while $a = P_A(b)$ means it is unique.

A sequence of alternating projections between nonempty closed sets A, B satisfies $b_k \in P_B(a_k), a_{k+1} \in P_A(b_k), k \in \mathbb{N}$. We occasionally switch to the following index-free notation, which is standard in optimization:

$$b \in P_B(a), a^+ \in P_A(b), b^+ \in P_B(a^+), \text{ etc.}$$

The sequence of alternating projections is then $\dots, a, b, a^+, b^+, a^{++}, b^{++}, \dots$. We refer to $a \rightarrow b \rightarrow a^+$, respectively $b \rightarrow a^+ \rightarrow b^+$, as the building blocks of the sequence, where it is always understood that $b \in P_B(a), a^+ \in P_A(b), b^+ \in P_B(a^+)$, etc.

Notions from nonsmooth analysis are covered by [22, 25]. The proximal normal cone to A at $a \in A$ is the set $N_A^p(a) = \{\lambda u : \lambda \geq 0, a \in P_A(a+u)\}$. The normal cone to A at $a \in A$ is the set $N_A(a)$ of vectors v for which there exist $a_k \in A$ with $a_k \rightarrow a$ and $v_k \in N_A^p(a_k)$ such that $v_k \rightarrow v$. The Fréchet normal cone $\widehat{N}_A(a)$ to A at $a \in A$ is the set of v for which $\limsup_{A \ni a' \rightarrow a} \frac{\langle v, a' - a \rangle}{\|a' - a\|} \leq 0$; cf. [22, (1.2)]. We have the inclusions $N_A^p(a) \subset \widehat{N}_A(a) \subset N_A(a)$; cf. [22, Chapter 2.D and (1.6)] or [4, Lemma 2.4]. For any function $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$, the epigraph of f is the set $\text{epi} f = \{(x, \xi) \in \mathbb{R}^n \times \mathbb{R} : \xi \geq f(x)\}$. The proximal subdifferential $\partial_p f(x)$ of a lower semi-continuous function f at $x \in \text{dom} f$ is the set of vectors $v \in \mathbb{R}^n$ such that $(v, -1) \in N_{\text{epi} f}^p(x, f(x))$; [22, (2.81)]. The subdifferential $\partial f(x)$ of f at $x \in \text{dom} f$

is the set of v satisfying $(v, -1) \in N_{\text{epif}}(x, f(x))$. The Fréchet subdifferential $\widehat{\partial}f(x)$ at $x \in \text{dom}f$ is the set of $v \in \mathbb{R}^n$ such that $(v, -1) \in \widehat{N}_{\text{epif}}(x, f(x))$, cf. [22, (1.51)].

3. TANGENTIAL AND NON-TANGENTIAL INTERSECTION

In this section we introduce the fundamental concept of separable intersection of sets A, B , which plays the central role in our convergence theory.

Definition 1. (Separable intersection). *We say that B intersects A separably at $x^* \in A \cap B$ with exponent $\omega \in [0, 2)$ and constant $\gamma > 0$ if there exists a neighborhood U of x^* such that for every building block $b \rightarrow a^+ \rightarrow b^+$ in U , the condition*

$$(1) \quad \langle b - a^+, b^+ - a^+ \rangle \leq (1 - \gamma \|b^+ - a^+\|^\omega) \|b - a^+\| \|b^+ - a^+\|$$

is satisfied. □

We say that B intersects A separably at x^* if (1) holds for *some* $\omega \in [0, 2)$, $\gamma > 0$. If it is also true that A intersects B separably, that is, if the analogue of (1) holds for building blocks $a \rightarrow b \rightarrow a^+$, then we obtain a symmetric condition, and in that case we say that A, B intersect separably at x^* .

Remark 1. Condition (1) discloses itself if we introduce the angle $\alpha = \angle(b - a^+, b^+ - a^+)$ and rewrite (1) in the more suggestive form

$$(1') \quad \frac{1 - \cos \alpha}{\|a^+ - b^+\|^\omega} \geq \gamma,$$

calling this the *angle condition* for the building block $b \rightarrow a^+ \rightarrow b^+$. For $\omega \in (0, 2)$ the interpretation of (1), or (1'), is that if the angle α between $b - a^+$ and $b^+ - a^+$ for two consecutive projection steps $b \rightarrow a^+ \rightarrow b^+$ shrinks down to 0 as the alternating sequence approaches x^* , then α should not shrink too fast. Namely, through (1'), the angle is linked to the shrinking distance between the sets. For $\omega = 0$ the meaning of (1') is that the angle α stays away from 0.

Remark 2. Suppose B intersects A separably with exponent $\omega \in [0, 2)$ and constant $\gamma > 0$ at x^* . Let $\omega' \in (\omega, 2)$ and $\gamma' \in (0, \gamma]$. Then B intersects A also ω' -separably with constant γ' . In consequence, 0-separability is the severest condition, while ω -separability gets less restrictive as ω increases.

As we shall see, for $\omega \geq 2$ property (1) can still be formulated, but turns out too weak to be meaningful. For an illustration see example 7.8.

Remark 3. Informally, when the angle $\alpha = \angle(b - a^+, b^+ - a^+)$ between two consecutive projection steps shrinks to zero, A, B must in some sense intersect *tangentially* at x^* . In contrast, when α stays away from 0, the case of 0-separability, one could say that A, B intersect transversally, or *at an angle*. In that case alternating projections are expected to behave well and converge linearly. Tangential intersection is the more embarrassing case, where convergence could be slowed down or even fail. Our concept of ω -separability gives new insight into the case of tangential intersection.

There has been considerable effort in the literature to *avoid* tangential intersection by making transversality assumptions. We mention transversal intersection in [20], the generalized non-separation property in [22], linearly regular intersection in [21], or the notion of constraint qualification in [4]. In the following we relate these notions to 0-separability.

Bauschke *et al.* [4, Definition 2.1] introduce an extension of the Mordukhovich normal cone called the B -restricted normal cone $N_A^B(x^*)$ to A at $x^* \in A$. They define $u \in N_A^B(x^*)$ if there exist $a_n \in A$, $a_n \rightarrow x^*$, and $u_n \rightarrow u$ such that

$$u_n = \lambda_n (b_n - a_n)$$

for some $\lambda_n > 0$ and $b_n \in B$ with $a_n \in P_A(b_n)$. They then establish basic inclusions between the restricted normal cone and various classical cones [4, Lemma 2.4]. In particular for any $a \in A$ and B one has $N_A^B(a) \subset N_A(a)$.

Now let \tilde{A} and \tilde{B} be non-empty subsets of \mathbb{R}^n . In [4, Definition 6.6] the authors say that $(A, \tilde{A}, B, \tilde{B})$ satisfies the CQ-condition at $x^* \in A \cap B$ if

$$(2) \quad N_A^{\tilde{B}}(x^*) \cap \left(-N_B^{\tilde{A}}(x^*) \right) \subset \{0\}.$$

This condition is to be understood as a transversality hypothesis, because we have the following

Proposition 1. (CQ implies 0-separability). *Let $P_A(\partial B \setminus A) \subset \tilde{A}$, $P_B(\partial A \setminus B) \subset \tilde{B}$, and suppose $(A, \tilde{A}, B, \tilde{B})$ satisfies the CQ-condition at $x^* \in A \cap B$. Then A, B intersect 0-separably at x^* .*

Proof: According to [4, Definition 2.1] the \tilde{B} -restricted proximal normal cone $\hat{N}_A^{\tilde{B}}(a)$ of A at $a \in A$ is the set of vectors u of the form $u = \lambda(\tilde{b} - a)$ for some $\lambda > 0$ and some $\tilde{b} \in \tilde{B}$ satisfying $a \in P_A(\tilde{b})$. The cone $\hat{N}_B^{\tilde{A}}(b)$ at $b \in B$ is defined analogously. Then by [4, Definition 6.1], specialized to the case of two sets, the CQ-number at x^* associated with $(A, \tilde{A}, B, \tilde{B})$ is

$$\theta_\delta(A, \tilde{A}, B, \tilde{B}) = \sup \left\{ \langle u, v \rangle : u \in \hat{N}_A^{\tilde{B}}(a), -v \in \hat{N}_B^{\tilde{A}}(b), \|u\| \leq 1, \|v\| \leq 1, a, b \in \mathcal{B}(x^*, \delta) \right\}$$

and the limiting CQ-number is

$$\theta(A, \tilde{A}, B, \tilde{B}) = \lim_{\delta \rightarrow 0^+} \theta_\delta(A, \tilde{A}, B, \tilde{B}).$$

The authors show in [4, Theorem 6.8] that for two sets the CQ-condition $N_A^{\tilde{B}}(x^*) \cap (-N_B^{\tilde{A}}(x^*)) \subset \{0\}$, implies $\theta(A, \tilde{A}, B, \tilde{B}) \leq 1$.

Using this, pick $\delta > 0$ such that $\theta_\delta(A, \tilde{A}, B, \tilde{B}) =: 1 - \gamma < 1$. Consider a building block $b \rightarrow a^+ \rightarrow b^+$ as in definition 1 with $b, a^+, b^+ \in U := \mathcal{B}(x^*, \delta)$. Then we have $b \in \tilde{B}$ and $a^+ \in \tilde{A}$. Hence $b - a^+ \in \hat{N}_A^{\tilde{B}}(a^+)$ and $a^+ - b^+ \in \hat{N}_B^{\tilde{A}}(b^+)$, and also

$$u = (b - a^+)/\|b - a^+\| \in \hat{N}_A^{\tilde{B}}(a^+) \text{ and } v = (b^+ - a^+)/\|b^+ - a^+\| \in -\hat{N}_B^{\tilde{A}}(b^+).$$

Therefore, if $\alpha = \angle(b - a^+, b^+ - a^+)$, then $\cos \alpha = \langle u, v \rangle \leq \theta_\delta(A, \tilde{A}, B, \tilde{B}) = 1 - \gamma$ by the definition of θ_δ , because $b, a^+, b^+ \in \mathcal{B}(x^*, \delta)$ and $\|u\| = \|v\| = 1$. That shows $1 - \cos \alpha \geq \gamma > 0$ and proves that B intersects A 0-separably at x^* with constant γ . The estimate for building blocks $a \rightarrow b \rightarrow a^+$ is analogous. \square

Example 7.3 shows that the converse of proposition 1 is not true. In fact, 0-separability seems more versatile in applications, while still guaranteeing linear convergence. We conclude by noting that linearly regular intersection in the sense of [21], and transversality in the sense of [20], imply 0-separability.

Following [21, section 2, (2.2)], A and B have linearly regular intersection at $x^* \in A \cap B$ if

$$(3) \quad N_A(x^*) \cap (-N_B(x^*)) = \{0\}.$$

This property is called *strong regularity* in [18] and *the basic qualification condition for sets* in [22, Definition 3.2 (i)]. As a consequence of $N_A^{\tilde{B}}(a) \subset N_A(a)$, $N_B^{\tilde{A}}(b) \subset N_B(b)$, linearly regular intersection implies that $(A, \tilde{A}, B, \tilde{B})$ satisfies the CQ-condition at x^* for any nonempty \tilde{A} and \tilde{B} in \mathbb{R}^n ; cf. [5]. By Proposition 1 we therefore have:

Corollary 1. (Linear regularity implies 0-separability). *Suppose A, B intersect linearly regularly at $x^* \in A \cap B$. Then they intersect 0-separably at x^* .* \square

As we mentioned before, in the context of alternating projections linear regularity and the CQ-condition are to be understood as transversality type hypotheses, indicating that the sets A, B intersect *at an angle at x^** , as opposed to intersecting tangentially. This is confirmed by relating 0-separability to the classical notion of transversality. Following [20, def. 3], two C^2 -manifolds A, B in \mathbb{R}^n intersect transversally at $x^* \in A \cap B$ if

$$(4) \quad T_A(x^*) + T_B(x^*) = \mathbb{R}^n,$$

where $T_M(x^*)$ is the tangent space to M at $x^* \in M$. We then have the following

Corollary 2. *Let A, B be C^2 -manifolds which intersect transversally at x^* . Then A and B intersect 0-separably at x^* .* \square

Proof: Indeed, as shown in [20, Theorem 18], classical transversality (4) implies linear regular intersection (3), hence we can apply Corollary 1. \square

After the initial version of this work was published, a related concept termed *intrinsic transversality* was proposed in [12]. Following [12, Def. 2.2], A, B are *intrinsically transversal* at $x^* \in A \cap B$ with constant $\kappa \in (0, 1]$ if there exists a neighborhood U of x^* such that for every $a^+ \in A \cap U \setminus B$ and every $b^+ \in B \cap U \setminus A$ the estimate

$$(5) \quad \max \left\{ d \left(\frac{a^+ - b^+}{\|a^+ - b^+\|}, N_B^p(b^+) \right), d \left(\frac{a^+ - b^+}{\|a^+ - b^+\|}, -N_A^p(a^+) \right) \right\} \geq \kappa > 0$$

is satisfied. This relates to 0-separability as follows.

Proposition 2. (Intrinsic transversality implies 0-separability). *Suppose A, B are intrinsically transversal at $x^* \in A \cap B$ with transversality constant $\kappa \in (0, 1]$. Then they intersect 0-separably at x^* with constant $\gamma = \kappa^2/2$.*

Proof: By assumption there exists a neighborhood U of x^* on which (5) is satisfied. Now let $b^+ \in P_B(a^+) \cap U$, $a^+ \neq b^+$, $a^+ \in A \cap U$. Then since $a^+ - b^+ \in N_B^p(b^+)$, we have $d \left(\frac{a^+ - b^+}{\|a^+ - b^+\|}, N_B^p(b^+) \right) = 0$, so by (5) we must have $d \left(\frac{a^+ - b^+}{\|a^+ - b^+\|}, -N_A^p(a^+) \right) \geq \kappa$. Since $b - a^+ \in N_A^p(a^+)$, we obtain

$$2 - 2 \frac{\langle b^+ - a^+, b - a^+ \rangle}{\|a^+ - b^+\| \|b - a^+\|} \geq \kappa^2,$$

and this readily gives

$$\langle b^+ - a^+, b - a^+ \rangle \leq \left(1 - \frac{\kappa^2}{2} \right) \|a^+ - b^+\| \|b - a^+\|,$$

which is (1) for the case $\omega = 0$ and $\gamma = \kappa^2/2$, as claimed. \square

We will resume the discussion of separable intersection of sets in section 6.

4. HÖLDER REGULARITY

In this section we introduce the concept of Hölder regularity. We then relate it to other regularity notions like Clarke regularity, prox-regularity, superregularity in the sense of [21], and its extension in [4].

Definition 2. (Hölder regularity). *Let $\sigma \in [0, 1)$. The set B is σ -Hölder regular with respect to A at $b^* \in A \cap B$ if there exists a neighborhood U of b^* and a constant $c > 0$ such that for every $a^+ \in A \cap U$, and every $b^+ \in P_B(a^+) \cap U$ one has*

$$(6) \quad \mathcal{B}(a^+, (1+c)r) \cap \{b \in P_A(a^+)^{-1} : \langle a^+ - b^+, b - b^+ \rangle > \sqrt{c}r^{\sigma+1} \|b - b^+\|\} \cap B = \emptyset,$$

where $r = \|a^+ - b^+\|$. We say that B is Hölder regular with respect to A if it is σ -Hölder regular with respect to A for every $\sigma \in [0, 1)$. \square

Remark 4. Using the angle $\beta = \angle(a^+ - b^+, b - b^+)$ and $r = \|a^+ - b^+\|$, condition (6) can be re-written in the following more suggestive form

$$(6') \quad \mathcal{B}(a^+, (1+c)r) \cap \{b \in P_A(a^+)^{-1} : \cos \beta > \sqrt{c}r^\sigma\} \cap B = \emptyset.$$

Geometrically, this means that the right circular cone with axis $a^+ - b^+$ and aperture $\beta = \arccos \sqrt{c}r^\sigma$ truncated by the ball $\mathcal{B}(a^+, (1+c)r)$ and the B -restricted proximal normal cone $\widehat{N}_A^B(a^+)$ contains no points of B other than b^+ .

In the remainder of this section we relate Hölder regularity to older geometric and analytic regularity concepts. We first consider notions related to 0-Hölder regularity. The case of σ -Hölder regularity with $\sigma > 0$ will be considered later.

Definition 3. (Superregularity [21, Proposition 4.4]). *A closed set B in \mathbb{R}^n is called superregular at $b^* \in B$ if for every $\epsilon > 0$ there exists $\delta > 0$ such that for all $b, b^+ \in \mathcal{B}(b^*, \delta) \cap B$ and $u \in N_B^p(b^+)$, the estimate $\langle u, b - b^+ \rangle \leq \epsilon \|u\| \|b - b^+\|$ is satisfied.*

Definition 4. ((A, ϵ, δ) -regularity [4, Definition 8.1, (i)]). *Let A, B be closed sets in \mathbb{R}^n . B is called (A, ϵ, δ) -regular at $b^* \in A \cap B$ if for all $b, b^+ \in \mathcal{B}(b^*, \delta) \cap B$ and every $u \in \widehat{N}_B^A(b^+)$, the estimate $\langle u, b - b^+ \rangle \leq \epsilon \|u\| \|b - b^+\|$ is satisfied.*

The two concepts are linked as follows: B is superregular at $b^* \in B$ if and only if for every $\epsilon > 0$ there exists $\delta > 0$ such that B is $(\mathbb{R}^n, \epsilon, \delta)$ -regular at b^* in the sense of Definition 4, see [4, Definition 8.1].

Proposition 3. (0-Hölder regularity from superregularity). *Suppose B is (A, ϵ, δ) -regular at $b^* \in A \cap B$. Then B is 0-Hölder regular at b^* with respect to A with constant $c = \epsilon^2$. In particular, if B is superregular at b^* , then B is 0-Hölder regular with respect to A with constant c that may be chosen arbitrarily small.*

Proof: Since superregularity of B at b^* implies that for every $\epsilon > 0$ there exists $\delta > 0$ such that B is (A, ϵ, δ) -regular at b^* , [4], it remains to prove the first part of the statement.

In order to check 0-Hölder regularity, we have to provide a neighborhood U of b^* and $c > 0$ such that (6) is satisfied with $\sigma = 0$. We choose $U = \mathcal{B}(b^*, \frac{\delta}{4(1+\epsilon^2)})$ and put $c = \epsilon^2$. To check (6) pick $a^+, b^+ \in U$ such that $b^+ \in P_B(a^+)$, $a^+ \in A$. That gives $r = \|b^+ - a^+\| \leq \frac{\delta}{2(1+c)}$. By the definition of the restricted normal cone we have $u := a^+ - b^+ \in \widehat{N}_B^A(b^+)$. Now let $b \in B$, $b \neq b^+$. We have to show that b is not an element of the set in (6) for $\sigma = 0$. Suppose $b \in \mathcal{B}(a^+, (1+c)r)$. Then we have to show $\langle a^+ - b^+, b - b^+ \rangle \leq \sqrt{c}r \|b - b^+\|$. Observe that $\|b - b^*\| \leq \|b - a^+\| + \|a^+ - b^*\| \leq (1+c)r + \frac{\delta}{4(1+c)} \leq (1+c)\frac{\delta}{2(1+c)} + \frac{\delta}{4(1+c)} < \delta$. Hence (A, ϵ, δ) -regularity implies $\langle u, b - b^+ \rangle \leq \epsilon \|u\| \|b - b^+\| = \sqrt{c} \|u\| \|b - b^+\|$, and the claim follows. \square

Remark 5. Example 7.1 shows that the converse of proposition 3 is not true. The difference between superregularity and its extension (A, ϵ, δ) -regularity on the one hand, and 0-Hölder regularity on the other, is the following: in (6) we exclude points in the intersection of a restricted right circular cone with vertex b^+ , axis $a^+ - b^+$, and aperture $\beta = \arccos \sqrt{cr^\sigma}$ and the shrinking ball $\mathcal{B}(a^+, (1+c)r)$. In contrast, (A, ϵ, δ) -regularity forbids many more points, namely all points in that same cone, but within the fixed ball $\mathcal{B}(b^*, \delta)$. In consequence, this type of regularity is not suited to deal with singularities pointing inwards, like the prototype in example 7.1. \square

Remark 6. If B is σ -Hölder regular at b^* with respect to A with constant $c > 0$ on the neighborhood U of b^* , and if $\sigma' < \sigma$, then for every $c' \in (0, c)$ there exists a neighborhood $V \subset U$ of b^* such that B is σ' -Hölder regular at b^* with constant c' . Indeed, if $b \in \mathcal{B}(a^+, (1+c')r)$ in (6), then also $b \in \mathcal{B}(a^+, (1+c)r)$, hence by assumption $\cos \beta \leq \sqrt{cr^\sigma} = \sqrt{cr^{\sigma-\sigma'}r^{\sigma'}} \leq \sqrt{c'r^{\sigma'}} < \sqrt{c'}$ if V is chosen so that $\sqrt{cr^{\sigma-\sigma'}} < \sqrt{c'}$. \square

We next justify our notion of Hölder regularity by proving that prox-regular sets are σ -Hölder regular for every $\sigma \in (0, 1]$. Recall that a set B in \mathbb{R}^n is *prox-regular* at $b^* \in B$ if there exists a neighborhood U of b^* such that $P_B(y)$ is single-valued for every $y \in U$, cf. [25, Chapter 13].

Consider $b \in B$ and let $d \in N_B^p(b)$ be a unit proximal normal to B at b . Define the reach of B at b along d as

$$(7) \quad R(b, d) = \sup\{R \geq 0 : b = P_B(b + td) \text{ for every } 0 \leq t \leq R\}.$$

Then $R(b, d) \in (0, \infty]$, and the case $R(b, d) = \infty$ occurs e.g. if B is convex and b a boundary point of B . We can say that $\mathcal{B}(b + R(b, d)d, R(b, d))$ is the largest ball with its centre on $b + \mathbb{R}_+d$ which touches B in b from outside, i.e., has no points from B in its interior.

It was shown in [24, Thm. 1.3 (h)] that B is prox-regular at $b^* \in B$ if and only if there exists $r > 0$ and a neighborhood U of b^* such that $R(b, d) \geq r$ for every $b \in U \cap B$ and every $d \in N_B^p(b)$ with $\|d\| = 1$. An immediate consequence is that sets of positive reach in the sense of Federer [15] are prox-regular; see e.g. [24, Theorem 1.3]. Therefore, prox-regularity is a local version of positive, or non-vanishing, reach.

We now relax the concept of non-vanishing reach to sets where the reach may vanish at some boundary points, but slowly so.

Definition 5. Let $\sigma \in (0, 1]$. The set B has σ -slowly vanishing reach with respect to the set A at $b^* \in A \cap B$ if there exists $0 \leq \tau < 1$ such that

$$(8) \quad \limsup_{A \ni a \rightarrow b^*, b \in P_B(a)} \frac{\|a - b\|^\sigma}{R(b, d)} \leq \tau,$$

where $d = (a - b)/\|a - b\|$. We say that the reach vanishes with exponent σ and rate τ . \square

Proposition 4. If B is prox-regular at $b^* \in A \cap B$, then it has slowly vanishing reach at b^* with respect to A with rate $\tau = 0$ and arbitrary exponent $\sigma \in (0, 1]$.

Proof: Let $\tau' > 0$. By [24, Thm. 1.3 (h)] prox-regularity at b^* implies that there exist $\epsilon > 0$ and $r > 0$ such that $R(b, d) \geq r$ for every $b \in B$ with $\|b - b^*\| \leq \epsilon$ and every $d \in N_B^p(b)$ with $\|d\| = 1$. By shrinking ϵ if necessary, we may assume $\epsilon^\sigma/(2^\sigma r) < \tau'$. Now let $a \in A \cap \mathcal{B}(b^*, \frac{\epsilon}{2})$ be arbitrary, choose $b \in P_B(a)$, and let $d = (a - b)/\|a - b\|$. Then as $b^* \in A \cap B$, we have $\|b - b^*\| \leq \|b - a\| + \|a - b^*\| \leq 2\|a - b^*\| \leq \epsilon$. Since $d \in N_B^p(b)$, the above gives us $R(b, d) \geq r$. Therefore, since $\|a - b\| = d_B(a) \leq \|a - b^*\|$, the quotient in (8) satisfies

$$\frac{\|a - b\|^\sigma}{R(b, d)} \leq \frac{\epsilon^\sigma}{2^\sigma r} < \tau',$$

and since $\tau' > 0$ was arbitrary, this shows that (8) is satisfied with $\tau = 0$. \square

Proposition 5. (Hölder regularity from slowly vanishing reach). *Let $\sigma \in (0, 1)$. Suppose B has σ -slowly vanishing reach with rate $\tau \in [0, 1)$ with respect to A at $b^* \in A \cap B$. Then B is $(1 - \sigma)$ -Hölder regular with respect to A with any constant $c > 0$ satisfying*

$$(9) \quad \frac{\tau}{2} \sqrt{2 + c} < 1.$$

In particular, c may be chosen arbitrarily small.

Proof: 1) We have to show that there exists a neighborhood U of b^* such that (6) is satisfied with c as in (9) and with exponent $1 - \sigma$.

By condition (9) we can choose $\tau' > \tau$ and $\epsilon > 0$ such that

$$\frac{\tau'}{2} \left(\epsilon + \sqrt{\epsilon^2 + 2 + c} \right) < 1.$$

By condition (8), and since $\tau < \tau'$, there exists a neighborhood U of b^* such that whenever $a^+, b^+ \in U$, $a^+ \in A$, $b^+ \in P_B(a^+)$ and $d = (a^+ - b^+)/\|a^+ - b^+\|$, then $r^\sigma/R(b^+, d) < \tau'$, where $r := \|a^+ - b^+\|$. On shrinking U further if necessary, we may arrange that $a^+, b^+ \in U$ implies $r^{1-\sigma} = \|a^+ - b^+\|^{1-\sigma} < \epsilon$. We will show that U is the neighborhood we need in condition (6).

2) To prove this pick $a^+, b^+ \in U$, $a^+ \in A$, $b^+ \in P_B(a^+)$, put $r = \|a^+ - b^+\|$, and let $b \in B$, $b \neq b^+$. We have to show that b is not an element of the set (6'). To check this, let β be the angle $\beta = \angle(a^+ - b^+, b - b^+)$. Since there is nothing to prove for $b \notin \mathcal{B}(a^+, (1+c)r)$, we assume $b \in \mathcal{B}(a^+, (1+c)r)$. Now we have to show that $\cos \beta \leq \sqrt{cr^{1-\sigma}}$. As this is clear for $\cos \beta \leq 0$, we may assume $\cos \beta > 0$.

Let us define

$$(10) \quad R := \frac{r}{2} \left(1 + \sqrt{1 + \frac{2c + c^2}{\cos^2 \beta}} \right),$$

where r, β are as before. We claim that the ball $\mathcal{B}(b^+ + Rd, R)$ contains b , where as above $d = (a^+ - b^+)/\|a^+ - b^+\|$. To prove this, note that by the cosine theorem, applied in the triangle a^+, b^+, b , we have

$$\|a^+ - b\|^2 = r^2 + \|b - b^+\|^2 - 2r\|b - b^+\| \cos \beta.$$

Since $\|a^+ - b\| \leq (1+c)r$, we obtain

$$r^2 + \|b - b^+\|^2 - 2r\|b - b^+\| \cos \beta \leq (1+c)^2 r^2,$$

which on completing squares turns out to be the same as

$$\|b - b^+\| \leq r \left(\cos \beta + \sqrt{2c + c^2 + \cos^2 \beta} \right) = 2R \cos \beta.$$

Here the last equality uses the definition (10) of R . We therefore obtain

$$\|b - b^+\|^2 \leq 2R \cos \beta \|b - b^+\|,$$

and using the cosine theorem again, now in the triangle $b^+ + Rd, b^+, b$, we deduce

$$\|b^+ + Rd - b\|^2 = R^2 + \|b - b^+\|^2 - 2R\|b - b^+\| \cos \beta \leq R^2.$$

This gives $b \in \mathcal{B}(b^+ + Rd, R)$ as claimed.

3) By the definition (7) of $R(b^+, d)$, any radius $R' < R(b^+, d)$ must give rise to a ball with $\mathcal{B}(b^+ + R'd, R') \cap B = \{b^+\}$. But as we have shown in part 2), the ball $\mathcal{B}(b^+ + Rd, R)$ contains b , so necessarily $R \geq R(b^+, d)$. Hence by the choice of U in part 1), $r^\sigma/R \leq$

$r^\sigma/R(b^+, d) < \tau'$, or what is the same, $r^\sigma < R\tau'$. Substituting the definition (10) of R and multiplying by $r^{-\sigma}$, we deduce

$$1 < r^{1-\sigma}\tau' \left(\frac{1}{2} + \frac{1}{2} \sqrt{1 + \frac{2c + c^2}{\cos^2 \beta}} \right).$$

Now suppose that $\cos \beta > \sqrt{cr^{1-\sigma}}$, contrary to what we wish to show. Then

$$\begin{aligned} 1 &< r^{1-\sigma}\tau' \left(\frac{1}{2} + \frac{1}{2} \sqrt{1 + \frac{2c + c^2}{cr^{2(1-\sigma)}}} \right) \\ &= \frac{\tau'}{2} \left(r^{1-\sigma} + \sqrt{r^{2(1-\sigma)} + 2 + c} \right) \\ &< \frac{\tau'}{2} \left(\epsilon + \sqrt{\epsilon^2 + 2 + c} \right) < 1, \end{aligned}$$

a contradiction. That proves the result. \square

Since prox-regularity at $b^* \in B$ implies slowly vanishing reach at b^* with respect to any closed set A containing b^* , we have the following immediate consequence.

Corollary 3. (Hölder regularity from prox-regularity). *Let B be prox-regular. Then B is σ -Hölder regular for every $\sigma \in [0, 1)$ with a constant $c > 0$ that may be chosen arbitrarily small.*

Proof: For $\sigma = 0$ this follows from Proposition 3, because prox-regularity implies superregularity. For $\sigma \in (0, 1)$ we obtain it by combining Propositions 4 and 5. \square

Consider the case of a Lipschitz domain B . Here Hölder regularity may be related to a property of the boundary ∂B of B .

Proposition 6. *Let $\sigma \in (0, 1)$. Suppose B is the epigraph of a locally Lipschitz function $f : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$. Let $x^* \in \mathbb{R}^{n-1}$ and suppose there exists a neighborhood V of x^* and $\mu > 0$ such that for every $x_0 \in V$ and every proximal subgradient $g \in \partial_p f(x_0)$ the one-sided Hölder estimate $f(x_0) + \langle g, x - x_0 \rangle - \mu \|x - x_0\|^{1+\sigma} \leq f(x)$ is satisfied for all $x \in V$. Then B is σ -Hölder regular at $(x^*, f(x^*)) \in B$ with respect to every closed set A containing $(x^*, f(x^*))$, and for every constant $c > 0$ satisfying $\mu \leq \sqrt{c}/(2+c)^\sigma$.*

Proof: We have to find a neighborhood U of $b^* = (x^*, f(x^*)) \in B$ such that (6) is satisfied with exponent σ and constant c satisfying $\mu \leq \sqrt{c}/(2+c)^\sigma$. Choose $\epsilon > 0$ such that $\mathcal{B}(x^*, \epsilon) \subset V$. Now choose $\delta > 0$ with $\delta < \epsilon/(2+c)$ and define $U = \mathcal{B}(b^*, \delta)$. We will show that U is as required.

In order to check (6), choose $a^+ \in A \setminus B$ and $b^+ \in P_B(a^+)$ such that $a^+, b^+ \in U$. As $b^+ \in P_B(a^+) \cap U$, we get $b^+ = (x_0, f(x_0)) \in B$ for some $x_0 \in V \subset \mathbb{R}^{n-1}$, while $a^+ \notin B = \text{epi} f$ implies $a^+ = (x_1, \xi_1)$ for some $\xi_1 < f(x_1)$. Since $a^+ - b^+$ is a proximal normal to B at b^+ , there exists a proximal subgradient $g \in \partial_p f(x_0)$ such that $(x_1, \xi_1) = (x_0, f(x_0)) + t(g, -1)$ for some $t > 0$. Using $\|a^+ - b^+\| = r$, we can therefore write

$$a^+ - b^+ = \left(\frac{rg}{\sqrt{1 + \|g\|^2}}, -\frac{r}{\sqrt{1 + \|g\|^2}} \right).$$

Now consider $b \in B$, $b \neq b^+$ such that $\|a^+ - b\| \leq (1+c)r$. To verify (6') we have to show that $\cos \beta \leq \sqrt{c}r^\sigma$, where $\beta = \angle(a^+ - b^+, b - b^+)$. Since there is nothing to prove

for $\cos \beta \leq 0$, we assume $\cos \beta > 0$. By the definition of B we have $b = (x, \xi)$ for some $x \in \mathbb{R}^{n-1}$ and $\xi \geq f(x)$. Now

$$\begin{aligned} \cos \beta &= \frac{\langle g, x - x_0 \rangle - \xi + f(x_0)}{\sqrt{1 + \|g\|^2} \sqrt{\|x - x_0\|^2 + (\xi - f(x_0))^2}} \\ &\leq \frac{\langle g, x - x_0 \rangle - f(x) + f(x_0)}{\|x - x_0\|} \\ &\leq \frac{\mu \|x - x_0\|^{1+\sigma}}{\|x - x_0\|} = \mu \|x - x_0\|^\sigma \leq \sqrt{cr}^\sigma. \end{aligned}$$

Here the first inequality uses the fact that $\xi \geq f(x)$. The second inequality uses the one-sided Hölder estimate from the hypothesis. In order to be allowed to use this estimate, we have to assure that $x \in V$. This follows from

$$\begin{aligned} \|x - x^*\| &\leq \|b - b^*\| \leq \|b - a^+\| + \|a^+ - b^*\| \\ &\leq (1 + c)\|a^+ - b^+\| + \|a^+ - b^*\| \\ &\leq (2 + c)\|a^+ - b^*\| \leq (2 + c)\delta < \epsilon. \end{aligned}$$

The third inequality can be seen as follows. We have

$$\begin{aligned} \|x - x_0\| &\leq \|b - b^+\| \leq \|b - a^+\| + \|a^+ - b^+\| \\ &\leq (2 + c)\|a^+ - b^+\| = (2 + c)r. \end{aligned}$$

Hence

$$\mu \|x - x_0\|^\sigma \leq \mu(2 + c)^\sigma r^\sigma \leq \sqrt{cr}^\sigma$$

by the choice of c . That completes the argument. \square

Remark 7. The nomenclature in Proposition 6 can be explained as follows. Lipschitz smoothness [14] of $-f$ at x_0 is a well-known second-order property equivalent to the second difference quotient

$$\Delta_2(x) = \frac{f(x) - f(x_0) - \langle g, x - x_0 \rangle}{\|x - x_0\|^2} \geq -\mu > -\infty$$

being bounded below for $g \in \partial f(x_0)$ and x in a neighborhood of x_0 . The Hölder estimate in Proposition 6 is the analogous but weaker condition $\Delta_{1+\sigma}(\cdot) \geq -\mu > -\infty$ for some $\sigma \in (0, 1)$. In analogy with [14] one could call this σ -Hölder smoothness of $-f$ at x_0 .

We consider the following natural modification of amenability from [25]. The set $B \subset \mathbb{R}^n$ is called σ -Hölder amenable at $x^* \in B$ if there exists a neighborhood U of x^* , a class $C^{1,\sigma}$ -mapping $G : \mathbb{R}^n \rightarrow \mathbb{R}^m$, and a closed convex set $C \subset \mathbb{R}^m$, such that $B \cap U = \{x \in U : G(x) \in C\}$ and $N_C(G(x^*)) \cap \ker(DG(x^*)^T) = \{0\}$, where $DG(x^*)$ denotes the first-order differential of G at x^* . A typical example in optimization is when B is defined by $C^{1,\sigma}$ equality and inequality constraints, where the Mangasarian-Fromowitz constraint qualification holds at x^* , see [24, Prop. 2.3], [25, Chap 10, F.], [21, Prop. 4.8].

Proposition 7. (Hölder regularity from Hölder amenability). *Suppose the closed set B is σ' -Hölder amenable at x^* for some $\sigma' \in (0, 1]$. Then B is σ -Hölder regular at x^* with respect to any closed set A containing x^* for every $\sigma \in (0, \sigma')$, and with arbitrary constant c . \square*

The proof may be adopted from on [21, Prop. 4.8] with minor changes, and we skip the details. This result suggests that Hölder regularity is settled between the weaker superregularity and the stronger prox-regularity. This is true as long as we consider this type of regularity as a property of B alone. We stress, however, that it is the combination with A and the shrinking distance between the sets in (6) which makes our definition 3 truly versatile in applications. This is corroborated by the following observation.

Proposition 8. (Hölder regularity from intrinsic transversality). *Suppose A, B are intrinsically transversal at x^* with constant of transversality $\kappa \in (0, 1]$. Then B is σ -Hölder regular at x^* with respect to A for every $\sigma \in [0, 1)$, with any constant $c > 0$ satisfying $c < \frac{\kappa^2}{1-\kappa^2}$.*

Proof: 1) By [12, Def. 2.2] and [12, Prop. 6.9] there exists a neighborhood V of x^* such that $\max\{\text{dist}(u, N_B(b)), \text{dist}(u, -N_A(a^+))\} \geq \kappa > 0$ for all $a^+ \in A \cap V \setminus B, b \in B \cap V \setminus A$, and $u = (a^+ - b)/\|a^+ - b\|$. Following entirely the argument in [12, page 6], one can now find a smaller neighborhood U of x^* such that the following is true: If $b \in B \cap U$ and $a^+ \in P_A(b) \cap U$, then even $d_B(a^+) \leq (1 - \kappa^2)\|b - a^+\|$.

2) We claim that U is the neighborhood required in σ -Hölder regularity with constant c . To check this, we have to show that the set (6) is empty. We assume that $b \in U$ is an element of that set. Then $b \in P_A(a^+)^{-1} \cap B$ and $b \in \mathcal{B}(a^+, (1+c)r)$. Hence we are in the situation of part 1), which means $r = d_B(a^+) \leq (1 - \kappa^2)\|b - a^+\| \leq (1 - \kappa^2)(1+c)r < r$, a contradiction. Hence the set (6) is empty. \square

5. CONVERGENCE

In this section we prove the main convergence result. Alternating projections converge locally for sets which intersect separably, if one of the sets is Hölder regular with respect to the other. The proof requires the following preparatory lemma.

Lemma 1. (Three-point estimate). *Suppose B intersects A separably at $x^* \in A \cap B$ with exponent $\omega \in [0, 2)$ and constant $\gamma > 0$ on the neighborhood U of x^* . Suppose B is also $\omega/2$ -Hölder regular at $x^* \in A \cap B$ with respect to A on U with constant $c > 0$ satisfying $c < \frac{\gamma}{2}$. Then there exists $0 < \ell < 1$, depending only on γ, c and U , such that*

$$(11) \quad \|a^+ - b^+\|^2 + \ell\|b - b^+\|^2 \leq \|a^+ - b\|^2$$

for every building block $b \rightarrow a^+ \rightarrow b^+$ in U .

Proof: 1) By the cosine theorem we have

$$\|a^+ - b\|^2 = \|a^+ - b^+\|^2 + \|b - b^+\|^2 - 2\|a^+ - b^+\|\|b - b^+\| \cos \beta,$$

where $\beta = \angle(b - b^+, a^+ - b^+)$. Hence in order to assure (11) we have to find $\ell \in (0, 1)$ such that

$$(12) \quad \frac{1-\ell}{2}\|b - b^+\| \geq \|a^+ - b^+\| \cos \beta$$

for all building blocks $b \rightarrow a^+ \rightarrow b^+$ in U . We consider two cases. Case I is when $\beta \in (\frac{\pi}{2}, \pi]$. Case II is $\beta \in [0, \frac{\pi}{2}]$.

2) We start by discussing case I. For angles $\beta \in (\frac{\pi}{2}, \pi]$ we have $\cos \beta < 0$, hence (12) is trivially true if we choose any $0 < \ell < 1$. For instance $\ell = \frac{1}{2}$ will do. To establish (12) we may now concentrate on case II, where $\beta \in [0, \frac{\pi}{2}]$.

3) In case II we want to use $\omega/2$ -Hölder regularity of B with respect to A . We subdivide case II in two subcases. Case IIa is when $\cos \beta \leq \sqrt{c}\|a^+ - b^+\|^{\frac{\omega}{2}}$. Case IIb is when $\cos \beta > \sqrt{c}\|a^+ - b^+\|^{\frac{\omega}{2}}$.

Let us start with case IIa, where $\cos \beta \leq \sqrt{c} \|a^+ - b^+\|^{\frac{\omega}{2}}$. Observe that

$$\begin{aligned} \|b - b^+\|^2 &= \|b - a^+\|^2 + \|a^+ - b^+\|^2 - 2\langle b - a^+, b^+ - a^+ \rangle \\ &= (\|b - a^+\| - \|a^+ - b^+\|)^2 + 2\|b - a^+\| \|a^+ - b^+\| (1 - \cos \alpha) \\ &\geq 2\|b - a^+\| \|a^+ - b^+\| (1 - \cos \alpha), \end{aligned}$$

where $\alpha = \angle(b - a^+, b^+ - a^+)$. By the angle condition (1') we have $1 - \cos \alpha \geq \gamma \|a^+ - b^+\|^\omega$ for every building block $b \rightarrow a^+ \rightarrow b^+$ in U . Hence

$$\|b - b^+\|^2 \geq 2\gamma \|b - a^+\| \|a^+ - b^+\|^{1+\omega} \geq 2\gamma \|a^+ - b^+\|^{2+\omega},$$

where the last estimate uses $b^+ \in P_B(a^+)$. Altogether we obtain

$$\|b - b^+\| \geq \sqrt{2\gamma} \|a^+ - b^+\|^{1+\frac{\omega}{2}} \geq \sqrt{\frac{2\gamma}{c}} \|a^+ - b^+\| \cos \beta,$$

bearing in mind that we are in case IIa. To assure (12) we put $\ell = 1 - \sqrt{\frac{2c}{\gamma}}$. Then $\ell \in (0, 1)$, because of the hypothesis $c < \frac{\gamma}{2}$.

4) Let us now deal with case IIb, where $\cos \beta > \sqrt{c} \|a^+ - b^+\|^{\frac{\omega}{2}}$. By Hölder regularity (6) of B with respect to A and since $a^+ \in P_A(b)$, we have $b \notin \mathcal{B}(a^+, (1+c)r)$, where $r = \|a^+ - b^+\|$. In other words, $\|b - a^+\| > (1+c)\|a^+ - b^+\|$. Using this and the cosine theorem again, we have

$$\begin{aligned} \|b - b^+\|^2 &= \|b - a^+\|^2 - \|a^+ - b^+\|^2 + 2\|b - b^+\| \|a^+ - b^+\| \cos \beta \\ &\geq c(c+2)\|a^+ - b^+\|^2 + 2\|b - b^+\| \|a^+ - b^+\| \cos \beta. \end{aligned}$$

Since $a^+ \neq b^+$, this may be rearranged as

$$(13) \quad \frac{\|b - b^+\|^2}{\|a^+ - b^+\|^2} - 2 \frac{\|b - b^+\|}{\|a^+ - b^+\|} \cos \beta - c(c+2) \geq 0.$$

Hence (13) implies that the polynomial $P(X) = X^2 - 2X \cos \beta - c(c+2)$ is nonnegative at $X = \frac{\|b - b^+\|}{\|a^+ - b^+\|}$. But for nonnegative X , nonnegativity $P(X) \geq 0$ is equivalent to

$$X \geq \cos \beta + \sqrt{\cos^2 \beta + c(c+2)} = \cos \beta \left(1 + \sqrt{1 + \frac{c(c+2)}{\cos^2 \beta}} \right).$$

Hence for $X = \frac{\|b - b^+\|}{\|a^+ - b^+\|}$ we know that

$$(14) \quad \frac{\|b - b^+\|}{\|a^+ - b^+\|} \geq \cos \beta \left(1 + \sqrt{1 + \frac{c(c+2)}{\cos^2 \beta}} \right).$$

Put $\Theta_{r,\beta} = 1 + \sqrt{1 + \frac{c(c+2)}{\cos^2 \beta}}$, then $\Theta_{r,\beta} \geq c+2$. Hence $\ell = \frac{c}{2+c}$ assures (12) in case IIb.

5) In conclusion, if we put $\ell = \min \left\{ \frac{1}{2}, 1 - \sqrt{\frac{2c}{\gamma}}, \frac{c}{2+c} \right\}$, with $c < \frac{\gamma}{2}$, then (12) is satisfied in all cases. \square

Theorem 1. (Local convergence). *Suppose B intersects A separably at $x^* \in A \cap B$ with exponent $\omega \in [0, 2)$ and constant γ and is $\omega/2$ -Hölder regular at x^* with respect to A and constant $c < \frac{\gamma}{2}$. Then there exists a neighborhood V of x^* such that every sequence of alternating projections between A and B which enters V , converges to a point $b^* \in A \cap B$.*

Proof: 1) By hypothesis there exists a neighborhood $U = \mathcal{B}(x^*, 4\epsilon)$ of $x^* \in A \cap B$ such that every building block $b \rightarrow a^+ \rightarrow b^+$ with $b, a^+, b^+ \in U$ satisfies the angle condition $1 - \cos \alpha \geq \gamma \|b^+ - a^+\|^\omega$, where $\alpha = \angle(b - a^+, b^+ - a^+)$. In addition, by shrinking U if necessary, we may assume that B is $\omega/2$ -Hölder regular at x^* on U with constant $c < \frac{\gamma}{2}$. Then by the three-point estimate (Lemma 1) there exists $\ell \in (0, 1)$ depending only on c, γ and U , such that $\|a^+ - b^+\|^2 + \ell \|b - b^+\|^2 \leq \|a^+ - b\|^2$ for every such building block. Since $\|a^+ - b\| \leq \|a - b\|$, we deduce the following four-point estimate

$$(15) \quad d_B(a^+)^2 + \ell \|b - b^+\|^2 \leq d_B(a)^2$$

for building blocks $a \rightarrow b \rightarrow a^+ \rightarrow b^+$ with $b, a^+, b^+ \in U$.

2) Define the constants $\theta = (\omega + 2)/4$ and $C = 1/((1 - \theta)\ell\sqrt{2\gamma})$. Choose $\delta > 0$ such that

$$9\delta + C2^{2(1-\theta)}\delta^{2(1-\theta)} < \frac{\epsilon}{4},$$

which implies $16\delta < \epsilon$. Then define the neighborhood V as $V = \mathcal{B}(x^*, \delta)$. We have to show that if the alternating sequence enters V , then it converges to a unique limit $b^* \in A \cap B$. By relabeling the sequence, we may without loss of generality assume that $b_0 \in V = \mathcal{B}(x^*, \delta)$. The case where the a_k 's reach V first is treated analogously.

We shall prove by induction that for every $k \geq 1$, we have

$$(16) \quad b_k, a_{k+1}, b_{k+1} \in \mathcal{B}(x^*, \epsilon)$$

and

$$(17) \quad \sum_{j=1}^k \|b_j - b_{j+1}\| \leq \frac{1}{2} \sum_{j=1}^k \|b_{j-1} - b_j\| + \frac{C}{2} (d_B(a_1)^{2(1-\theta)} - d_B(a_{k+1})^{2(1-\theta)}).$$

Let us first do the induction step and suppose that hypotheses (16), (17) are true at $k - 1$ for some $k \geq 2$. We have to show that they also hold at k .

2.1) Firstly we check (16) at k . By (16) at $k - 1$ we know that $b_{k-1}, a_k, b_k \in \mathcal{B}(x^*, \epsilon)$. So it remains to prove $a_{k+1}, b_{k+1} \in \mathcal{B}(x^*, \epsilon)$. We claim that $b_k \in \mathcal{B}(x^*, \frac{\epsilon}{4})$. Indeed, using the induction hypothesis (17) at $k - 1$ gives

$$\sum_{j=1}^{k-1} \|b_j - b_{j+1}\| \leq \frac{1}{2} \sum_{j=1}^{k-1} \|b_j - b_{j-1}\| + \frac{C}{2} (d_B(a_1)^{2(1-\theta)} - d_B(a_k)^{2(1-\theta)}).$$

Hence

$$\begin{aligned} \sum_{j=1}^{k-1} \|b_j - b_{j+1}\| &\leq \|b_0 - b_1\| + C d_B(a_1)^{2(1-\theta)} - C d_B(a_k)^{2(1-\theta)} - \|b_{k-1} - b_k\| \\ &\leq \|b_0 - b_1\| + C d_B(a_1)^{2(1-\theta)}. \end{aligned}$$

Therefore,

$$(18) \quad \begin{aligned} \|b_k - x^*\| &\leq \|b_k - b_1\| + \|b_1 - x^*\| \leq \sum_{j=1}^{k-1} \|b_{j+1} - b_j\| + \|b_1 - x^*\| \\ &\leq \|b_0 - b_1\| + C d_B(a_1)^{2(1-\theta)} + \|b_1 - x^*\|. \end{aligned}$$

Since $b_0 \in \mathcal{B}(x^*, \delta)$, we have $\|b_0 - a_1\| \leq \|b_0 - x^*\| \leq \delta$, hence $a_1 \in \mathcal{B}(x^*, 2\delta)$. Then $\|a_1 - b_1\| \leq \|a_1 - x^*\| \leq 2\delta$, which gives $\|b_1 - x^*\| \leq 4\delta$. It follows that $\|b_0 - b_1\| \leq 5\delta$. Now since $d_B(a_1) = \|a_1 - b_1\| \leq 2\delta$, going back to (18), we obtain

$$\|b_k - x^*\| \leq 9\delta + C2^{2(1-\theta)}\delta^{2(1-\theta)} < \frac{\epsilon}{4},$$

which is our above claim. Now this implies

$$\|a_{k+1} - x^*\| \leq \|a_{k+1} - b_k\| + \|b_k - x^*\| \leq 2\|b_k - x^*\| < \frac{\epsilon}{2} < \epsilon,$$

and

$$\|b_{k+1} - x^*\| \leq \|a_{k+1} - b_{k+1}\| + \|a_{k+1} - x^*\| \leq 2\|a_{k+1} - x^*\| < \epsilon.$$

This proves $a_{k+1} \in \mathcal{B}(x^*, \epsilon)$ and $b_{k+1} \in \mathcal{B}(x^*, \epsilon)$ and therefore (16) at k .

2.2) Let us now prove that (17) is true at k . Using the induction hypothesis (16) at $k-1$, that is, $b_{k-1}, a_k, b_k \in \mathcal{B}(x^*, \epsilon)$, we apply part 1) of the proof to the building block $b_{k-1} \rightarrow a_k \rightarrow b_k$, which gives

$$(19) \quad \frac{1 - \cos \alpha_k}{\|a_k - b_k\|^\omega} \geq \gamma,$$

where $\alpha_k = \angle(b_{k-1} - a_k, b_k - a_k)$. By part 2.1), which is already proved, we have $b_k, a_{k+1}, b_{k+1} \in \mathcal{B}(x^*, \epsilon)$ and $\mathcal{B}(x^*, \epsilon) \subset U$, so that we can apply the four-point estimate of part 1) to the building block $b_k \rightarrow a_{k+1} \rightarrow b_{k+1}$. This gives

$$(20) \quad d_B(a_{k+1})^2 + \ell \|b_k - b_{k+1}\|^2 \leq d_B(a_k)^2.$$

Now using the cosine theorem and (19) we obtain

$$\begin{aligned} \|b_{k-1} - b_k\|^2 &= \|b_{k-1} - a_k\|^2 + \|a_k - b_k\|^2 - 2\|b_{k-1} - a_k\| \|a_k - b_k\| \cos \alpha_k \\ &= (\|b_{k-1} - a_k\| - \|a_k - b_k\|)^2 + 2\|b_{k-1} - a_k\| \|a_k - b_k\| (1 - \cos \alpha_k) \\ &\geq (\|b_{k-1} - a_k\| - \|a_k - b_k\|)^2 + 2\gamma \|b_{k-1} - a_k\| \|a_k - b_k\|^{\omega+1} \\ &\geq 2\gamma \|b_{k-1} - a_k\| \|a_k - b_k\|^{\omega+1}. \end{aligned}$$

Since $b_k \in P_B(a_k)$ and $b_{k-1} \in B$, we have $\|b_{k-1} - a_k\| \geq \|a_k - b_k\| = d_B(a_k)$. Hence $\|b_{k-1} - b_k\|^2 \geq 2\gamma d_B(a_k)^{\omega+2}$, or what is the same

$$(21) \quad \|a_k - b_k\|^{-(\omega+2)/2} \|b_{k-1} - b_k\| \geq \sqrt{2\gamma}.$$

Recalling that $\theta = (\omega + 2)/4$ we have $\theta \in [\frac{1}{2}, 1)$, because $\omega \in [0, 2)$. By concavity of the function $s \mapsto s^{1-\theta}/(1-\theta)$ we have $s_1^{1-\theta} - s_2^{1-\theta} \geq (1-\theta)s_1^{-\theta}(s_1 - s_2)$. Applying this to $s_1 = d_B(a_k)^2$ and $s_2 = d_B(a_{k+1})^2$, we obtain

$$\begin{aligned} d_B(a_k)^{2(1-\theta)} - d_B(a_{k+1})^{2(1-\theta)} &\geq (1-\theta)d_B(a_k)^{-2\theta} (d_B(a_k)^2 - d_B(a_{k+1})^2) \\ &= (1-\theta)\|a_k - b_k\|^{-2\theta} (\|a_k - b_k\|^2 - \|a_{k+1} - b_{k+1}\|^2) \\ &\geq (1-\theta)\ell\sqrt{2\gamma} \frac{\|b_k - b_{k+1}\|^2}{\|b_{k-1} - b_k\|}, \end{aligned}$$

where the last estimate uses (20) and (21). Multiplying by $\|b_k - b_{k-1}\|$ and recalling that $C = 1/((1-\theta)\ell\sqrt{2\gamma})$, this gives

$$C (d_B(a_k)^{2(1-\theta)} - d_B(a_{k+1})^{2(1-\theta)}) \|b_k - b_{k-1}\| \geq \|b_k - b_{k+1}\|^2.$$

By comparison of the arithmetic and geometric mean, $a^2 \leq bc$ implies $a \leq \frac{1}{2}b + \frac{1}{2}c$ for positive a, b, c , hence we obtain

$$(22) \quad \|b_k - b_{k+1}\| \leq \frac{1}{2}\|b_k - b_{k-1}\| + \frac{C}{2} (d_B(a_k)^{2(1-\theta)} - d_B(a_{k+1})^{2(1-\theta)}).$$

By the induction hypothesis we have (17) at $k-1$, that is,

$$\sum_{j=1}^{k-1} \|b_j - b_{j+1}\| \leq \frac{1}{2} \sum_{j=1}^{k-1} \|b_{j-1} - b_j\| + \frac{C}{2} (d_B(a_1)^{2(1-\theta)} - d_B(a_k)^{2(1-\theta)}).$$

Adding this and (22) gives (17) at index k .

2.3) Let us now prove that the hypotheses (16) and (17) hold at $k = 1$. Concerning (16), since $b_1 \in \mathcal{B}(x^*, 4\delta)$ and $\|b_1 - a_2\| \leq 4\delta \leq \frac{\epsilon}{4}$, we have $a_2 \in \mathcal{B}(x^*, \frac{\epsilon}{2})$. Then using $\|a_2 - b_2\| \leq \frac{\epsilon}{2}$ gives $b_2 \in \mathcal{B}(x^*, \epsilon)$, so (16) is true at $k = 1$.

Concerning the validity of (17) at $k = 1$, observe that using $b_0, b_1, a_1 \in \mathcal{B}(x^*, \frac{\epsilon}{4})$ we may repeat the argument in the induction step starting before formula (20) with $k = 1$ in the place of k . The conclusion is formula (22) at $k = 1$, that is,

$$\|b_1 - b_2\| \leq \frac{1}{2}\|b_1 - b_0\| + \frac{C}{2} (d_B(a_1)^{2(1-\theta)} - d_B(a_2)^{2(1-\theta)}),$$

and this is precisely (17) at $k = 1$. This concludes the induction argument.

3) Having proved (16), (17) for all indices $k \geq 1$, we see from (18) that the series $\sum_{j=1}^{\infty} \|b_j - b_{j+1}\|$ converges, which means b_k is a Cauchy sequence, which converges to a limit $b^* \in B \cap \mathcal{B}(x^*, \epsilon)$. Using relation (21) we conclude that a_k converges to the same limit $b^* \in A \cap B$. \square

Our next result gives the convergence rate for $\omega \in (0, 2)$. The case $\omega = 0$, where linear convergence is obtained, will be treated separately in Theorem 3.

Corollary 4. (Rate of convergence). *Under the hypotheses of Theorem 1, with $\omega \in (0, 2)$, the convergence rates are $\|b_k - b^*\| = \mathcal{O}\left(k^{-\frac{2-\omega}{2\omega}}\right)$ and $\|a_k - b^*\| = \mathcal{O}\left(k^{-\frac{2-\omega}{2\omega}}\right)$.*

Proof: Summing (22) from $k = N$ to $k = M$ gives

$$-\frac{1}{2}\|b_N - b_{N-1}\| + \frac{1}{2} \sum_{k=N}^{M-1} \|b_k - b_{k+1}\| + \|b_M - b_{M+1}\| \leq \frac{C}{2} (d_B(a_N)^{2(1-\theta)} - d_B(a_{M+1})^{2(1-\theta)}).$$

Passing to the limit $M \rightarrow \infty$ gives

$$-\frac{1}{2}\|b_N - b_{N-1}\| + \frac{1}{2} \sum_{k=N}^{\infty} \|b_k - b_{k+1}\| \leq \frac{C}{2} d_B(a_N)^{2(1-\theta)}.$$

Introducing $S_N = \sum_{k=N}^{\infty} \|b_k - b_{k+1}\|$, this becomes

$$-\frac{1}{2}(S_{N-1} - S_N) + \frac{1}{2}S_N \leq \frac{C}{2} d_B(a_N)^{2(1-\theta)}.$$

Consequently,

$$\frac{1}{2}S_N \leq \frac{1}{2}(S_{N-1} - S_N) + C d_B(a_N)^{2(1-\theta)}.$$

Now using estimate (21), we have $d_B(a_N)^{2(1-\theta)} \leq (2\gamma)^{-\frac{1-\theta}{2\theta}} \|b_{N-1} - b_N\|^{\frac{1-\theta}{\theta}}$. Putting $C' := C(2\gamma)^{-\frac{1-\theta}{2\theta}}$ and substituting this gives

$$(23) \quad \frac{1}{2}S_N \leq \frac{1}{2}(S_{N-1} - S_N) + C' (S_{N-1} - S_N)^{\frac{1-\theta}{\theta}}.$$

Since $\theta \in (\frac{1}{2}, 1)$, we have $(1-\theta)/\theta \in (0, 1)$. Moreover, $S_N \rightarrow 0$ implies $S_{N-1} - S_N \rightarrow 0$, so the second term $(S_{N-1} - S_N)^{\frac{1-\theta}{\theta}}$ on the right of (23) dominates the first term $\frac{1}{2}(S_{N-1} - S_N)$. That means, there exists another constant $C'' > 0$ such that

$$S_N^{\frac{\theta}{1-\theta}} \leq C''(S_{N-1} - S_N)$$

for all $N \in \mathbb{N}$. We claim that there exists yet another constant C''' such that

$$(24) \quad 1 \leq C'''(S_{N-1} - S_N)S_N^{-\frac{\theta}{1-\theta}} \leq C''' \left(S_N^{-\frac{2\theta-1}{1-\theta}} - S_{N-1}^{-\frac{2\theta-1}{1-\theta}} \right).$$

Assuming this proved, summation of (24) from $N = 1$ to $N = M$ gives

$$M \leq C''' \left(S_M^{-\frac{2\theta-1}{1-\theta}} - S_1^{-\frac{2\theta-1}{1-\theta}} \right).$$

Hence for yet two other constants C'''' , C''''' ,

$$S_M \leq C'''' \left[S_1^{-\frac{2\theta-1}{1-\theta}} + M \right]^{-\frac{1-\theta}{2\theta-1}} \leq C''''' M^{-\frac{1-\theta}{2\theta-1}}.$$

Since $\|b_M - b^*\| \leq S_M$ by the triangle inequality, that proves the claimed speed of convergence.

In order to prove (24) we divide the set of indices into $\mathcal{I} = \{N : 2S_N \geq S_{N-1}\}$ and $\mathcal{J} = \{N : 2S_N < S_{N-1}\}$. For $N \in \mathcal{I}$ we have

$$\begin{aligned} (S_{N-1} - S_N) S_N^{-\frac{\theta}{1-\theta}} &\leq 2^{\frac{\theta}{1-\theta}} (S_{N-1} - S_N) S_{N-1}^{-\frac{\theta}{1-\theta}} \\ &\leq 2^{\frac{\theta}{1-\theta}} \int_{S_N}^{S_{N-1}} S^{-\frac{\theta}{1-\theta}} dS \\ &= 2^{\frac{\theta}{1-\theta}} \frac{1-\theta}{2\theta-1} \left(S_N^{-\frac{2\theta-1}{1-\theta}} - S_{N-1}^{-\frac{2\theta-1}{1-\theta}} \right), \end{aligned}$$

proving (24). In contrast, for $N \in \mathcal{J}$ we have

$$S_N^{-\frac{2\theta-1}{1-\theta}} - S_{N-1}^{-\frac{2\theta-1}{1-\theta}} \geq 2^{\frac{2\theta-1}{1-\theta}} S_{N-1}^{-\frac{2\theta-1}{1-\theta}} - S_{N-1}^{-\frac{2\theta-1}{1-\theta}} = \left(2^{\frac{2\theta-1}{1-\theta}} - 1 \right) S_{N-1}^{-\frac{2\theta-1}{1-\theta}} \rightarrow \infty$$

in view of $S_{N-1} \rightarrow 0$, $\frac{2\theta-1}{1-\theta} > 0$ and $2^{\frac{2\theta-1}{1-\theta}} - 1 > 0$. So on the set \mathcal{J} estimate (24) is also satisfied. Finally, the same estimate for a_k follows from $\|a_{k+1} - b^*\| \leq \|a_{k+1} - b_k\| + \|b_k - b^*\| \leq 2\|b_k - b^*\|$. \square

Theorem 2. (Local convergence with linear rate). *Let A, B intersect 0-separably at x^* with constant $\gamma \in (0, 2)$. Suppose B is 0-Hölder regular at x^* with respect to A with constant $c < \frac{\gamma}{2}$. Then there exists a neighborhood V of x^* such that every sequence of alternating projections that enters V converges R-linearly to a point $b^* \in A \cap B$.*

Proof: Applying Lemma 1 and Theorem 1 in the case $\omega = 0$, we obtain convergence of the sequence a_k, b_k to a point $b^* \in A \cap B$ from summability of $\sum_k \|b_{k-1} - b_k\|$. Now from the proof of Corollary 4, we see that in the case $\theta = \frac{1}{2}$ equation (23) simplifies to

$$\frac{1}{2} S_N \leq \frac{1}{2} (S_{N-1} - S_N) + C' (S_{N-1} - S_N),$$

or what is the same

$$S_N \leq \frac{1 + 2C'}{2 + 2C'} S_{N-1}.$$

This proves Q-linear convergence of S_N to 0, hence R-linear convergence of $b_N \rightarrow b^*$. \square

Remark 8. Theorem 2 extends the results in [21, Thm. 5.2] and [4, Thm. 3.14] in two ways. Firstly, as seen in example 7.1, 0-Hölder regularity includes sets B which have singularities pointing inwards, where superregularity [21] and its extension in [4] fail. Secondly, 0-separability is weaker than linear regularity or the CQ in [4], see example 7.4.

We now obtain the following consequence of Theorem 1, originally proved in [5] for more general families of sets. When specialized to the case of two sets we have

Corollary 5. (Bauschke et al. [5, Theorem 3.14]). *Suppose $(A, \tilde{A}, B, \tilde{B})$ satisfies the CQ-condition (2) at $x^* \in A \cap B$, where $P_A(\partial B \setminus A) \subset \tilde{A}$, $P_B(\partial A \setminus B) \subset \tilde{B}$. Moreover, suppose for every $\epsilon > 0$ there exists $\delta > 0$ such that B is (A, ϵ, δ) regular at x^* . Then there exists a neighborhood V of x^* such that every alternating sequence which enters V converges R -linearly to a point in $A \cap B$. \square*

Remark 9. While [4, Thm. 3.14] is slightly more restrictive than Theorem 2 as far as the regularity and transversality hypotheses are concerned, the authors of [4, 5] go on the other hand further than our present contribution in two ways. They discuss the case of more than two sets, and they quantify the size of the neighborhood on which the observed convergence rate is attained. A fine analysis of the proof of Lemma 1 and Corollary 4 should in principle allow a similar more quantitative version of Theorem 2.

As already shown in [5] one readily derives

Corollary 6. (Lewis, Luke, Malick [21]). *Suppose A, B intersect linearly regularly and B is superregular. Then alternating projections converge locally R -linearly to a point in the intersection. \square*

The following is now a consequence of Theorem 2, using Propositions 2 and 8.

Corollary 7. (Drusvyatskiy, Ioffe, Lewis [12]). *Suppose A, B intersect intrinsically transversally at x^* . Then there exists a neighborhood U of x^* such that every sequence of alternating projections entering U converges R -linearly to a point in the intersection.*

Proof: By Proposition 2 the sets A, B intersect 0-separably at x^* with constant $\gamma = \kappa^2/2 \in (0, 1]$. By Proposition 8, B is 0-Hölder regular with respect to A at x^* with any constant $c < \kappa^2/(1 - \kappa^2) = \gamma/(2 - \gamma)$. Choosing $c < \gamma/2$ allows us therefore to apply Theorem 2. \square

Remark 10. Drusvyatskiy *et al.* [12] stress that their approach gives local R -linear convergence under a transversality hypothesis alone, while the older [5, 21] as well as our present approach still need regularity assumptions. However, this statement should be read with care, because Propositions 2 and 8 show that intrinsic transversality amalgamates transversality and regularity aspects. In particular, it is more restrictive than 0-Hölder regularity in tandem with 0-separability, so that Theorem 2 is stronger than the main result in [12].

6. SUBANALYTIC SETS

Following [8], a subset A of \mathbb{R}^n is called *semianalytic* if for every $x \in \mathbb{R}^n$ there exists an open neighborhood V of x such that

$$(25) \quad A \cap V = \bigcup_{i \in I} \bigcap_{j \in J} \{x \in V : \phi_{ij}(x) = 0, \psi_{ij}(x) > 0\}$$

for finite sets I, J and real-analytic functions $\phi_{ij}, \psi_{ij} : V \rightarrow \mathbb{R}$. The set B in \mathbb{R}^n is called *subanalytic* if for every $x \in \mathbb{R}^n$ there exist a neighborhood V of x and a bounded semianalytic subset A of some $\mathbb{R}^n \times \mathbb{R}^m$, $m \geq 1$, such that $B \cap V = \{x \in \mathbb{R}^n : \exists y \in \mathbb{R}^m \text{ such that } (x, y) \in A\}$. Finally, an extended real-valued function $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ is called subanalytic if its graph is a subanalytic subset of $\mathbb{R}^n \times \mathbb{R}$.

We consider the function $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ defined as $f(x) = i_A(x) + \frac{1}{2}d_B(x)^2$, where i_A is the indicator function of A ; cf. [22, p. 84].

Lemma 2. *Let $f = i_A + \frac{1}{2}d_B^2$. Let $a^+ \in A$ be projected from $b \in B$ and $v = \lambda(b - a^+) \in N_A^p(a^+)$, where $\lambda \geq 0$. Then $v + a^+ - P_B(a^+) \subset \widehat{\partial}f(a^+)$.*

Proof: Note that $a^+ - P_B(a^+) \subset \widehat{\partial}(\frac{1}{2}d_B^2)(a^+)$ by [22, 1.3.3] or [25, Example 8.53]. Next observe that $\widehat{\partial}i_A(a^+) = \widehat{N}_A(a^+)$ by [22, Proposition 1.79] or [25, Exercice 8.14]. Hence $v \in N_A^p(a^+) \subset \widehat{N}_A(a^+) = \widehat{\partial}i_A(a^+)$ using [4, Lemma 2.4]. Finally, by the sum rule [19, Prop. 1.12] we have $\widehat{\partial}i_A(a^+) + \widehat{\partial}\frac{1}{2}d_B^2(a^+) \subset \widehat{\partial}f(a^+)$, which completes the proof. \square

Definition 6. Let $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ be lower semi-continuous with closed domain such that $f|_{\text{dom}f}$ is continuous. We say that f satisfies the Łojasiewicz inequality with exponent $\theta \in [0, 1)$ at the critical point x^* of f if there exists $\gamma > 0$ and a neighborhood U of x^* such that $|f(x) - f(x^*)|^{-\theta} \|g\| \geq \gamma$ for every $x \in U$ and every $g \in \widehat{\partial}f(x)$. \square

Here x^* is critical if $0 \in \partial f(x^*)$, see [22, 25].

Lemma 3. Suppose $f = i_A + \frac{1}{2}d_B^2$ satisfies the Łojasiewicz inequality with exponent $\theta \in [0, 1)$ at $a^* = b^* \in A \cap B$ and constant $\gamma > 0$. Then in fact $\theta > \frac{1}{2}$. Moreover, B intersects A separably with exponent $\omega = 4\theta - 2 \in (0, 2)$ and constant $\gamma' = 2^{-2\theta-1}\gamma^2$.

Proof: Note that $f(a^*) = 0$. Therefore there exists a neighborhood U of $a^* \in A \cap B$ such that

$$(26) \quad f(a^+)^{-\theta} \|g\| \geq \gamma > 0$$

for every $a^+ \in A \cap U$ and every $g \in \widehat{\partial}f(a^+)$. Now let $a \rightarrow b \rightarrow a^+ \rightarrow b^+$ be a building block with $a, b, a^+, b^+ \in U$. From Lemma 2, $g = v + a^+ - b^+ \in \widehat{\partial}f(a^+)$ for every $v \in N_A^p(a^+)$ of the form $v = \lambda(b - a^+)$. This uses the fact that $a^+ \in P_A(b)$. Hence by (26) we have

$$2^\theta d_B(a^+)^{-2\theta} \|\lambda(b - a^+) + a^+ - b^+\| \geq \gamma > 0$$

for every $\lambda \geq 0$. We deduce

$$(27) \quad d_B(a^+)^{-2\theta} \min_{\lambda \geq 0} \|\lambda(b - a^+) + a^+ - b^+\| \geq 2^{-\theta} \gamma.$$

Let us for the time being consider angles $\alpha = \angle(b - a^+, b^+ - a^+)$ smaller than 90° . Then the minimum value in (27) is $d_B(a^+)^{-2\theta} \|a^+ - b^+\| \sin \alpha$. Therefore

$$(28) \quad \frac{\sin \alpha}{d_B(a^+)^{2\theta-1}} \geq 2^{-\theta} \gamma.$$

Since $1 - \cos \alpha \geq \frac{1}{2} \sin^2 \alpha$, estimate (28) implies

$$(29) \quad \frac{1 - \cos \alpha}{d_B(a^+)^{4\theta-2}} \geq 2^{-2\theta-1} \gamma^2.$$

This shows that we must have $\theta > \frac{1}{2}$, because the numerator tends to 0, so the denominator has to go to zero, too, which it does for $4\theta - 2 > 0$.

Let us now discuss the case where $\alpha \geq 90^\circ$. We claim that the same estimate (29) is still satisfied. Since $\cos \alpha < 0$, the numerator $1 - \cos \alpha$ in (29) is ≥ 1 . Moreover, the infimum in (27) is now attained at $\lambda = 0$ with the value $\|a^+ - b^+\| = d_B(a^+)$. Hence estimate (27) implies $d_B(a^+)^{1-2\theta} \geq 2^{-\theta} \gamma$, hence $d_B(a^+)^{2-4\theta} \geq 2^{-2\theta} \gamma^2 > 2^{-2\theta-1} \gamma^2$, so that (29) is satisfied. This completes the proof. \square

Theorem 3. Let A, B be closed subanalytic sets. Then A, B intersect separably.

Proof: We assume $A \cap B \neq \emptyset$, otherwise there is nothing to prove. Consider the function $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$ defined as $f(x) = i_A(x) + \frac{1}{2}d_B(x)^2$. Then f has closed domain A and is continuous on A , which makes it amenable to definition 6. Every $x^* \in A \cap B$ is a critical

point of f . Since A, B are subanalytic sets, f is subanalytic. That can be seen as follows. First observe that d_B is subanalytic by [27, I.2.1.11]. Then d_B^2 is subanalytic as the product of two subanalytic functions by [27, I.2.1.9]. Finally, $\text{graph}(f) = (A \times \mathbb{R}) \cap \text{graph}(\frac{1}{2}d_B^2)$ shows that f is subanalytic.

Now we invoke Theorem 3.1 of [9], which asserts that f satisfies the Łojasiewicz inequality at x^* for some $\theta \in (0, 1)$. Hence (26) is true for every $g \in \partial f(a^+)$, and therefore also for every $g \in \widehat{\partial} f(a^+)$. Applying Lemma 3, we deduce that B intersects A separably with $\omega = 4\theta - 2$. Interchanging the roles of A and B , it follows also that A intersects B separably. \square

Corollary 8. (Local convergence for subanalytic sets). *Let A, B be subanalytic. Suppose B is Hölder regular at $x^* \in A \cap B$ with respect to A . Then there exists a neighborhood V of x^* such that every sequence of alternating projections a_k, b_k which enters V converges to some $b^* \in A \cap B$ with rate $\|b_k - b^*\| = \mathcal{O}(k^{-\rho})$ for some $\rho \in (0, \infty)$.* \square

Corollary 9. *Let A, B be closed subanalytic sets and suppose B has slowly vanishing reach with respect to A . Let $x^* \in A \cap B$, then there exists a neighborhood U of x^* such that every sequence of alternating projections a_k, b_k which enters U converges to some $b^* \in A \cap B$ with rate $\|b_k - b^*\| = \mathcal{O}(k^{-\rho})$ for some $\rho \in (0, \infty)$.* \square

Recall from [8, 27] that a subset A of \mathbb{R}^n is called *semialgebraic* if for every $x \in \mathbb{R}^n$ there exists a neighborhood V of x such that (25) is satisfied with ϕ_{ij}, ψ_{ij} polynomials. Naturally, this means that every semialgebraic set is semianalytic, hence subanalytic. By combining Theorems 1 and 3, we therefore obtain the following result.

Corollary 10. (Borwein, Li, Yao [10]). *Let A, B be closed convex semialgebraic sets with nonempty intersection. Then there exists $\rho \in (0, \infty)$ such that every sequence of alternating projections converges with rate $\|b_k - b^*\| = \mathcal{O}(k^{-\rho})$.* \square

As a variant of the method of alternating projects consider the averaged projection method. Given closed sets C_1, \dots, C_m , the method generates a sequence x_n by the recursion $x_{n+1} \in \frac{1}{m}(P_{C_1}(x_n) + \dots + P_{C_m}(x_n))$.

Corollary 11. *Let C_1, \dots, C_m be subanalytic sets in \mathbb{R}^d , and let $c^* \in C_1 \cap \dots \cap C_m$. Then there exists a neighborhood U of c^* such that whenever a sequence x_n of averaged projections enters U , then it converges to some $x^* \in C_1 \cap \dots \cap C_m$ with rate $\|x_k - x^*\| = \mathcal{O}(k^{-\rho})$ for some $\rho \in (0, \infty)$.*

Proof: We follow a standard procedure and define closed sets in the product space $\mathbb{R}^d \times \dots \times \mathbb{R}^d$ (m times) as $B = \{(x, \dots, x) : x \in \mathbb{R}^d\}$, and $A = C_1 \times \dots \times C_m$. Note that A is again subanalytic by [27, I.2.1.1], whereas B is convex and subanalytic. We have $P_B(x_1, \dots, x_m) = (\frac{1}{m}(x_1 + \dots + x_m), \dots, \frac{1}{m}(x_1 + \dots + x_m))$, while $P_A(x_1, \dots, x_m) = (P_{C_1}(x_1), \dots, P_{C_m}(x_m))$. Therefore a sequence of averaged projections between C_1, \dots, C_m generates a sequence of alternating projections between A, B .

Since B is convex, it is prox-regular hence Hölder regular with respect to A , so by Corollary 8 there exists a neighborhood $\mathcal{U} = U \times \dots \times U$ of $(c^*, \dots, c^*) \in A \cap B$ such that every alternating sequence which enters \mathcal{U} converges to some $(x^*, \dots, x^*) \in A \cap B$ with rate $\mathcal{O}(k^{-\rho})$ for some $\rho \in (0, \infty)$. Now consider an averaged projection sequence x_k entering U . It follows that $(x_k, \dots, x_k) \in \mathcal{U}$, hence x_k converges to x^* with that same rate. \square

Remark 11. We mention a related averaged projection method in [1, Corollary 12], where the authors use the Kurdyka-Łojasiewicz inequality. The employed technique indicates

that results in the spirit of Theorem 3 could be obtained for more general classes of sets definable in an o-minimal structure [1].

We conclude this section with an application of Theorem 1, demonstrating its versatility as a convergence test in practical situations. Let \mathbb{C}^N be a finite dimensional unitary space, and consider the discrete Fourier transform

$$\widehat{x}(\omega) = \frac{1}{\sqrt{N}} \sum_{t=0}^{N-1} e^{2\pi i t \omega / N} x(t), \quad \omega = 0, \dots, N-1$$

as a unitary linear operator $x \rightarrow \widehat{x}$ of \mathbb{C}^N . The phase retrieval problem [13, 16] consist in estimating an unknown signal $x \in \mathbb{C}^N$ whose Fourier amplitude $|\widehat{x}(\omega)| = a(\omega)$, $\omega = 0, \dots, N-1$, is known. In physical terminology, identifying x means retrieving its unknown phase $\widehat{x}(\omega)/|\widehat{x}(\omega)|$ in frequency domain.

Formally, given a function $a(\cdot) : \{0, \dots, N-1\} \rightarrow [0, \infty)$, we have to find an element of the set $B = \{x \in \mathbb{C}^N : |\widehat{x}(\omega)| = a(\omega) \text{ for all } \omega = 0, \dots, N-1\}$. Since this problem is underdetermined, additional information about x in a different Fourier plane or in the time domain is added. We represent it in the abstract form $x \in A$ for a closed set A . Then the phase retrieval problem is to find $x \in A \cap B$.

The famous Gerchberg-Saxton error reduction algorithm [16] computes a solution of the phase retrieval problem by generating a sequence of estimates as follows: Given $x \in \mathbb{C}^N$, compute \widehat{x} and correct its Fourier amplitude by putting $y(\omega) = a(\omega) \widehat{x}(\omega)/|\widehat{x}(\omega)|$ if $\widehat{x}(\omega) \neq 0$, and $y(\omega) = a(\omega)$ if there is extinction $\widehat{x}(\omega) = 0$. For short, $y = a\widehat{x}/|\widehat{x}|$ with the convention $0/|0| = 1$. Then compute the inverse discrete Fourier transform \widetilde{y} of y and build the new iterate x^+ by projecting \widetilde{y} on the set A , that is $x^+ \in P_A(\widetilde{y})$. In condensed notation:

$$(30) \quad x^+ \in P_A((a\widehat{x}/|\widehat{x}|)^\sim).$$

Corollary 12. (Gerchberg-Saxton error reduction). *Suppose the constraint $x \in A$ is represented by a closed subanalytic set A . Let $x^* \in A \cap B$ be a solution of the phase retrieval problem. Then there exists $\epsilon > 0$ such that whenever a Gerchberg-Saxton sequence x_k enters $\mathcal{B}(x^*, \epsilon)$, then it converges to a solution $\bar{x} \in A \cap B$ of the phase retrieval problem with rate of convergence $\|x_k - \bar{x}\| = \mathcal{O}(k^{-\rho})$ for some $\rho \in (0, \infty)$.*

Proof: With the convention $0/|0| = 1$, the mapping $x \mapsto (a\widehat{x}/|\widehat{x}|)^\sim$ is an orthogonal projection on the set $B = \{x \in \mathbb{C}^N : |\widehat{x}(\cdot)| = a(\cdot)\}$. (See for instance [3, (8),(10)], where the authors consider even the function space case). Therefore the Gerchberg-Saxton algorithm (30) is an instance of the alternating projection methods between the subanalytic set A and the Fourier amplitude set B . We show that B is subanalytic and prox-regular. Local convergence with rate $\mathcal{O}(k^{-\rho})$ then follows from Corollary 9.

As far as subanalyticity of B is concerned, observe that on identifying \mathbb{C}^N with \mathbb{R}^{2N} via $\widehat{x}(\omega) = \widehat{x}_1(\omega) + i\widehat{x}_2(\omega)$, we have

$$B = \bigcap_{\omega=0}^{N-1} \{x \in \mathbb{C}^N : \widehat{x}_1(\omega)^2 + \widehat{x}_2(\omega)^2 - a(\omega)^2 = 0\},$$

which is clearly a representation of the form (25), since the discrete Fourier transform $x \mapsto \widehat{x}$ is analytic.

To show prox-regularity of B , we have to show that the projection on B is single-valued in a neighborhood of B . With the same identification $\mathbb{C}^N \cong \mathbb{R}^{2N}$ evoked before, the projection on B splits into N projections in \mathbb{R}^2 , given as $(\widehat{x}_1(\omega), \widehat{x}_2(\omega)) \rightarrow a(\omega) \left(\frac{\widehat{x}_1(\omega)}{\sqrt{\widehat{x}_1(\omega)^2 + \widehat{x}_2(\omega)^2}}, \frac{\widehat{x}_2(\omega)}{\sqrt{\widehat{x}_1(\omega)^2 + \widehat{x}_2(\omega)^2}} \right)$. In the case $a(\omega) = 0$ this is the projection onto

the origin, which is clearly single-valued. For $a(\omega) > 0$ this is the orthogonal projection onto the sphere of radius $a(\omega)$ in \mathbb{R}^2 , which is single-valued except at the origin $(\hat{x}_1(\omega), \hat{x}_2(\omega)) = (0, 0)$. This means the projection on B is unique on the neighborhood $U = \{x \in \mathbb{C}^N : |\hat{x}(\cdot)| \geq a(\cdot)/2\}$ of B , proving prox-regularity of B . \square

Remark 12. The constraint $x \in A$ may represent additional measurements, or it may include prior information about the unknown image. In the original work [16], $x \in A$ represents Fourier amplitude information from a second Fourier plane, which is a constraint analogous to $x \in B$. The constraint $x \in A$ may also represent prior information about the support $\text{supp}(x)$ of the unknown signal x in physical domain. It may for instance be known that $\text{supp}(x) \subset S$, where S is a subset of $\{0, \dots, N-1\}$ with $\text{card}(S) \ll N$, or with a periodic structure. This is known as an atomicity constraint in crystallographic phase retrieval [13]. For $A = \{x \in \mathbb{C}^N : x(t) = 0 \text{ for } t \notin S\}$, P_A is simply truncation $y \rightarrow y \cdot 1_S$. Here the Gerchberg-Saxton error correction method has the explicit form

$$x^+(t) = \begin{cases} (a\hat{x}/|\hat{x}|)^\sim(t) & \text{if } t \in S \\ 0 & \text{else} \end{cases}$$

Other choices of the constraint $x \in A$ have been discussed in the literature, see e.g. [13]. Our convergence result requires only subanalyticity of A , a condition which is always satisfied in practice.

7. EXAMPLES

Example 7.1. (Packman gulping an ice-cream cone the wrong way). Consider packman $B = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1, x \leq |y|\}$ the instant before it scarfs down the ice-cream cone section $A = \{(x, y) \in \mathbb{R}^2 : 0 \leq x^2 + y^2 \leq 1, 2|y| \leq x\}$ fitting symmetrically into its notch. We have $A \cap B = \{(0, 0)\}$, leaving an angular gap of 15° on both sides.

Let $a^+ = (x, \frac{1}{2}x) \in \partial A$, then $b^+ = P_B(a^+) = (\frac{3}{4}x, \frac{3}{4}x) \in \partial B$, which means $r = \|b^+ - a^+\| = \frac{\sqrt{2}}{4}x$. It is easy to see that condition (6) is satisfied for every $c < 1$ and arbitrary $\sigma \in [0, 1)$, i.e., B is Hölder regular with respect to A . This example shows that Hölder regularity applies to sets which have inward corners and fail Clarke regularity.

Note that since B is not Clarke regular at $x^* = (0, 0)$, it is not superregular in the sense of [21]. What is more, B is not (A, ϵ, δ) -regular in the sense of [4] at $x^* = (0, 0)$, regardless how $\epsilon, \delta > 0$ are chosen, because the cone $b^+ + \{v : \langle a^+ - b^+, v \rangle \leq \epsilon \|a^+ - b^+\| \|v\|\}$ with vertex at the projected point $b^+ = (\frac{3}{4}x, \frac{3}{4}x) \in B$ hits B at points $b' \in B$ other than b on the opposite side of A , regardless how small ϵ is chosen. And this cannot be prevented by shrinking the neighborhood $\mathcal{B}(x^*, \delta)$. Note that A, B intersect 0-separably at $(0, 0)$, hence alternating projections converge linearly by Theorem 2. This cannot be obtained from the results in [4, 21].

In [17, Def. 2.9] Hesse and Luke discuss a related concept called (ϵ, δ) -subregularity. In [17, Thm 3.11] they prove linear convergence under (ϵ, δ) -subregularity of B with respect to $A \cap B$, in tandem with a transversality assumption at x^* , called local linear regularity. Note that B above is $(0, 1)$ -subregular with respect to $A \cap B = \{x^*\}$ at $x^* = (0, 0)$. However, if we change the set A to $A = \{(x, y) : 0 \leq x^2 + y^2 \leq 1, -\frac{1}{2}x \leq y \leq x, x \geq 0\}$, then B is no longer (ϵ, δ) -subregular with respect to $A \cap B$ at x^* , regardless how small $\epsilon \in (0, 1)$, $\delta > 0$ are chosen. Since now $S = A \cap B = \{(x, y) : x^2 + y^2 \leq 1, y = x, x \geq 0\}$, fixing $\delta > 0$, we can find $x \in B \cap \mathcal{B}(x^*, \delta)$, $x = (\xi, -\xi)$ with $\xi > 0$, and $\bar{x} = (\eta, \eta) \in S \cap \mathcal{B}(x^*, \delta)$ with $\eta > 0$ such that $\cos \angle(v_x, \bar{x} - x)$ is arbitrarily close to 1. Note that B is still 0-Hölder regular at x^* with respect to A , so Theorem 2 is still applicable. \square

Example 7.2. (Regularity cannot be dispensed with). Following [6], consider the spiral $z(\phi) = (1 + e^{-\phi})e^{i\phi}$, $\phi \in [0, \infty)$ in the complex plane which approaches the unit circle $S = \{|z| = 1\}$ from outside. Define a sequence $z_n = z(\phi_n)$ with $\phi_1 < \phi_2 < \dots \rightarrow \infty$ such that $\|z_{n+1} - z_n\| < \|z_n - z_{n-1}\| \rightarrow 0$, $P_{\{z_k : k \neq n\} \cup S}(z_n) = z_{n+1}$, and such that every $z \in S$ is an accumulation point of the z_n . In [6] an explicit construction with these properties is obtained recursively as

$$(31) \quad z_n = z(\phi_n), \quad \|z(\phi_{n+1}) - z(\phi_n)\| = r_n, \quad r_{n+1} = e^{-\phi_{n+1}} \frac{1 - e^{-2\pi}}{2}.$$

Let $A = \{z_{2n} : n \in \mathbb{N}\} \cup S$, $B = \{z_{2n-1} : n \in \mathbb{N}\} \cup S$, then $A \cap B = S$. Note that for starting points $|z_0| > 1$, the sequence of alternating projections between A and B is a tail of the sequence z_n , so none of the alternating sequences converges. Note that $\angle(z_{n+1} - z_n, z_{n-1} - z_n) \rightarrow \pi$, hence A, B intersect 0-separably at every $x^* \in S = A \cap B$. The CQ in the sense of [4] is satisfied at every $x^* \in A \cap B$. Namely, for $z \in S$, $N_A^B(z) = N_B^A(z) = \mathbb{R}_+(-iz)$. Indeed, as $a_n = P_A(b_n)$ approaches z , the direction $u_n = (b_n - a_n)/\|b_n - a_n\|$ approaches a direction perpendicular to z , and since the spiral turns counterclockwise, this direction is $-iz$. Therefore $N_A^B(z) \cap (-N_B^A(z)) = \{0\}$ for every $z \in S$.

Since the sequence z_n fails to converge, we conclude that this must be due to the lack of regularity at points in S . Indeed, Hölder regularity fails for every $0 \leq \sigma < 1$. This can be seen as follows. Since the angle $\angle(b - a^+, b^+ - a^+)$ for the building block $b \rightarrow a^+ \rightarrow b^+$ approaches π , the corresponding angle $\beta = \angle(b - b^+, a^+ - b^+)$ goes to 0, so $\cos \beta \rightarrow 1$, and for $\sigma \in (0, 1)$ we cannot find $c > 0$ such that $\cos \beta \leq \sqrt{c}r^\sigma$. Therefore, in order to assure (6), we would need $b \notin \mathcal{B}(a^+, (1+c)r)$, where $r = \|a^+ - b^+\|$. This however, would imply linear convergence of the alternating sequence, which fails. As a consequence of Proposition 3, the other regularity concepts fail, as does intrinsic transversality. \square

Example 7.3. (Discrete spiral I). We consider a discrete approximation of the logarithmic spiral, generated by 8 equally spaced rays emanating from the origin. Starting on one of the rays, we project perpendicularly on the neighboring ray, going counterclockwise. We label the projected points a_1, b_1, a_2, \dots . This defines two sets $A = \{a_i : i \in \mathbb{N}\} \cup \{(0, 0)\}$ and $B = \{b_i : i \in \mathbb{N}\} \cup \{(0, 0)\}$ with $A \cap B = \{(0, 0)\}$ such that $P_B(a_i) = b_i$ and $P_A(b_i) = a_{i+1}$. Every sequence of alternating projections between A and B not starting at the origin is a tail of the sequence a_n, b_n and converges to $(0, 0)$.

Since $\alpha = \angle(b - a^+, b^+ - a^+) = 135^\circ$, A, B intersect 0-separably at $x^* = (0, 0)$. We check whether the intersection satisfies the CQ in the sense of [4]. Consider one of the rays on which a point a^+ is situated. Then $u = b - a^+ \in \widehat{N}_A^B(a^+)$ is perpendicular to $a^+ - x^*$, i.e., perpendicular to the ray in question. As u is the same for all a^+ on that ray, we have $u \in N_A^B(0, 0)$. Altogether, $N_A^B(0, 0) = \text{lin}\{u_1, u_3, -u_1, -u_3\}$ for four directions spaced 90° . Similarly, $N_B^A(0, 0) = \{u_2, u_4, -u_2, -u_4\}$ spaced 90° , and intertwined with the directions of $N_A^B(0, 0)$. We have $N_A^B(0, 0) = -N_B^A(0, 0)$, and similarly for $N_B^A(0, 0)$, and since $N_B^A(0, 0) \cap \widetilde{N}_A^B(0, 0) = \{(0, 0)\}$, the intersection does indeed satisfy the CQ in the sense of [4] for $\widetilde{A} = A, \widetilde{B} = B$.

How about regularity at $(0, 0)$? Naturally, A, B are not superregular at $(0, 0)$, because they are not Clarke regular. Concerning (A, ϵ, δ) -regularity of B in the sense of [4], suppose in a building block $b \rightarrow a^+ \rightarrow b^+$ we wish to set up a cone with apex b^+ and axis $b^+ + \mathbb{R}_+(a^+ - b^+)$ by choosing its aperture small enough through the choice of ϵ such that all previous points of A are avoided, then we have to choose smaller and smaller angles β to do this, so this type of regularity fails.

On the other hand, we have σ -Hölder regularity for every $\sigma \in [0, 1)$. Suppose we start at $a_1 = (1, 0)$, then $b_1 = (\frac{1}{2}, \frac{1}{2})$ and $a_2 = (0, \frac{1}{2})$. After a tour of 360° , the spiral comes

back to the horizontal ray $\mathbb{R}^+(1, 0)$ at $a_5 = (\frac{1}{16}, 0)$. So while at the beginning the spiral turns within the square $[-1, 1]^2$, from the second tour onward it will stay in the square $[-\frac{1}{16}, \frac{1}{16}]^2$. As the circle $\mathcal{B}(b_1, \frac{7}{16}\sqrt{2})$ contains no points of the small square in its interior, the distance of b_1 to the small square being $R = \frac{7}{16}\sqrt{2}$, writing $R = (1+c)r$ we conclude that we can take $c = \frac{7}{8}\sqrt{2} - 1 > 0$ in (6). Now up to a scaling and a rotation the situation is precisely the same for *every* building block $a \rightarrow b \rightarrow a^+$ starting in a square of length $2\|a\|$. After one 360° -tour we end up at a^{++++} on the same ray as a , and from there on the spiral will stay in that smaller square of length $2\frac{1}{16}\|a\| = 2\|a^{++++}\|$. As a consequence of theorem 2, the sequence converges to $(0, 0)$ with linear rate. None of the approaches of [4, 20, 21] allows to derive this. \square

Example 7.4. (Discrete spiral II). We can modify the above construction by fixing $\phi \in (0, \frac{\pi}{4})$ and generating rays $k\phi$, $k \in \mathbb{N}$. Turning counterclockwise, and keeping only the projected points, we generate iterates a_k, b_k with the property that a_k has angle $2k\phi \bmod 2\pi$ with the horizontal, b_k has angle $(2k+1)\phi \bmod 2\pi$. We put $A = \{a_k : k \in \mathbb{N}\} \cup \{(0, 0)\}$, $B = \{b_k : k \in \mathbb{N}\} \cup \{(0, 0)\}$, then $A \cap B = \{(0, 0)\}$ and $P_B(a_k) = b_k$, $P_A(b_k) = a_{k+1}$ by adapting the argument in example 7.3. The sequence represents again a discrete version of the logarithmic spiral, turning inwards counterclockwise. However, if we now choose ϕ such that $\phi/(2\pi)$ is irrational, there will be no periodicity, and the set of directions $a_k/\|a_k\|$ will be dense in \mathbb{S}^1 , and so for $b_k/\|b_k\|$. We have $\angle(b^+ - a^+, b - a^+) = \pi - \phi$, which means A, B intersect 0-separably at $(0, 0)$. They intersect *at an angle*, this angle being $\pi - \phi$. However, A, B do not intersect linearly regularly in the sense of [4, 21]. Indeed, let us fix $\psi \in [0, 2\pi)$ and $u = (\cos \psi, \sin \psi)$. Then there exist rays $2k\phi$ arbitrarily close to ψ and a_k on these rays, projected from b_{k-1} on ray $(2k-1)\phi$. That means, $u_k = (b_k - a_k)/\|b_k - a_k\|$ gets arbitrarily close to the direction $u^\perp = (-\sin \psi, \cos \psi)$, so $u^\perp \in N_A^B(0, 0)$. This shows $N_A^{\tilde{B}}(0, 0) = \mathbb{R}^2$ and $N_B^{\tilde{A}}(0, 0) = \mathbb{R}^2$ for $\tilde{A} = P_A(\partial B \setminus A)$, $\tilde{B} = P_B(\partial A \setminus B)$, so linear regularity and extensions fail. \square

Example 7.5. (Spiral and cylinder [7]). Consider the cylinder $B = \{(x_1, x_2, x_3) : x_1^2 + x_2^2 \leq 1, 0 \leq x_3 \leq 1\}$ and the spiral $A = \{((1 + e^{-t}) \cos t, (1 + e^{-t}) \sin t, e^{-t/2}) : t \geq 0\} \cup S$, where $S = \{(\cos \alpha, \sin \alpha, 0) : \alpha \in [0, 2\pi]\}$. Clearly $A \cap B = S$. As shown in [7], any sequence of alternating projections between A and B started at $a \in A \setminus S$ wanders down following the spiral, turning infinitely often around the cylinder with shrinking $a_n - b_n \rightarrow 0$. In particular, every $x^* \in S$ is an accumulation point of a_n, b_n , so convergence fails. Since B , being convex, is clearly Hölder regular with respect to A , we deduce that the angle condition (1) must fail, so in particular A is not subanalytic. This is interesting, as A is the projection of an unbounded semianalytic set in \mathbb{R}^4 . For a picture see [7].

Example 7.6. (Failure of intrinsic transversality). We consider the sets $A = \{2^{-2n} : n \in \mathbb{N}\} \cup \{0\}$, $B = \{2^{-2n+1} : n \in \mathbb{N}\} \cup \{0\}$ in \mathbb{R} , so that $A \cap B = \{0\}$. The sequence of alternating projections is $1, \frac{1}{2}, \frac{1}{4}, \dots$ and converges Q-linearly to 0. We have $N_A^B(0) = N_B^A(0) = \mathbb{R}_+$, hence A, B intersect with the CQ in the sense of [4] at 0, hence also 0-separably. Note that B is not (A, ϵ, δ) -regular at 0 in the sense of [4], but it is σ -Hölder-regular for every $\sigma \in [0, 1)$. Note that intrinsic transversality fails here, because it uses the cones $N_A(a), N_B(b)$, which in this case are too large because they coincide with the whole line.

We modify this example as follows. Let $a_n = 2^{-n}$, $A = \{a_n : n \in \mathbb{N}\} \cup \{0\}$, $b_n = \frac{1}{2}(a_n + a_{n+1}) - \delta_n$, $B = \{b_n : n \in \mathbb{N}\} \cup \{0\}$, where $\delta_n < 2^{-n}(a_n - b_n)$. Then $\|a_{n+1} - b_n\|$ shrinks only by a factor $1 - \delta_n \rightarrow 1$ with respect to $\|b_n - a_n\|$, while shrinkage between $\|a_{n+1} - b_n\|$ and $\|a_{n+1} - b_{n+1}\|$ is by a factor close to $\frac{1}{2}$. This shows that an alternating

sequence may converge R-linearly without a fixed shrinkage factor $1 - \kappa^2$ in every half step. Note that Theorem 2 still applies in this case.

Example 7.7. We give an example where A, B intersect tangentially, but not ω -separably for any $\omega \in [0, 2)$. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable with f' continuous at 0, $f(0) = 0$, $f(x) > 0$ for $x \neq 0$, and define $A = \text{epi} f = \{(x, y) \in \mathbb{R}^2 : y \geq f(x)\}$, $B = \{(x, y) \in \mathbb{R}^2 : y \leq 0\}$, then $A \cap B = \{(0, 0)\}$. We consider a building block $b \rightarrow a^+ \rightarrow b^+$. Let $b^+ = (x, 0)$, then $a^+ = (x, f(x))$. Suppose $b = (y, 0)$, then $y = x + f(x)f'(x)$. Then the quotient q in (1') reads

$$q(x) = \frac{\sqrt{1 + f'(x)^2} - 1}{f(x)^\omega \sqrt{1 + f'(x)^2}} = \frac{f'(x)^2}{f(x)^\omega \sqrt{1 + f'(x)^2} (\sqrt{1 + f'(x)^2} + 1)} \leq \frac{f'(x)^2}{2f(x)^\omega}.$$

Therefore, if the angle condition (1) is to hold for some ω , then $\liminf_{x \rightarrow 0} \frac{f'(x)^2}{f(x)^\omega} \geq \gamma > 0$. It is possible to construct f such that this fails for every $\omega \in [0, 2)$. Take for instance

$$f(x) = \begin{cases} e^{-\frac{1}{x^2}} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases},$$

then $q(x) \leq 2x^{-6} \exp(-x^{-2}(2 - \omega)) \rightarrow 0$ as $x \rightarrow 0$ for $0 < \omega < 2$. Separability with $\omega = 0$ is also impossible because $f'(x) \rightarrow 0$ as $x \rightarrow 0$. In conclusion, the sets A and B intersects tangentially, but not separably for any $\omega \in [0, 2)$.

Example 7.8. Using the same function f and A, B , observe that for $\omega \geq 2$ the quotient $q(x)$ stays away from 0, so that condition (1') is satisfied. This explains why values $\omega \geq 2$ are not meaningful in definition 1.

REFERENCES

- [1] H. Attouch, J. Bolte, P. Redont, A. Soubeyran. Proximal alternating minimization and projection methods for nonconvex problems: an approach based on the Kurdyka-Łojasiewicz inequality. *Mathematics of Operations Research* 35(2) (2010), 438–457.
- [2] H. H. Bauschke, J. M. Borwein. On projection algorithms for solving convex feasibility problems. *SIAM Review* 38 (1996), 367–426.
- [3] H.H. Bauschke, P.L. Combettes, D.R. Luke. Phase retrieval, error reduction algorithm, and Fienup variants: a view from convex optimization. *J. Opt. Soc. Am. Series A* 19(7) (2002), 1334–1345.
- [4] H.H. Bauschke, D.R. Luke, H.M. Phan, X. Wang. Restricted normal cones and the method of alternating projections: theory. *Set-Valued and Variational Analysis* 21 (2013), 431–473.
- [5] H.H. Bauschke, D.R. Luke, H.M. Phan, X. Wang. Restricted normal cones and the method of alternating projections: applications. *Set-Valued and Variational Analysis* 21 (2013), 475–501.
- [6] H.H. Bauschke, D. Noll. On cluster points of alternating projections. *Serdica Math. Journal* 39 (2013), 355–364.
- [7] H.H. Bauschke, D. Noll. On the local convergence of the Douglas-Rachford algorithm. *Archiv der Mathematik* 102(6) (2014), 589–600.
- [8] E. Bierstone, P. Milman. Semianalytic and subanalytic sets. *IHES Publ. Math.* 67 (1988), 5–42.
- [9] J. Bolte, A. Daniilidis, A.S. Lewis. The Łojasiewicz inequality for nonsmooth subanalytic functions with applications to subgradient dynamical systems. *SIAM J. Opt.* 17(4) (2007), 1205–1223.
- [10] J.M. Borwein, G. Li, L. Yao. Analysis of the convergence rate for the cyclic projection algorithm applied to basic semialgebraic convex sets. *SIAM Journal on Optimization* 24(1) (2014), 498–527.
- [11] P.L. Combettes, H.J. Trussell. Method of successive projections for finding a common point of sets in metric space. *Journal of Optimization Theory and Applications* 67 (1990), 487–507.
- [12] D. Drusvyatskiy, A.D. Ioffe, A.S. Lewis. Alternating projections and coupling slope. arXiv:1401.7569v1 [math.OC] 29 Jan 2014.
- [13] V. Elser. Solution of the crystallographic phase problem by iterated projections. *Acta Crystallographica Section A.* 59 (2003), 201–209.
- [14] M. Fabian. Lipschitz smooth points of convex functions and isomorphic characterizations of Hilbert space. *Proc. London Math. Soc.* s3-51 (1985), 113–126.
- [15] H. Federer. Hausdorff measure and Lebesgue area. *Proc. Nat. Acad. Sci. U.S.A.* 37 (1951), 90–94.

- [16] R. W. Gerchberg and W. O. Saxton. A practical algorithm for the determination of the phase from image and diffraction plane pictures. *Optik* 35 (1972), 237–246.
- [17] R. Hesse, D. R. Luke. Nonconvex notions of regularity and convergence of fundamental algorithms for feasibility problems. *SIAM J. Opt.* 23(4) (2013), 2397–2419.
- [18] A.Y. Kruger. About regularity of collections of sets. *Set-Valued Analysis* 14(2) (2006), 187–206.
- [19] A.Y. Kruger. On Fréchet subdifferentials. *Optimization and related topics, 3. Journal of Mathematical Sciences* 116(3) (2003), 3325–3358.
- [20] A.S. Lewis, J. Malick. Alternating projections on manifolds. *Math. Oper. Res.* 33 (2008), 216–234.
- [21] A.S. Lewis, R. Luke, J. Malick. Local linear convergence for alternating and averaged non convex projections. *Found. Comp. Math.* 9 (2009), 485–513.
- [22] B.S. Mordukhovich. *Variational Analysis and Generalized Differentiation*. Springer, New York, 2006.
- [23] J. von Neumann. *Functional Operators, vol. II*. Princeton University Press, Princeton, 1950.
- [24] R.A. Poliquin, R.T. Rockafellar, L. Thibault. Local differentiability of distance functions. *Trans. Amer. Math. Soc.* 352 (2000), 5231–5249.
- [25] R.T. Rockafellar, R.J.B. Wets. *Variational Analysis*. Grundlehren der mathematischen Wissenschaften, Springer Verlag 317, 2009.
- [26] H.A. Schwarz. Über einige Abbildungsaufgaben. *Gesammelte Mathematische Abhandlungen* 11 (1869), 65–83.
- [27] M. Shiota. *Geometry of Subanalytic and Semialgebraic Sets*. Birkhäuser Verlag, 1997.