

Asymptotic behavior of divergences and Cameron-Martin theorem on loop spaces

Xiang Dong Li*
University of Oxford, Oxford, UK

August 14, 2002

Abstract

We first prove the L^p -convergence ($p \geq 1$) and a Fernique type exponential integrability of divergence functionals for all Cameron-Martin vector fields with respect to the pinned Wiener measure on loop spaces over a compact Riemannian manifold. We then prove that the Driver flow is a smooth transform on path spaces in the sense of the Malliavin calculus and has an ∞ -quasi-continuous modification which can be quasi-surely well-defined on path spaces. This leads us to construct the Driver flow on loop spaces through the corresponding flow on path spaces. Combining these two results with the Cruzeiro lemma [4] [24], we give an alternative proof of the quasi-invariance of the pinned Wiener measure under Driver's flow on loop spaces which was established earlier by Driver [7] and Enchev-Stroock [9] by Doob's h -processes approach together with the short time estimates of the gradient and the Hessian of the logarithmic heat kernel on compact Riemannian manifolds. We also establish the L^p -convergence ($p \geq 1$) and a Fernique type exponential integrability theorem for the stochastic anti-development of pinned Brownian motions on compact Riemannian manifold with an explicit exponential exponent. Our results generalize and sharpen some earlier results due to E. Hsu [15] and L. Gross [13]. Our method do not need any heat kernel estimate and is based on quasi-sure analysis and Sobolev estimates on path spaces.

Contents

1 Introduction

2

*AMS 2000 subject classifications: 60H07, 58G32. *Keywords and phrases:* Divergence, pinned Wiener measure, Airault-Malliavin-Sugita-Watanabe inequality, exponential integrability, Driver flow, quasi-invariance, stochastic anti-development. Research partially supported by a Postdoctoral Research Fellowship of the University of Oxford.

2	The Airault-Malliavin-Sugita-Watanabe inequality	7
3	Sobolev estimates of divergence functionals on path spaces	9
4	The $L^p(\nu)$-convergence of divergence functionals	14
5	Exponential integrability of divergence functionals	18
6	Smoothness of Driver's flow on path spaces	21
7	Cameron-Martin theorem on loop spaces	28
8	Stochastic anti-development of pinned Brownian motions	31
9	Driver's flow on loop spaces via Doob's h-theory	34

1 Introduction

Let M be a compact connected Riemannian manifold, $m_0 \in M$ be a fixed point. Let ∇ be the Levi-Civita connection on M . The orthonormal frame bundle over M is denoted by $O(M)$. The path space (respectively, the loop space) over M is defined by $P_{m_0}(M) = \{\gamma \in C([0, 1], M) : \gamma(0) = m_0\}$ (respectively, $L_{m_0}(M) = \{\gamma \in P_{m_0}(M) : \gamma(1) = m_0\}$). Let H be the \mathbb{R}^d -valued Cameron-Martin space, i.e., the set of absolutely continuous functions $h : [0, 1] \rightarrow \mathbb{R}^d$ such that $h(0) = 0$ and $\dot{h} \in L^2([0, 1])$. Let H_0 its subspace with zero values at time 1, i.e., $H_0 = \{h \in H : h(1) = 0\}$.

Let Δ be the Laplace-Beltrami operator on M . The Wiener measure on $P_{m_0}(M)$, denoted by μ , is the law of M -valued Brownian motion (with generator $\Delta/2$) starting at m_0 . The pinned Wiener measure on $L_{m_0}(M)$, denoted by ν , is the law of the conditional Brownian motion $\{\gamma(s), s \in [0, 1]\}$ on M such that $\gamma(0) = \gamma(1) = m_0$. Intuitively, we have

$$\nu(\cdot) = \mu(\cdot | \gamma(1) = m_0).$$

Rigorously, if $p_t(x, y)$ denotes the heat kernel on M , then for any $\alpha < 1$,

$$\frac{d\nu}{d\mu} \Big|_{\mathcal{F}_\alpha}(\gamma) = \frac{p_{1-\alpha}(\gamma(\alpha), m_0)}{p_1(m_0, m_0)}. \quad (1.1)$$

where $\mathcal{F}_\alpha = \sigma(\gamma(s), s \in [0, \alpha])$. For details, see e.g. Bismut [2] and Driver [7].

For μ -a.s. $\gamma \in P_{m_0}(M)$ (respectively, ν -a.s. $\gamma \in L_{m_0}(M)$), one can use the Itô SDE theory to define the stochastic parallel transport $U_s(\gamma) : T_{m_0}M \rightarrow T_{\gamma(s)}M$ as the unique $O(M)$ -valued stochastic process satisfying the following covariant SDE

$$\nabla_{\circ d\gamma(s)} U_s(\gamma) = 0$$

with the initial condition $U_0(\gamma) = Id_{T_{m_0}M}$, where $Id_{T_{m_0}M}$ is the identity transform over $T_{m_0}M$. See e.g. [2]. For all $h \in H$ (respectively, $h \in H_0$), the Cameron-Martin vector field D_h on $P_{m_0}(M)$ (respectively, $L_{m_0}(M)$) is defined by: for μ -a.s. $\gamma \in P_{m_0}(M)$ (respectively, ν -a.s. $\gamma \in L_{m_0}(M)$),

$$D_h(\gamma)(s) = U_s(\gamma)h(s), \quad \forall s \in [0, 1]. \quad (1.2)$$

In [6] [8] [14], the classical Cameron-Martin theorem has been generalized to the path space $(P_{m_0}(M), \mu)$. That is to say, for all fixed $h \in H$, the vector field D_h generates a global flow $\{\Phi_t, t \in \mathbb{R}\}$ which can be μ -a.s. well-defined on $P_{m_0}(M)$ (see Section 6 below) under which the Wiener measure is quasi-invariant and an integration by parts formula holds. Concerning the same issue on the loop space $(L_{m_0}(M), \nu)$, Driver [7] proved that for any $h \in \mathcal{C}^1 \cap H_0$ (the set of Lipschitz Cameron-Martin vectors $h \in H_0$), the vector field D_h generates a global flow $\{\tilde{\Phi}_t : t \in \mathbb{R}\}$ on $L_{m_0}(M)$ such that for ν -a.s. $\gamma \in L_{m_0}(M)$,

$$\dot{\tilde{\Phi}}_t(\gamma) = D_h(\tilde{\Phi}_t(\gamma)), \quad \tilde{\Phi}_0(\gamma) = \gamma,$$

and the Wiener measure ν is quasi-invariant under the flow $\tilde{\Phi}_t$, i.e., the measure $(\tilde{\Phi}_t)_*\nu$ is equivalent to ν . Moreover, an integration by parts formula holds on $(L_{m_0}(M), \nu)$: for two cylindrical functionals F and G on $L_{m_0}(M)$,

$$E_\nu(D_h FG) = E_\nu(F(-D_h G + \delta(h)G)),$$

and

$$\frac{d(\tilde{\Phi}_t)_*\nu}{d\nu}(\gamma) = \exp\left(\int_0^t \delta(h)(\tilde{\Phi}_{-s}(\gamma))ds\right), \quad (1.3)$$

where $\delta(h)$ is the so-called divergence functional on $(L_{m_0}(M), \nu)$ defined as follows: for ν -a.s. $\gamma \in L_{m_0}(M)$,

$$\delta(h)(\gamma) = \int_0^1 (\dot{h}(s) + \frac{1}{2} Ric_{U_s(\gamma)}(h(s)), dx(s)). \quad (1.4)$$

Here Ric denotes the Ricci curvature form over $O(M)$ and $\{x(s), s \in [0, 1]\}$ is the stochastic anti-development of $\{\gamma(s), s \in [0, 1]\}$, denoted by $x = I^{-1}(\gamma)$, and is given by the following Stratonovich stochastic integral

$$x(s) = \int_0^s U_r^{-1}(\gamma) \circ d\gamma_r, \quad s \in [0, 1]. \quad (1.5)$$

The complete theory of integration by parts formula on the loop space $(L_{m_0}(M), \nu)$ for all the Cameron-Martin vector fields D_h with $h \in H_0$ was first proved by Enchev and Stroock [9] where the authors have also proved the quasi-invariance of the pinned

Wiener measure under the flow generated by D_h . In [15] [16], E. Hsu gave another approach to integration by parts formula on loop spaces which avoids the problem of the quasi-invariance of the pinned Wiener measure on the loop space. Let us mention that all the approaches appeared in [7] [9] [15] [16] relied strongly on the short time upper bound estimates on the gradient and the Hessian of the logarithmic of the heat kernel and all these authors used the Doob h -processes method for conditional Brownian motion on a compact Riemannian manifold. See also Section 9 below.

The purpose of this paper is to study the asymptotic behavior of divergence functionals, Driver flow and Cameron-Martin theorem on loop spaces as well as some related problems by a different approach. First, we will use the Airault-Malliavin-Sugita-Watanabe inequality (see Section 2) and some Sobolev estimates on the divergence functionals on the path space (see Section 3) to prove the L^p -convergence ($p \geq 1$) and the Fernique type exponential integrability of divergence functionals with respect to the pinned Wiener measure on loop spaces, see Section 4 and Section 5. Second, we will prove that the Driver flow $\{\Phi_t, t \in \mathbb{R}\}$ is a smooth transform on path spaces in the sense of the Malliavin calculus and has ∞ -quasi-continuous version denoted by $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$ which can be quasi-surely well defined up to a slim subset of the path space $P_{m_0}(M)$. Moreover, we prove that if $h \in H_0$ then $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$ actually realizes the Driver flow generated by the vector field D_h on the loop space $(L_{m_0}(M), \nu)$, see Section 6 and Section 7. Third, we will combine these two results with the Cruzeiro lemma [4] [24] to give an alternative approach to the complete Cameron-Martin theorem on the loop space $(L_{m_0}(M), \nu)$ avoiding to use any heat kernel estimate, see Section 7. Finally, we use our method to establish the $L^p(\nu)$ -convergence ($p \geq 1$) and a Fernique type exponential integrability theorem for the stochastic anti-development of pinned Brownian motions on a compact Riemannian manifold equipped with any torsion-skew symmetric (TSS) connection, see Section 8. Our results generalize and sharpen some earlier results due to Hsu [15] [16] and L. Gross [13]. Our method is inspired by M. P. Malliavin and P. Malliavin [25] where the authors first established the quasi-invariance of the pinned Wiener measure on the loop group over a compact Lie group under the left or the right action of a finite energy loop (which is nothing else than the Driver flow on the loop group). In some sense, it leads us to get *sharp or better* estimate than the direct approach based on Doob's h -theory and heat kernel estimates, see Section 8 and Section 9.

To state our main results, let us follow Hsu [15] [16] to introduce a sequence of functionals as follows: for any $h \in H$ and $s < 1$, let

$$\delta_s(h)(\gamma) = \int_0^s (\dot{h}(s) + \frac{1}{2} Ric_{U_s(\gamma)}(h(s)), dx(s)), \quad x = I^{-1}(\gamma).$$

Note that $\delta_s(h)$ is μ -a.s. well-defined on $P_{m_0}(M)$. Since ν is equivalent to μ on \mathcal{F}_s , cf. (1.1), we see that $\delta_s(h)$ is also well-defined for ν -a.s. $\gamma \in L_{m_0}(M)$.

Now we are in a position to state our main results of this paper.

Theorem 1.1 *Let $h \in H_0$. Then the divergence functional $\delta(h)$ (formally given by (1.4)) can be realized as the $L^p(\nu)$ -limit of $\delta_s(h)$ as $s \rightarrow 1$ for all $p \geq 1$. In fact, for all $p \geq 1$, there is a constant C_p such that for all $h \in H_0$,*

$$\|\delta_s(h) - \delta(h)\|_{L^p(\nu)} \leq C_p \left(\int_s^1 |\dot{h}(t)|^2 dt \right)^{\frac{1}{2}},$$

and

$$\|\delta(h)\|_{L^p(\nu)} \leq C_p \|h\|_{H_0}.$$

Moreover, for all

$$\lambda < \lambda_0 = \frac{1}{(2 + \|Ric\|_\infty) \|h\|_H},$$

we have

$$E_\nu [\exp(\lambda |\delta(h)|^2)] < +\infty,$$

or, equivalently,

$$\lim_{t \rightarrow \infty} \frac{1}{t^2} \log \nu(\{\gamma \in L_{m_0}(M) : |\delta(h)| > t\}) = -\frac{1}{(2 + \|Ric\|_\infty) \|h\|_H}.$$

Corollary 1.2 *For all $p > 1$, the gradient operator D on the loop space $L_{m_0}(M)$ is closable from $L^p(L_{m_0}(M), \nu)$ into $L^p(L_{m_0}(M), H_0, \nu)$.*

Corollary 1.3 *For all $h \in H_0$ and all $\lambda > 0$, we have*

$$E_\nu [\exp(\lambda |\delta(h)|)] < \infty.$$

Theorem 1.4 *For any $h \in H$, there exists an ∞ -quasi-continuous version of the Driver flow $\{\Phi_t, t \in \mathbb{R}\}$ which can be well-defined up to a slim subset of $P_{m_0}(M)$.*

Let $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$ be a fixed ∞ -quasi-continuous version of $\{\Phi_t, t \in \mathbb{R}\}$. By the disintegration principle of the Wiener measure, $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$ can be ν -a.s. well-defined on $L_{m_0}(M)$.

Theorem 1.5 *Let $h \in H_0$. Then $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$ is the flow generated by the vector field D_h on $L_{m_0}(M)$. Moreover, the pinned Wiener measure ν is quasi-invariant under $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$. More precisely,*

$$\frac{d(\tilde{\Phi}_t)_* \nu}{d\nu}(\gamma) = \exp \left(\int_0^t \delta(h)(\tilde{\Phi}_{-s}(\gamma)) ds \right), \quad \nu - \text{a.s. } \gamma \in L_{m_0}(M),$$

and for all $p > 1$ with the conjugate exponent q , i.e., $\frac{1}{p} + \frac{1}{q} = 1$, we have

$$\|K_t\|_{L^p(\nu)}^p \leq E [e^{pqt|\delta(h)|}].$$

Remark 1.1 *All the above results (as well as the Sobolev norms and capacities comparison inequalities (6.17) and (6.18) in Section 6 below) remain true if we replace the Levi-Civita connection by any torsion skew-symmetric (TSS) connection. In this case, we need only to replace the Ricci curvature Ric of the Levi-Civita connection by the Ricci curvature $\widehat{\text{Ric}}$ of the dual connection $\widehat{\nabla}$ given by*

$$\widehat{\nabla}_X Y = \nabla_X Y - T(X, Y), \quad X, Y \in \Gamma(TM),$$

where T is the torsion of our given TSS connection ∇ . Indeed, we have announced Theorem 1.1 in [20] (without giving the precise value λ_0) in this setting with an equivalent expression of the divergence functional $\delta(h)$ as used in Driver[7]. In particular, we recapture the Malliavin-Malliavin theorem on the quasi-invariance of the pinned Wiener measure on loop group over a compact Lie group.

The following result generalizes and sharpens some earlier results due to L. Gross [13] where the author established the $L^p(\nu)$ -convergence and a Fernique type exponential integrability theorem for the stochastic anti-development of pinned Brownian motions on a compact Lie group.

Theorem 1.6 *Let M be a compact Riemannian manifold equipped with a TSS connection, $m \in \mathbb{N}$, $m \geq 2$, $\alpha \in (\frac{1}{2m}, \frac{1}{2})$. Then for any*

$$\lambda < \lambda_0 := \frac{1}{2} \inf\{\|w\|_H^2 : w \in X, \|w\|_{2m, \alpha} = 1\},$$

we have

$$E_\nu [\exp(\lambda \|x\|_{2m, \alpha}^2)] < +\infty,$$

where

$$\|x\|_{2m, \alpha} = \left[\int_0^1 \int_0^1 \frac{\|x(t) - x(s)\|_{\mathbb{R}^d}}{|t - s|^{1+2m\alpha}} dt ds \right]^{\frac{1}{2m}},$$

and

$$x(s) = \int_0^1 U_s^{-1}(\gamma) \circ d\gamma(s), \quad s \in [0, 1], \quad \nu - \text{a.s. } \gamma \in L_{m_0}(M).$$

Moreover, for any $\lambda < \frac{1}{2}$, we have

$$E_\nu \left[\exp \left(\lambda \sup_{s \in [0, 1]} \|x(s)\|^2 \right) \right] < +\infty.$$

In addition, $x(s)$ converges to $x(1)$ in $L^p(\nu)$ for all $p \geq 1$ as s tends to 1 and there exists a constant C_p such that

$$\|x(s) - x(1)\|_{L^p(\nu)} \leq C_p (1 - s)^{\frac{1}{2}}.$$

2 The Airault-Malliavin-Sugita-Watanabe inequality

Let $X = \{x \in C([0, 1], \mathbb{R}^d) : x(0) = 0\}$ be the Wiener space, μ_0 be the Wiener measure on X . For any $r \in \mathbb{N}$ and $p > 1$, we let $W^{r,p}(X)$ denote the (r, p) -Sobolev space on the Wiener space X with the Sobolev norm $\|\cdot\|_{W^{r,p}(X)}$. Let A_1, \dots, A_d be the canonical horizontal vector fields on $O(M)$, $r_0 \in O(M)$ be a fixed orthonormal frame over m_0 . Consider the horizontal SDE on $O(M)$:

$$\begin{cases} dr_x(s) &= \sum_{i=1}^d A_i(r_x(s)) \circ dx^i(s) \\ r_x(0) &= r_0. \end{cases}$$

Let $\gamma_x(s) := \pi(r_x(s))$, $s \in [0, 1]$. Then it is well-known that $\{\gamma_x(s), s \in [0, 1]\}$ is a Brownian motion on M starting at m_0 . The Wiener measure μ on the path space $P_{m_0}(M) = \{\gamma \in C([0, 1], M) : \gamma(0) = m_0\}$ is given by the law of $\{\gamma_x(s), s \in [0, 1]\}$, i.e., $\mu = I_*\mu_0$, where $I : X \rightarrow P_{m_0}(M)$ is the Itô map given by

$$I(x) = \gamma_x, \quad \mu_0 - \text{a.s. } x \in X.$$

Consider the following M -valued Wiener functional Φ :

$$\Phi(x) = \gamma_x(1).$$

By [1], $\Phi \in W^{\infty,\infty}(X, M)$ and Φ is non-degenerated, i.e.,

$$(\text{Det}[\Phi(x)])^{-1} \in W^\infty(X, M),$$

where $W^{\infty,\infty}(X, M)$ is the set of all smooth M -valued Wiener functionals in the sense of Malliavin calculus, and

$$\text{Det}[\Phi(x)] = \sqrt{\det[\nabla\Phi(x) \cdot \nabla\Phi(x)^\tau]},$$

where the determinant on the right hand side is taken with respect to the Riemannian metric on $T_{\Phi(x)}M$, $\nabla\Phi(x)^\tau$ denotes the adjoint of $\nabla\Phi(x) : H \rightarrow T_{\Phi(x)}M$.

Recall that if $f \in W^{\infty,\infty}(X, M)$, then f has an ∞ -quasi-continuous modification which can be well-defined outside a slim subset of X . Moreover, if f_1, f_2 are two ∞ -quasi-continuous modifications of f , then f_1 and f_2 only differ on a slim set. Let Φ^* be any quasi-continuous modification of Φ . The following co-area formula is well-known (see e.g. [1] [24]): there exists a family of area measures denoted by $\{da^y(\cdot), y \in M\}$ (where each $da^y(\cdot)$ is supported on the submanifold $S_y = \Phi^{*-1}(y)$) such that for any $u \in W^\infty(X)$ and $v \in C^\infty(M)$,

$$\int_X u(x)v(\Phi(x))[\text{Det}\Phi](x)d\mu_0(x) = \int_M v(y) \int_{\Phi^{*-1}(y)} u^*(x)da^y(x)dy,$$

where u^* denotes any ∞ -quasi-continuous modification of u . Let

$$\nu_y(dx) = (\text{Det}[\Phi](x))^{-1} da^y(x).$$

Then for all $y \in M$, ν_y is a Borel probability measure supported on the submanifold $S_y = \Phi^{*-1}(y)$. Moreover, ν_y has no charge on any slim subset of X .

By [33] [21], the Itô map $I : X \rightarrow P_{m_0}(M)$ is smooth in the sense of the Malliavin calculus and has an ∞ -quasi-continuous modification. Throughout this paper, we let \tilde{I} denote a fixed ∞ -quasi-continuous modification of the Ito map I . Using the dyadic polygonal approximation of M -valued Brownian motion, and by a similar argument used in Section 4 in Shigekawa [30], one can prove that the stochastic anti-development map given by Eq. (1.5) has an ∞ -quasi-continuous version (denoted by \tilde{I}^{-1}) which can be quasi-surely well defined on $P_{m_0}(M)$ and satisfies $\tilde{I} \circ I = Id_X$ quasi-surely (i.e., except on a slim set of X). By this and using the capacity comparison inequality due to the author [21], we can easily prove that for two different versions of ∞ -quasi-continuous modification of I , saying I_1 and I_2 , $I_1|_{S_{m_0}}(S_{m_0})$ only differ from $I_2|_{S_{m_0}}(S_{m_0})$ on a slim subset of $P_{m_0}(M)$. [Indeed, let $S = \{x : I_1(x) \neq I_2(x)\}$. Then S is a slim set of X . Let $O \subset X$ be an open set containing S and with capacity $C_{r,p}(O) < \epsilon$ for all $r \in \mathbb{N}$ and $p > 1$. By the capacity comparison inequality between the path space and the Wiener space (see [21]), we have

$$\hat{C}_{r,p}(I_i(O)) \leq \alpha C_{2r,p+1}(I_i^{-1} \circ I_i(O)), \quad i = 1, 2,$$

where $\alpha = \alpha(r, p)$ is a constant and $\hat{C}_{r,p}$ is the (r, p) -capacity on the path space $P_{m_0}(M)$ (for its definition, see [21]). Note that $I_i^{-1} \circ I_i = Id_X$ holds quasi-surely on X , $i = 1, 2$. Hence $C_{2r,p+1}(I_i^{-1} \circ I_i(O)) = C_{2r,p+1}(O) < \epsilon$, $i = 1, 2$. Since ϵ is arbitrary, we get $\hat{C}_{r,p}(I_i(S)) = 0$, $\forall r \in \mathbb{N}, p > 1, i = 1, 2$.] Thus, $\tilde{I}|_{S_{m_0}}$, the restriction of \tilde{I} on the submanifold $S_{m_0} = \{x \in X : \gamma_x(1) = m_0\}$, is ν_{m_0} -a.s. well defined. Moreover, $\tilde{I}|_{S_{m_0}} : (S_{m_0}, \nu_{m_0}) \rightarrow (L_{m_0}(M), \nu)$ is a measure theoretic isomorphism. That is to say, $\tilde{I}|_{S_{m_0}}(S_0)$ only differs from $L_{m_0}(M)$ on a slim set of the path space $P_{m_0}(M)$ and

$$\nu = (\tilde{I}|_{S_{m_0}})_* \nu_{m_0}.$$

The following result is due to Airault-Malliavin [1] and Sugita [33].

Theorem 2.1 *There exist a pair $(p, r) \in (1, +\infty) \times \mathbb{N}$ and a constant $C > 0$ such that for any $f \in W^\infty(X, \mathbb{R}^+)$, we have*

$$\int_{S_{m_0}} f^*(x) \nu_{m_0}(dx) \leq C \|f\|_{W^{r,p}(X)},$$

where f^* is any ∞ -quasi-continuous modification of f .

In fact, using the Watanabe generalized distribution theory on Wiener space, we can even precise the constant C and the value of the pair (r, p) appeared in Theorem 2.1 as follows.

Theorem 2.2 (*Airault-Malliavin-Sugita-Watanabe inequality*) *For all $p > 1$ and $f \in W^{2[\frac{d}{2}]+2,p}(X)$, we have the*

$$\left| \int_{S_{m_0}} f^*(x) d\nu_{m_0}(x) \right| \leq C \|\delta_{m_0} \circ \Phi\|_{-2([\frac{d}{2}]+1), \frac{p}{p-1}} \|f\|_{2[\frac{d}{2}]+2,p}.$$

where

$$C = [p_1(m_0, m_0)]^{-1}.$$

Proof. For any $y \in M$, let δ_y be the Dirac delta function at point y . Then $\delta \in \mathcal{S}_{-2r}(M)$ for all $r \geq [\frac{d}{2}]+1$, where $d = \dim M$ and $\mathcal{S}_{-2r}(M)$ is the topological dual of $\mathcal{S}_{2r}(M)$ (the completion of the Schwartz space $\mathcal{S}(M)$ of rapidly decreasing C^∞ -functions on M by the norm $\|\cdot\|_{2r}$ defined by $\|\phi\|_{2r} = \|(1+|x|^2-\Delta)^r \phi\|_\infty$). See e.g. the proof of Theorem 4.2 in H. Sugita [32] (p. 47–48) and S. Watanabe [34] Remark 2.2 (p. 8). Let $F \in W^{\infty,\infty}(X, M)$ be a smooth non-degenerate Wiener functional. Then for all $k = 0, 1, \dots$ and $p > 1$, the map $y \in M \rightarrow \delta_y(F) \in W^{-2[\frac{d}{2}]-2-2k,p}(X)$ is $2k$ -times continuous differentiable. Hence for any $f \in D^{2[\frac{d}{2}]+2+2k,p}(X)$ we have

$$\left| \int_X f(x) \delta_y(F(x)) d\mu_0(x) \right| \leq \|\delta_y \circ F\|_{-2[\frac{d}{2}]-2-2k, \frac{p}{p-1}} \|f\|_{2[\frac{d}{2}]+2+2k,p}.$$

In particular, using the fact that

$$\nu_{m_0}(dx) = \frac{\delta_{m_0}(\Phi(x))}{\int_X \delta_{m_0}(\Phi(x)) d\mu_0(x)} d\mu_0(x),$$

we deduce the Airault-Malliavin-Sugita-Watanabe inequality with the constant C given by (cf. Watanabe [34])

$$C = \left[\int_X \delta_{m_0}(\Phi(x)) d\mu_0(x) \right]^{-1} = [p_1(m_0, m_0)]^{-1}.$$

□

3 Sobolev estimates of divergence functionals on path spaces

Following [21], for any $r \in \mathbb{N}$ and $p > 1$ we let $D^{r,p}(P_{m_0}(M))$ denote the (r, p) -Sobolev space on $P_{m_0}(M)$ with the Sobolev norm $\|\cdot\|_{D^{r,p}(P_{m_0}(M))}$ defined by

$$\|F\|_{D^{r,p}(P_{m_0}(M))} = \sum_{k=0}^r \left\| \|D^k F\|_{H^{\otimes k}} \right\|_p.$$

For any fixed $t \in [0, 1]$, regarding $x \rightarrow r_x(t)$ as an $O(M)$ -valued Wiener functional, we have $r_x(t) \in W^{\infty, \infty}(X, O(M))$. More precisely, for any $n \in \mathbb{N}$ and any $h, h_1, \dots, h_n \in H$, the following H -directional derivatives exist:

$$\begin{aligned} \nabla_h r_x(t) &:= \left. \left\{ \frac{d}{d\epsilon} r_{x+\epsilon h}(t) \right\} \right|_{\epsilon=0}, \\ \dots &\dots \\ \nabla_{h_1 \dots h_n} r_x(t) &:= \left. \left\{ \frac{D^n}{\partial \epsilon_1 \dots \partial \epsilon_n} r_{x + \sum_{i=1}^n \epsilon_i h_i}(t) \right\} \right|_{\epsilon_1 = \dots = \epsilon_n = 0}. \end{aligned}$$

where $\frac{D}{d\epsilon_i}$ denotes the Levi-Civita covariant-derivative along the smooth curve $\epsilon_i \mapsto r_{x+\epsilon_i h_i}(t)$ on $O(M)$. Moreover, we have the following

Proposition 3.1 ([21]) *There exist $D_{s_1 \dots s_n}^{j_1 \dots j_n} r_x(t) \in L^2([0, 1]^n, T_{r_x(t)} O(M))$ such that*

(1) *$D_{s_1 \dots s_n}^{j_1 \dots j_n} r_x(t)$ is adapted with respect to $\mathcal{F}_t = \sigma(x(s), s \in [0, t])$ for any fixed $s_1, \dots, s_n \in [0, 1]$ and $0 \leq j_1, \dots, j_n \leq n$. Moreover,*

$$D_{s_1 \dots s_n}^{j_1 \dots j_n} r_x(t) = 0 \quad \text{if } s_1 \vee \dots \vee s_n \in [t, 1],$$

(2) *for any $h_1, \dots, h_n \in H$,*

$$\langle \nabla^n r_x(t), h_1 \otimes \dots \otimes h_n \rangle_{H^{\otimes n}} = \sum_{j_1, \dots, j_n} \int_0^t \dots \int_0^t D_{s_1 \dots s_n}^{j_1 \dots j_n} r_x(t) \dot{h}_1^{j_1}(s_1) \dots \dot{h}_n^{j_n}(s_n) ds_1 \dots ds_n,$$

(3) *for any $p \geq 1$, we have*

$$\sup_{s_1, \dots, s_n \in [0, 1]} E \left[\sup_{s_1 \vee \dots \vee s_n \leq s \leq 1} \left\| D_{s_1 \dots s_n}^{j_1 \dots j_n} r_x(s) \right\|^p \right] < +\infty, \quad (3.1)$$

where $\|D_{s_1 \dots s_n}^{j_1 \dots j_n} r_x(s)\|$ denotes the Riemannian norm of the vector field $D_{s_1 \dots s_n}^{j_1 \dots j_n} r_x(s)$ with respect to the Sasaki Riemannian metric on $O(M)$ (for its definition, see [21]).

Let e_1, \dots, e_d be the standard orthonormal basis of \mathbb{R}^d , R be the Riemannian curvature tensor on M . For any $r \in O(M)$, the Ricci curvature over the frame r is a real matrix given by

$$Ric_r(a) = \sum_{i=1}^d r^{-1} \circ R(re_i, ra) \circ re_i, \quad \forall a \in \mathbb{R}^d.$$

Let

$$J(x, t) = \frac{1}{2} Ric_{r_x(t)}.$$

By the chain rule and Proposition 3.1, we have $J(x, t) \in W^\infty(X, M(d, d))$, where $M(d, d)$ denotes the set of all $d \times d$ real matrix. Moreover, we have the following

Proposition 3.2 *The Malliavin derivatives $D_{s_1 \dots s_n}^{j_1 \dots j_n} J(x, t)$ belong to $L^2([0, 1]^n, M(d, d))$ and are adapted with respect to $\mathcal{F}_t = \sigma(x(s), s \in [0, t])$ for any fixed $s_1, \dots, s_n \in [0, 1]$ and any $0 \leq j_1, \dots, j_n \leq n$. For any $h_1, \dots, h_n \in H$,*

$$\langle \nabla^n J(x, t), h_1 \otimes \dots \otimes h_n \rangle = \int_0^t \dots \int_0^t D_{s_1 \dots s_n}^{j_1 \dots j_n} J(x, t) \dot{h}_1^{j_1}(s_1) \dots \dot{h}_n^{j_n}(s_n) ds_1 \dots ds_n.$$

Moreover, for any $p \geq 1$,

$$\sup_{s_1, \dots, s_n \in [0, 1]} E \left[\sup_{s_1 \vee \dots \vee s_n \leq t \leq 1} \left\| D_{s_1 \dots s_n}^{j_1 \dots j_n} J(x, t) \right\|_{H.S}^p \right] < +\infty. \quad (3.2)$$

Proof. The proof can be easily given by the chain rule and using Proposition 3.1. In particular, (3.1) yields (3.2). \square

Let $h \in H$ and D_h be the vector fields on $P_{m_0}(M)$ defined by (1.2). By integration by parts formula, see e.g. [2] [6] [10] [8] [14], the $L^2(\mu)$ adjoint of D_h is given by $D_h^* = -D_h + \delta(h)$, where $\delta(h)$ is the divergence functional. Moreover, for μ -a.s. $\gamma \in P_{m_0}(M)$, we have

$$\delta(h)(\gamma) = \int_0^1 (\dot{h}(\tau) + \frac{1}{2} Ric_{r_x(\tau)} h(\tau), dx(\tau)), \quad x = I^{-1}(\gamma).$$

Now we state the main result in this section.

Theorem 3.3 *For any $r \in \mathbb{N}, p > 1$, there is a constant $C > 0$ such that for all $h \in H$ we have*

$$\|\delta(h)\|_{D^{r,p}(P_{m_0}(M))} \leq C \|h\|_H.$$

Proof. Let $\widehat{\delta(h)} = \delta(h) \circ I$. By the Sobolev norm comparison theorem (see [21]), we have

$$\|\delta(h)\|_{D^{r,p}(P_{m_0}(M))} \leq \alpha_{r,p} \|\widehat{\delta(h)}\|_{W^{2r,p+1}(X)},$$

where $\alpha_{r,p}$ is a constant. Hence we need only to prove that for any $r \in \mathbb{N}, p \geq 2$,

$$\|\widehat{\delta(h)}\|_{W^{r,p}(X)} \leq C \|h\|_H. \quad (3.3)$$

By induction and direct computation, it can be easily to show that for any $n \geq 2$ and $h, h_1, \dots, h_n \in H$, we have

$$D_s \widehat{\delta(h)}(x) = \dot{h}(s) + J(x, s)h(s) + \int_0^1 \langle D_s J(x, t)h(t), dx(t) \rangle,$$

$$D_{s_1 \dots s_n}^{j_1 \dots j_n} \widehat{\delta(h)}(x) = \int_0^1 \langle D_{s_1 \dots s_n}^{j_1 \dots j_n} J(x, t) h(t), dx(t) \rangle + \sum_{i=1}^n \left(D_{s_1 \dots \widehat{s}_i \dots s_n}^{j_1 \dots \widehat{j}_i \dots j_n} J(x, s_i) h(s_i) \right)^{j_i}, \quad (3.4)$$

where we use the notation $(a_1, \dots, a_d)^j := a_j$, $j = 1, \dots, n$.

By the Burkholder-Davies-Gundy inequality, we have $\|\widehat{\delta(h)}\|_p \leq C\|h\|_H$. It remains to prove that for any $n \geq 1, p \geq 2$, there is a constant $C > 0$ such that

$$\left\| \|\nabla^n \widehat{\delta(h)}\|_{H.S} \right\|_{L^p(\mu_0)} \leq C\|h\|_H. \quad (3.5)$$

Below we only give a proof of (3.5) for $n \geq 2, p \geq 2$. The proof for the case of $n = 1, p \geq 2$ is analogue. By definition, we have

$$\begin{aligned} \|\nabla^n \widehat{\delta(h)}\|_{H.S}^p &= \left[\sum_{j_1, \dots, j_n \in [0,1]^n} \int \left| D_{s_1 \dots s_n}^{j_1 \dots j_n} \widehat{\delta(h)} \right|^2 ds_1 \dots ds_n \right]^{\frac{p}{2}} \\ &\leq 2^{\frac{p}{2}} \left[\sum_{j_1, \dots, j_n \in [0,1]^n} \int \left| \int_0^1 \langle D_{s_1 \dots s_n}^{j_1 \dots j_n} J(x, t) h(t), dx(t) \rangle \right|^2 ds \right]^{\frac{p}{2}} \\ &\quad + 2^{\frac{p}{2}} \left[\sum_{j_1, \dots, j_n \in [0,1]^n} \sum_{i=1}^n \int \left| (D_{s_1 \dots \widehat{s}_i \dots s_n}^{j_1 \dots \widehat{j}_i \dots j_n} J(x, s_i) h(s_i))^{j_i} \right|^2 ds \right]^{\frac{p}{2}} \\ &\leq I_1 + I_2. \end{aligned}$$

By the Hölder inequality and the Burkholder-Davis-Gundy inequality, we have

$$\begin{aligned} I_1 &= E \left[\int_{[0,1]^n} \left| \int_0^1 \langle D_{s_1 \dots s_n}^{j_1 \dots j_n} J(x, t) h(t), dx(t) \rangle \right|^2 ds \right]^{\frac{p}{2}} \\ &\leq \int_{[0,1]^n} E \left| \int_0^1 \langle D_{s_1 \dots s_n}^{j_1 \dots j_n} J(x, t) h(t), dx(t) \rangle \right|^p ds \\ &\leq C \int_{[0,1]^n} E \left[\int_0^1 |D_{s_1 \dots s_n}^{j_1 \dots j_n} J(x, t) h(t)|^2 dt \right]^{\frac{p}{2}} ds \\ &\leq C \|h\|_H^p \int_{[0,1]^n} E \left(\sup_{t \in [v_{s_i}, 1]} |D_{s_1 \dots s_n}^{j_1 \dots j_n} J(x, t)|^p \right) ds \end{aligned}$$

$$\begin{aligned}
& \text{(by (3.2))} \\
& \leq C \|h\|_H^p.
\end{aligned}$$

Similarly, we have

$$\begin{aligned}
I_2 &= E \left[\int_{[0,1]^n} \left| (D_{s_1 \dots \widehat{s}_i \dots s_n}^{j_1 \dots \widehat{j}_i \dots j_n} J(x, s_i) h(s_i))^{j_i} \right|^2 ds \right]^{\frac{p}{2}} \\
&\leq E \left[\int_0^1 \left(\sup_{s_i \in [\vee s_j, 1]} \left| (D_{s_1 \dots \widehat{s}_i \dots s_n}^{j_1 \dots \widehat{j}_i \dots j_n} J(x, s_i) \right|^2 h(s_i) ds_i \right) \right]^{\frac{p}{2}} \\
&\leq \|h\|_H^p \sup_{s_j \in [0,1]} E \left[\sup_{s_i \in [\vee s_j, 1]} \left| (D_{s_1 \dots \widehat{s}_i \dots s_n}^{j_1 \dots \widehat{j}_i \dots j_n} J(x, s_i) \right|^p \right] \\
&\quad \text{(by (3.2))} \\
&\leq C \|h\|_H^p.
\end{aligned}$$

Combining the above inequalities for I_1 and I_2 , we obtain (3.5) and hence (3.3). \square

Proposition 3.4 *For any $n \in \mathbb{N}, p > 1$, there is a constant $C > 0$ such that for all $h \in H$ we have*

$$\left\| \int_{s_1}^{s_2} \left\langle \dot{h}(t) + \frac{1}{2} Ric_{r_x(t)}(h(t)), dx(t) \right\rangle \right\|_{D^{r,p}(P_{m_0}(M))} \leq C \left[\left(\int_{s_1}^{s_2} |\dot{h}(t)|^2 dt \right)^{\frac{1}{2}} + \left(\int_{s_1}^{s_2} |h(t)|^2 dt \right)^{\frac{1}{2}} \right]. \quad (3.6)$$

Proof. The proof is similar to the one of Theorem 3.3. \square

Corollary 3.5 *The functional $\delta(h)$ has an ∞ -quasi-continuous modification which can be ν -a.s. well-defined on $L_{m_0}(M)$.*

Proof. By Theorem 3.3, $\delta(h) \in D^{\infty, \infty}(P_{m_0}(M)) = \bigcap_{r \in \mathbb{N}, p > 1} D^{r,p}(P_{m_0}(M))$. Thus, $\delta(h)$ is a smooth functional on $P_{m_0}(M)$. Hence it has an ∞ -quasi-continuous modification (see [21]) which can be well-defined outside of a slim set. Thus we obtain a ν -a.s. well-defined functional on the loop space $L_{m_0}(M)$. \square

4 The $L^p(\nu)$ -convergence of divergence functionals

For any $h \in H_0$ and $s < 1$, the following functionals are well-defined for μ -a.s. $\gamma \in P_{m_0}(M)$,

$$\delta_s(h)(\gamma) = \int_0^s (\dot{h}(\tau) + \frac{1}{2} Ric_{r_x(\tau)} h(\tau), dx(\tau)), \quad x = I^{-1}(\gamma). \quad (4.1)$$

Since μ and ν are equivalent on $\mathcal{F}_s = \sigma(\gamma_x(\tau), \tau \leq s)$, $\delta_s(h)$ is also ν -a.s. well-defined on $L_{m_0}(M)$. The main technique part in E. Hsu's proof of the integration by parts formula on the loop space, see Proposition 4.1 in [15], is to prove that as $s \rightarrow 1$ the sequence $\{\delta_s(h)\}$ converges in $L^1(\nu)$ to a limit which belongs to $L^2(\nu)$. In view of this, for ν -a.s. $\gamma \in L_{m_0}(M)$, Hsu defined $\delta(h)(\gamma)$ as the $L^1(\nu)$ -limit of $\delta_s(h)(\gamma)$ and then prove that $\delta(h)$ is nothing else than the divergence functional appeared in an integration by parts formula on $L_{m_0}(M)$.

The purpose of this section is to prove that for all $p \geq 1$ as $s \rightarrow 1$ the sequence $\{\delta_s(h)\}$ converges to $\widetilde{\delta}(h)$ in $L^p(\nu)$, where $\widetilde{\delta}(h)$ is an ∞ -quasi-continuous modification of $\delta(h)$ constructed by the quasi-sure analysis principle which is ν -a.s. well-defined on $L_{m_0}(M)$ (see Corrolary 3.5 above). Moreover, we prove that $\widetilde{\delta}(h)$ satisfies the Driver-Enchev-Stroock-Hsu integration by parts formula on the loop space.

Theorem 4.1 *For any $p \geq 1$, there exists a constant $C_p > 0$ such that for all $h \in H$, we have*

$$\|\widetilde{\delta}(h)\|_{L^p(\nu)} \leq C_p \|h\|_H. \quad (4.2)$$

Moreover, for any $h \in H_0$,

$$\|\delta_s(h) - \widetilde{\delta}(h)\|_{W^{r,p}(X)} \leq C \left(\int_s^1 |\dot{h}(t)|^2 dt \right)^{\frac{1}{2}}. \quad (4.3)$$

$$\|\delta_s(h) - \widetilde{\delta}(h)\|_{L^p(\nu)} \leq C \left(\int_s^1 |\dot{h}(t)|^2 dt \right)^{\frac{1}{2}}. \quad (4.4)$$

Proof. By the Hölder inequality, we need only to prove Theorem 4.1 for $p = 2n$, $n \in \mathbb{N}$. By Theorem 2.1, there exist $r \in \mathbb{N}$ and $q > 1$ such that

$$\|\widetilde{\delta}(h)\|_{L^p(\nu)}^p \leq C \left\| \left(\widetilde{\delta}(h) \right)^p \right\|_{W^{r,q}(X)}.$$

It remains to prove that for any $r \in \mathbb{N}$ and $q > 1$,

$$\left\| \nabla^r \left(\widetilde{\delta}(h) \right)^p \right\|_{L^q(\mu_0)} \leq C \|h\|_H^p. \quad (4.5)$$

By the Burkholder-Davies-Gundy inequality, we have

$$\left\| \left(\widehat{\delta(h)} \right)^p \right\|_{L^q(\mu)} \leq C \|h\|_H^p. \quad (4.6)$$

Thus (4.5) holds for $r = 0$ and $q > 1$. Now we prove (4.5) for $r = 1$. By the Hölder inequality, we have

$$\begin{aligned} \left\| \nabla(\widehat{\delta(h)})^p \right\|_{L^q(\mu_0)} &= \left\| p(\widehat{\delta(h)})^{p-1} \nabla \widehat{\delta(h)} \right\|_{L^q(\mu_0)} \\ &\leq p \left\| \left(\widehat{\delta(h)} \right)^{p-1} \right\|_{L^{2q}(\mu_0)} \left\| \nabla \widehat{\delta(h)} \right\|_{L^{2q}(\mu_0)} \\ &\leq pC \left\| \widehat{\delta(h)} \right\|_{L^{2(p-1)q}(\mu_0)}^{p-1} \|h\|_H \\ &\quad \text{(using (3.5) and (4.6))} \\ &\leq pC \|h\|_H^p. \end{aligned}$$

In general, for any $k \in \mathbb{N}$ we have

$$\begin{aligned} \nabla^{k+1} f^{2n} &= 2n \nabla^k f^{2n-1} \otimes \nabla f + 2n f^{2n-1} \nabla^{k+1} f \\ &= 2n(2n-1) \nabla^{k-1} f^{2n-2} \otimes \nabla f \otimes \nabla f \\ &\quad + 2n(2n-1) f^{2n-2} \nabla^k f \otimes \nabla f + 2n f^{2n-1} \nabla^{k+1} f. \end{aligned}$$

Thus, by induction and using the Hölder inequality together with (3.5) and (4.6), we can prove (4.5) for $p = 2n$ and all $r \in \mathbb{N}$.

For any $s < 1$, we can easily prove that $\delta_s(h) \circ I \in W^{\infty, \infty}(X)$. Hence $\delta_s(h) \in D^{\infty, \infty}(P_{m_0}(M))$. Let $\delta_s(h)^*$ be any ∞ -quasi-continuous modification of $\delta_s(h)$. Then $\delta_s(h)^*$ is also well-defined for $\nu - a.s. \gamma \in L_{m_0}(M)$ and $\delta_s(h) = \delta_s(h)^*$ holds ν -a.s. on $L_{m_0}(M)$. By Theorem 2.1 and by the same argument used in the proof of (4.2) together with Proposition 3.4, we can prove that

$$\begin{aligned} \|\delta_{s_2}(h)^* - \delta_{s_1}(h)^*\|_{L^p(\nu)}^p &= \|(\delta_{s_2}(h) - \delta_{s_1}(h))^*\|_{L^p(\nu)}^p \\ &\leq C \|\delta_{s_2}(h) - \delta_{s_1}(h)\|_{W^{r,q}(X)}^p \\ &= C \left\| \int_{s_1}^{s_2} (\dot{h}(t) + J(x, t)h(t), dx(t)) \right\|_{W^{r,q}}^p \\ &\leq C \left[\left(\int_{s_1}^{s_2} |\dot{h}(t)|^2 dt \right)^{p/2} + \left(\int_{s_1}^{s_2} |h(t)|^2 dt \right)^{p/2} \right], \end{aligned} \quad (4.7)$$

where C is a constant which only depends on p . Now for $h \in H_0$, we have $\int_s^1 |h(t)|^2 dt \leq \int_s^1 |\dot{h}(t)|^2 dt$. Taking $s_2 = 1$ in (4.7), we obtain (4.3) and (4.4). \square

Now we introduce the gradient operator on $P_{m_0}(M)$ and $L_{m_0}(M)$. Let F be a cylindrical functional on $P_{m_0}(M)$ (respectively, $L_{m_0}(M)$) given by

$$F(\gamma) = f(\gamma_{s_1}, \dots, \gamma_{s_k}),$$

where $f \in C^\infty(M^k)$, $0 = s_0 < s_1 < \dots < s_k < s_{k+1} = 1$ is a finite partition of $[0, 1]$, $k \in \mathbb{N}$. For μ -a.s. $\gamma \in P_{m_0}(M)$ (respectively, ν -a.s. $\gamma \in L_{m_0}(M)$), we define the gradient $DF(\gamma)$ of F as the unique element of H (respectively, H_0) such that for any $h \in H$ (respectively, $h \in H_0$),

$$\langle DF(\gamma), h \rangle_H = D_h F(\gamma).$$

Here

$$D_h F(\gamma) = \sum_{i=1}^k \langle \text{grad}^{(i)} f(\gamma(s_i)), U_{s_i}(\gamma) h(s_i) \rangle, \quad \nu - \text{a.s. } \gamma \in L_{m_0}(M),$$

and $\text{grad}^{(i)} f(\gamma(s_i))$ denotes the gradient of f with respect to the i -th variable, $i = 1, \dots, k$.

Below we prove that $\widetilde{\delta(h)}$ satisfies the Driver-Enchev-Stroock-Hsu integrate by parts formula on the loop space.

Theorem 4.2 *Let F, G be two cylindrical functionals on $L_{m_0}(M)$, $h \in H_0$. Then*

$$\langle D_h F, G \rangle_{L^2(\nu)} = \langle F, D_h^* G \rangle_{L^2(\nu)},$$

where

$$D_h^* = -D_h + \widetilde{\delta(h)}.$$

Proof. First, we suppose $h \in H_1 = \{h \in H_0 : \text{supp}(h) \subset\subset (0, 1)\}$. Then there exists some $\alpha \in [0, 1)$ such that $h(\tau) = 0, \forall \tau \in [\alpha, 1]$. Without loss of the generality we may suppose that F and G are \mathcal{F}_α -measurable. To simplify the notation, let $p_t(x) = p_t(m_0, x)$, $x \in M$. Note that

$$D_h p_{1-\alpha}(\gamma(\alpha)) = \text{grad} p_{1-\alpha}(\gamma(\alpha)) U_\alpha(\gamma) h(\alpha) = 0.$$

By (1.1) and the integration by parts formula on the path space, we have

$$\begin{aligned}
& \int_{L_{m_0}(M)} D_h F(\gamma) G(\gamma) d\nu(\gamma) = \int_{P_{m_0}(M)} D_h F(\gamma) G(\gamma) \frac{p_{1-\alpha}(\gamma(\alpha))}{p_1(m_0)} d\mu(\gamma) \\
& = \int_{P_{m_0}(M)} F(\gamma) D_h^* \left(G(\gamma) \frac{p_{1-\alpha}(\gamma(\alpha))}{p_1(m_0)} \right) d\mu(\gamma) \\
& = \int_{P_{m_0}(M)} F(\gamma) \left[-D_h \left(G(\gamma) \frac{p_{1-\alpha}(\gamma(\alpha))}{p_1(m_0)} \right) + \widetilde{\delta}(h) \left(G(\gamma) \frac{p_{1-\alpha}(\gamma(\alpha))}{p_1(m_0)} \right) \right] d\mu(\gamma) \\
& = \int_{P_{m_0}(M)} F(\gamma) \left(-D_h G(\gamma) + \widetilde{\delta}(h) G(\gamma) \right) \frac{p_{1-\alpha}(\gamma(\alpha))}{p_1(m_0)} d\mu(\gamma) \\
& \quad - \int_{P_{m_0}(M)} F(\gamma) G(\gamma) D_h \left(\frac{p_{1-\alpha}(\gamma(\alpha))}{p_1(m_0)} \right) d\mu \\
& = \int_{P_{m_0}(M)} F(\gamma) \left(-D_h G(\gamma) + \widetilde{\delta}(h) G(\gamma) \right) \frac{p_{1-\alpha}(\gamma(\alpha))}{p_1(m_0)} d\mu(\gamma) \\
& = \int_{L_{m_0}(M)} F(\gamma) \left(-D_h G(\gamma) + \widetilde{\delta}(h) G(\gamma) \right) d\nu(\gamma).
\end{aligned}$$

Hence Theorem 4.2 holds for $h \in H_1$.

Next, for any $h \in H_0$, since H_1 is dense in H_0 , there exist $h_n \in H_1$ such that $\|h_n - h\|_{H_0} \rightarrow 0$. By (4.2) in Theorem 4.1, and using the fact that $\widetilde{\delta}(h_n) - \widetilde{\delta}(h) = \widetilde{\delta}(h_n - h)$, for any $p > 1$, we have $\|\widetilde{\delta}(h_n) - \widetilde{\delta}(h)\|_{L^p(\nu)} \leq C \|h_n - h\|_{H_0}$. In particular, $\lim_{n \rightarrow \infty} \|\widetilde{\delta}(h_n) - \widetilde{\delta}(h)\|_{L^2(\nu)} = 0$. Hence

$$\langle F, \widetilde{\delta}(h_n) G \rangle_{L^2(\nu)} \rightarrow \langle F, \widetilde{\delta}(h) G \rangle_{L^2(\nu)}, \quad n \rightarrow \infty.$$

On the other hand, by the definition of D_h and the Lebesgue dominated convergence theorem, we have

$$\begin{aligned}
\langle D_{h_n} F, G \rangle_{L^2(\nu)} & \longrightarrow \langle D_h F, G \rangle_{L^2(\nu)}, \quad n \rightarrow \infty, \\
\langle D_{h_n} G, F \rangle_{L^2(\nu)} & \longrightarrow \langle D_h G, F \rangle_{L^2(\nu)}, \quad n \rightarrow \infty.
\end{aligned}$$

Note that $\langle D_{h_n} F, G \rangle_{L^2(\nu)} + \langle D_{h_n} G, F \rangle_{L^2(\nu)} = \langle F, \widetilde{\delta}(h_n) G \rangle_{L^2(\nu)}$. Letting $n \rightarrow \infty$, we prove that Theorem 4.2 holds for all $h \in H_0$. \square

Remark 4.1 *Combining Theorem 4.1 and Theorem 4.2 with Proposition 4.1 in Hsu [15] concerning the $L^1(\nu)$ -convergence of $\delta_s(h)$ to $\delta(h)$, we conclude that $\widetilde{\delta}(h)$ is nothing else than the divergence function defined in Driver [7] (in the case where*

$h \in \mathcal{C}^1 \cap H_0$), Enchev-Stroock [9] and Hsu [15] [16]. In view of this and to simplify the notation, as we have used in the statement of Theorem 1.1, in the rest part of this paper, we will use the notation $\delta(h)$ in stead of $\delta(\widetilde{h})$.

As a consequence of Theorem 4.2 and the $L^p(\nu)$ -integrability of the divergence functional $\delta(h)$ (see Theorem 4.1), by a standard argument as used in Fang-Malliavin [10] or Hsu [15] [16], we have the following result which allows us to introduce the first order Sobolev spaces over the loop space.

Theorem 4.3 *For all $p > 1$, D is closable from $L^p(L_{m_0}(M), \nu)$ to $L^p(L_{m_0}, H, \nu)$.*

5 Exponential integrability of divergence functionals

In this section we prove a Fernique type exponential integrability theorem for the divergence functional on the loop space.

Theorem 5.1 *For all*

$$\lambda < \lambda_0 = \frac{1}{(2 + \|Ric\|_\infty)\|h\|_H},$$

we have

$$E_\nu [\exp(\lambda|\delta(h)|^2)] < +\infty, \quad (5.1)$$

or, equivalently,

$$\lim_{t \rightarrow \infty} \frac{1}{t^2} \log \nu(\{\gamma \in L_{m_0}(M) : |\delta(h)| > t\}) = \frac{1}{(2 + \|Ric\|_\infty)\|h\|_H},$$

where $\|Ric\|_\infty$ denotes the upper bound of the Ricci curvature.

To prove Theorem 5.1, we shall use the following lemma which provides us with a very useful tool to study the exponential integrability of some functionals with the pinned Wiener measure on loop spaces.

Lemma 5.2 *Let $f \in W^{n, n^{2p}}(X)$, $n \in \mathbb{N}$, $p > 1$. Then*

$$\left\| e^{f^2} \right\|_{W^{n, p}(X)} \leq C(n) \left\| e^{(1+\epsilon)f^2} \right\|_{W^{n, \frac{n^{2p\epsilon}}{1+\epsilon}}(X)}^{\frac{1}{1+\epsilon}} \|f\|_{W^{n, \frac{n^{2p\epsilon}}{1+\epsilon}}(X)}^n. \quad (5.2)$$

Let $F \in D^{\infty, \infty}(P_{m_0}(M))$. Then there exist a constant $C > 0$ and a pair $(n, p) \in \mathbb{N} \times (1, +\infty)$ such that

$$E_\nu \left[e^{\widetilde{F}^2} \right] \leq C \left\| e^{(1+\epsilon)F^2} \right\|_{L^p(\mu)}^{\frac{1}{1+\epsilon}} \|F \circ I\|_{W^{n, \frac{n^{2p\epsilon}}{1+\epsilon}}(X)}^n, \quad (5.3)$$

where \widetilde{F} denotes any ∞ -quasi-continuous modification of F . Moreover, we can take $n = 2\lfloor \frac{d}{2} \rfloor + 2$ and $p > 1$ which can be arbitrary close to 1.

Proof of Lemma 5.2. By the chain rule, we have

$$\nabla^n e^{f^2} = e^{f^2} \sum_{r_1+\dots+r_n=n} c_{r_1,\dots,r_n} \nabla^{r_1} f^2 \otimes \dots \otimes \nabla^{r_n} f^2,$$

where c_{r_1,\dots,r_n} are some combinatory constants which can be given explicitly, and the summation is taken over all $\{(r_1, \dots, r_n) \in \mathbb{N}^n : 0 \leq r_1, \dots, r_n \leq n, r_1 + \dots + r_n = n\}$. Using the Hölder inequality, for any $\epsilon > 0$, we have

$$\left\| \|\nabla^n e^{f^2}\|_{H^{\otimes n}} \right\|_p \leq C(n) \|e^{f^2}\|_{p(1+\epsilon)} \sum_{r_1+\dots+r_n=n} \|\nabla^{r_1} f^2\|_{\frac{np\epsilon}{1+\epsilon}} \dots \|\nabla^{r_n} f^2\|_{\frac{np\epsilon}{1+\epsilon}}.$$

Moreover, for any $r \in \mathbb{N}$,

$$\nabla^r f^2 = \sum_{j_1+\dots+j_r=r} C_{j_1,\dots,j_r} \nabla^{j_1} f \otimes \dots \otimes \nabla^{j_r} f.$$

Hence

$$\begin{aligned} \|\nabla^r f^2\|_p &\leq C(r) \sum_{j_1+\dots+j_r=r} \|\nabla^{j_1} f\|_{rp} \dots \|\nabla^{j_r} f\|_{rp} \\ &\leq C(r) \|f\|_{W^{r,rp}(X)}^r. \end{aligned}$$

Thus,

$$\begin{aligned} \left\| \|\nabla^n e^{f^2}\|_{H^{\otimes n}} \right\|_p &\leq C(n) \|e^{(1+\epsilon)f^2}\|_p^{\frac{1}{1+\epsilon}} \sum_{r_1+\dots+r_n=n} \|f\|_{W^{r_1, \frac{r_1 np\epsilon}{1+\epsilon}}(X)}^{r_1} \dots \|f\|_{W^{r_n, \frac{r_n np\epsilon}{1+\epsilon}}(X)}^{r_n} \\ &\leq C(n) \|e^{(1+\epsilon)f^2}\|_p^{\frac{1}{1+\epsilon}} \|f\|_{W^{n, \frac{n^2 p\epsilon}{1+\epsilon}}(X)}^n. \end{aligned}$$

The inequality (5.2) follows. Combining (5.2), Theorem 2.1 with Remark 2.2, we prove the inequality (5.3). \square

Proposition 5.3 *For all*

$$\lambda < \lambda_0 = \frac{1}{(2 + \|Ric\|_\infty) \|h\|_H},$$

we have

$$E_\mu [\exp(\lambda |\delta(h)|^2)] < +\infty.$$

Proof. By random time changing, there exists a Brownian motion $\{B_t, t \in [0, \infty)\}$ which is adapted to the standard Brownian filtration $\mathcal{F}_s = \sigma(x(s), s \in [0, t])$ (here we allow $t \in [0, \infty)$) such that

$$\delta(h) \circ I(x) = B_T,$$

where

$$T = \left[\int_0^1 |\dot{h}(\tau) + \frac{1}{2} Ric_{r_x(\tau)} h(\tau)|^2 d\tau \right]^{\frac{1}{2}} \leq (1 + \frac{\|Ric\|_\infty}{2}) \|h\|_H.$$

By the refinement version of the well-known Fernique lemma ([11], see also Theorem 3.3 in [19]), we have

$$\begin{aligned} E_\mu [exp(\lambda \delta(h))] &= E [exp(\lambda \|B_T\|^2)] \\ &\leq E \left[exp \left(\lambda \sup_{s \in [0, (2 + \|Ric\|_\infty) \|h\|_H / 2]} \|B_t\|^2 \right) \right] \\ &< +\infty, \end{aligned}$$

provided that

$$\lambda(2 + \|Ric\|_\infty) \frac{\|h\|_H}{2} < \frac{1}{2},$$

i.e.,

$$\lambda < \lambda_0 = \frac{1}{(2 + \|Ric\|_\infty) \|h\|_H}.$$

□

Proof of Theorem 5.1. Applying Lemma 5.2 to $F = \delta(h)$, for any $p > 1$ and $\epsilon > 0$, there exists a constant $C > 0$ such that:

$$E_\nu [exp(\lambda |\delta(h)|^2)] \leq C \left\| exp(\lambda(1 + \epsilon) |\widehat{\delta(h)}|^2) \right\|_{W^{n,p}(X)}^{\frac{1}{1+\epsilon}} \left\| \widehat{\delta(h)} \right\|_{W^{n, \frac{n^2 p \epsilon}{1+\epsilon}}(X)}^n,$$

where $n = 2\lfloor \frac{d}{2} \rfloor + 2$. By Proposition 5.3 we have

$$\left\| e^{\lambda(1+\epsilon)p |\widehat{\delta(h)}|^2} \right\|_{W^{n,p}(X)} < +\infty,$$

provided that

$$\lambda < \frac{1}{p(1 + \epsilon)(2 + \|Ric\|_\infty) \|h\|_H}.$$

On the other hand, Theorem 3.3 says

$$\left\| \widehat{\delta(h)} \right\|_{W^{n, \frac{n^2 p \epsilon}{1+\epsilon}}(X)} < C \|h\|_H.$$

Thus, for all $\lambda < [p(1 + \epsilon)(2 + \|Ric\|_\infty) \|h\|_H]^{-1}$, we have $E_\nu [exp(\lambda |\delta(h)|^2)] < +\infty$. Since we can chose p arbitrary close to 1 and ϵ arbitrary close to 0, we deduce the desired inequality (5.1) for all $\lambda < \lambda_0 = [(2 + \|Ric\|_\infty) \|h\|_H]^{-1}$. □

As a consequence of Theorem 5.1 we have the following result.

Theorem 5.4 For any $\lambda > 0$ and any $h \in H_0$, we have

$$E_\nu [\exp(\lambda|\delta(h)|)] < +\infty.$$

Proof. By Theorem 5.1 we have

$$\lim_{t \rightarrow \infty} \frac{1}{t^2} \log \nu (\{\gamma \in L_{m_0}(M) : |\delta(h)| > t\}) = -\frac{1}{(2 + \|Ric\|_\infty)\|h\|_H}.$$

This yields that

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \nu (\{\gamma \in L_{m_0}(M) : |\delta(h)| > t\}) = -\infty.$$

Theorem 5.4 follows. □

6 Smoothness of Driver's flow on path spaces

Recall that by [6] [8] [14], for all $h \in H$, the vector field D_h generates a global flow $\{\Phi_t, t \in \mathbb{R}\}$, the so-called Driver flow on $P_{m_0}(M)$, such that for μ -a.s. $\gamma \in P_{m_0}(M)$,

$$\begin{aligned} \dot{\Phi}_t(\gamma) &= D_h(\Phi_t(\gamma)), \\ \Phi_0(\gamma) &= \gamma. \end{aligned} \tag{6.1}$$

In the case where $h \in \mathcal{C}^1 \cap H_0$ is a Lipschitz Cameron-Martin vector, Driver[7] constructed the flow generated by D_h on the loop space $L_{m_0}(M)$ by using the technique of enlargement of filtration. In [9], Enchev-Stroock gave another approach to construct the flow corresponding to D_h for all $h \in H_0$. As we have pointed out before, their approaches relied on the gradient estimate and the Hessian estimate of the logarithmic of the heat kernel on compact Riemannian manifold. For details, see Section 9.

In this section, without using any heat kernel estimate, we construct the Driver flow of D_h on the loop space for all $h \in H_0$ through an ∞ -quasi-continuous modification of the corresponding flow on the path space. To this end, we shall first prove that for any $h \in H_0$ the Driver flow generated by D_h is a smooth transform on the path space $P_{m_0}^{2m, \alpha}(M)$ in the sense of the Malliavin calculus, where for any $m \in \mathbb{N}$, $m \geq 2$, $\alpha \in (\frac{1}{2m}, \frac{1}{2})$,

$$P_{m_0}^{2m, \alpha}(M) = \left\{ \gamma \in P_{m_0}(M) : \int_0^1 \int_0^1 \frac{d(\gamma(t), \gamma(s))^{2m}}{|t-s|^{1+2m\alpha}} dt ds < \infty \right\}.$$

By [21] and the reference therein, the Wiener measure ν is supported on $P_{m_0}^{2m, \alpha}(M)$ and the Sobolev spaces theory on $P_{m_0}^{2m, \alpha}(M)$ is as the same as (i.e., quasi-homeomorphic

to) the one on $P_{m_0}(M)$. Moreover, $P_{m_0}^{2m,\alpha}(M)$ is a M -type 2 Banach manifold modelled on $X^{2m,\alpha}$ whose norm $\|\cdot\|_{2m,\alpha}$ is smooth in $X^{2m,\alpha} \setminus \{0\}$, where

$$X^{2m,\alpha} := \left\{ x \in X : \int_0^1 \int_0^1 \frac{\|x(t) - x(s)\|_{\mathbb{R}^d}^{2m}}{|t-s|^{1+2m\alpha}} dt ds < \infty \right\},$$

on which we consider the fractional Hölder norm $\|\cdot\|_{2m,\alpha}$ given by

$$\|x\|_{2m,\alpha} = \left(\int_0^1 \int_0^1 \frac{\|x(t) - x(s)\|_{\mathbb{R}^d}^{2m}}{|t-s|^{1+2m\alpha}} dt ds \right)^{\frac{1}{2m}}.$$

Theorem 6.1 *For any $h \in H$, Driver's flow $\Phi_t : P_{m_0}^{2m,\alpha}(M) \rightarrow P_{m_0}^{2m,\alpha}(M)$ is a smooth mapping in the sense of the Malliavin calculus, i.e.,*

$$\Phi_t \in D^{\infty,\infty}(P_{m_0}^{2m,\alpha}(M), P_{m_0}^{2m,\alpha}(M)).$$

To prove this theorem, let us first introduce the set $\mathcal{SM}(\mathbb{R}^d)$ of all \mathbb{R}^d -valued semimartingales with the Doob-Meyer decomposition $\xi(s) = \int_0^s O(r)dr + \int_0^s A(r)dr$, where $\{(O(s), A(s)), s \in [0, 1]\}$ is an adapted $M(d, d) \times \mathbb{R}^d$ -valued process such that

$\|\xi\|^2 := E \left[\sup_{s \in [0,1]} \|O(s)\|_{M(d,d)}^2 \right] + E \left[\int_0^1 \|A(s)\|_{\mathbb{R}^d}^2 ds \right]$ is finite. By definition, we have

$$\begin{aligned} \|\|\xi(\cdot)\|_{2m,\alpha}\|_p &= \left\| \left\| \int_0^\cdot O(r)dx(r) + \int_0^\cdot A(r)dr \right\|_{2m,\alpha} \right\|_p \\ &\leq \left\| \left\| \int_0^\cdot O(r)dx(r) \right\|_{2m,\alpha} \right\|_p + \left\| \left\| \int_0^\cdot A(r)dr \right\|_{2m,\alpha} \right\|_p. \end{aligned}$$

Using the Burkholder-Davies-Gundy inequality we have

$$\begin{aligned} E \left[\left\| \int_0^\cdot O(r)dx(r) \right\|_{2m,\alpha}^p \right] &\leq C_p E \left[\left\| \int_0^\cdot |O(r)|^2 dr \right\|_{2m,\alpha}^{\frac{p}{2}} \right] \\ &\leq C_p E \left[\sup_{r \in [0,1]} \|O(r)\|^{2 \cdot \frac{p}{2}} \left\| \int_0^\cdot dr \right\|_{2m,\alpha}^{\frac{p}{2}} \right] \\ &\leq C(p, m, \alpha) E \left[\sup_{r \in [0,1]} \|O(r)\|^p \right]. \end{aligned}$$

Using the Cauchy-Schwartz inequality we have

$$\left| \int_{s_1}^{s_2} A(r)dr \right|^{2m} \leq \left[\int_0^1 |A(r)|^2 dr \right]^m |s_1 - s_2|^m.$$

Thus

$$\begin{aligned} \int_0^1 \int_0^1 \frac{|\int_{s_1}^{s_2} A(r) dr|^{2m}}{|s_1 - s_2|^{1+2m\alpha}} ds_1 ds_2 &\leq \left[\int_0^1 |A(r)|^2 dr \right]^m \int_0^1 \int_0^1 \frac{ds_1 ds_2}{|s_1 - s_2|^{1+2m\alpha-m}} \\ &\leq C(m, \alpha) \left[\int_0^1 |A(r)|^2 dr \right]^m. \end{aligned}$$

This yields that

$$\begin{aligned} \left\| \left\| \int_0^1 A(r) dr \right\|_{2m, \alpha} \right\|_p &\leq [C(m, \alpha)]^{\frac{1}{2m}} \left\{ E \left[\int_0^1 |A(r)|^2 dr \right]^{pm \times \frac{1}{2m}} \right\}^{\frac{1}{p}} \\ &\leq C(m, \alpha) \left\{ E \left[\int_0^1 |A(r)|^2 dr \right]^{\frac{p}{2}} \right\}^{\frac{1}{p}}. \end{aligned}$$

Therefore

$$\| \|\xi\|_{2m, \alpha} \|_p \leq C(m, p, \alpha) \left\{ E \left[\sup_{r \in [0, 1]} |O(r)|^p \right] \right\}^{\frac{1}{p}} + C(m, p, \alpha) \left\{ E \left[\int_0^1 |A(r)|^2 dr \right]^{\frac{p}{2}} \right\}^{\frac{1}{p}}.$$

Hence for $m \geq 2$, $\alpha \in (\frac{1}{2m}, \frac{1}{2})$ and $p \geq 1$, we have

$$E \left[\|\xi\|_{2m, \alpha}^p \right] \leq C(m, p, \alpha) E \left[\sup_{r \in [0, 1]} \|O(r)\|^p \right] + C(m, p, \alpha) E \left[\int_0^1 |A(r)|^2 dr \right]^{\frac{p}{2}}.$$

For any $p \geq 1$, define the norm $\|\cdot\|_p$ on $\mathcal{SM}(\mathbb{R}^d)$ as

$$\|\xi\|_p^p := E \left[\sup_{r \in [0, 1]} \|O(r)\|^p \right] + E \left[\int_0^1 |A(r)|^2 dr \right]^{\frac{p}{2}}.$$

Then

$$E \left[\|\xi\|_{2m, \alpha}^p \right] \leq C(m, p, \alpha) E \left[\|\xi\|_p^p \right]. \quad (6.2)$$

Proof of Theorem 6.1. Set $\xi_t = I^{-1} \circ \Phi_t \circ I$. By [21], we have $I \in D^{\infty, \infty}(X^{2m, \alpha}, P_{m_0}^{2m, \alpha}(M))$ and $I^{-1} \in D^{\infty, \infty}(P_{m_0}^{2m, \alpha}(M), X^{2m, \alpha})$. Hence it remains to prove $\xi_t \in W^{\infty, \infty}(X^{2m, \alpha}, X^{2m, \alpha})$. By [6] [14], ξ_t satisfies the following ODE on $X^{2m, \alpha}$, where the stochastic integral is taken in the sense of Itô,

$$\begin{aligned} \frac{\partial}{\partial t} \xi_t(s) &= h(s) + \frac{1}{2} \int_0^s Ric_{U_t(\tau)}(h(\tau)) d\tau + \int_0^s q_h(t, \tau) d\xi_t(\tau) \\ \xi_0(x) &= x. \end{aligned} \quad (6.3)$$

Here $U_t(s)$ is the stochastic parallel transport along $\gamma_t(s) = \Phi_t(\gamma)(s)$, $q_h(t, s)$ is given by the following Stratonovich stochastic integral

$$q_h(t, s) = \int_0^s \Omega_{U_t(\tau)}(h(\tau), \circ d\xi_t(\tau)).$$

Using the Picard iteration and by a similar argument as used in Driver [6] and Hsu[15], we can prove that for any $k \in K$, $D_k \xi_t(s)$ exists for all $s \in [0, 1]$ and all $t \in \mathbb{R}$. Moreover, $D_k \xi_t$ satisfies the following equation

$$\begin{aligned} \frac{\partial}{\partial t} D_k \xi_t(s) &= \int_0^s q_h(t, \tau) dD_k \xi_t(\tau) + \frac{1}{2} \int_0^s D_k Ric_{U_t(\tau)}(h(\tau)) d\tau + \int_0^s D_k q_h(t, \tau) d\xi_t(\tau), \\ D_k \xi_0(s) &= k(s). \end{aligned} \tag{6.4}$$

For any $T > 0$, $p > 1$, we can prove

$$\sup_{t \in [-T, T]} E \left[\left\| \|D\xi_t(\cdot)\|_H \right\|_{2m, \alpha}^p \right] < \infty. \tag{6.5}$$

Indeed, let $\{\xi_t^n(s), s \in [0, 1], t \in [-T, T]\}$ be given as in [6] and [14]. Let $(O_t^n(s), A_t^n(s))$ be the Doob-Meyer decomposition of ξ_t^n . Let $\gamma_t^n = I(\xi_t^n)$, $U_t^n = U(\gamma_t^n)$ and $q_h^n(t, s) = \int_0^s \Omega_{U_t^n(r)}(h(r), \circ d\xi_t^n(r))$, $s \in [0, 1]$, $t \in [-T, T]$. Then for any fixed $s \in [0, 1]$ and $t \in [-T, T]$, it is easy to see that $(O_t^n(s), A_t^n(s)) \in W^{\infty, \infty}(X, O(d) \times \mathbb{R}^d)$. Similarly to [14], we can easily prove that

$$\sup_{t \in [-T, T]} E \left[\left\| \xi_t^n - \xi_t^{n-1} \right\|^{2p} \right] \leq \frac{(CT)^n}{n!}, \tag{6.6}$$

where

$$E \left[\left\| \xi_t^n - \xi_t^{n-1} \right\|^p \right] := E \left[\sup_{s \in [0, 1]} \|O_t^n - O_t^{n-1}\|_{M(d, d)}^p \right] + E \left[\int_0^1 \|A_t^n(s) - A_t^{n-1}(s)\|^2 ds \right]^{\frac{p}{2}}.$$

Moreover, for all $k \in H$ we can easily show that

$$\begin{aligned} D_k O_t^n(s) &= O_t^n(s) \int_0^t [O_u^n(s)]^{-1} D_k q_h^{n-1}(u, s) O_u^n(s) du, \\ D_k A_t^n(s) &= O_t^n(s) \int_0^t [O_u^n(s)]^{-1} D_k q_h^{n-1}(u, s) A_u^n(s) du \\ &\quad + \frac{1}{2} O_t^n(s) \int_0^t [O_u^n(s)]^{-1} D_k Ric_{U_u^{n-1}(s)}(h(s)) du. \end{aligned}$$

Putting $k \in H$ such that $\dot{k} = 1_{[\tau, 1]} e_\alpha$ into the above formulas, we obtain the explicit expressions of the Malliavin derivatives $D_\tau^\alpha O_t^n(s)$ and $D_\tau^\alpha A_t^n(s)$. Set

$$E \left[\left\| \|D_\tau^\alpha \xi_t^n\| \right\|^p \right] := E \left[\|O_t^n\|_{M(d, d)}^p \right] + E \left[\sup_{s \in [0, 1]} \|D_\tau^\alpha O_t^n(s)\|_{M(d, d)}^p \right] + E \left[\int_0^1 \|D_\tau^\alpha A_t^n(s)\|_{\mathbb{R}^d}^2 ds \right]^{\frac{p}{2}}.$$

By standard argument and the Burkholder-Davies-Gundy inequality, and using the fact that $O_t^n(s) \in O(d)$ and $\sup_{n \in \mathbb{N}} \|A_t^n(s)\| \leq C(1 + |\dot{h}(s)|)$ (see [14]), it is straightforward to prove that

$$\begin{aligned}
E \left[\sup_{s \in [0,1]} \|D_\tau^\alpha U_t^n(s)\|^p \right] &\leq cE [\|D_\tau^\alpha \xi_t^n\|^p], \quad E \left[\sup_{s \in [0,1]} \|D_\tau^\alpha q_h^n(t, s)\|^p \right] \leq cE [\|D_\tau^\alpha \xi_t^n\|^p], \\
E \left[\sup_{s \in [0,1]} \|D_\tau^\alpha U_t^n(s) - D_\tau^\alpha U_t^{n-1}(s)\|^p \right] &\leq cE \left[\sup_{s \in [0,1]} \|D_\tau^\alpha \xi_t^n(s) - D_\tau^\alpha \xi_t^{n-1}(s)\|^p \right] \\
&\quad + c \left\{ E \left[\sup_{s \in [0,1]} \|D_\tau^\alpha U_t^n(s)\|^{2p} \right] \right\}^{\frac{1}{2}} \left\{ E [\|\xi_t^n - \xi_t^{n-1}\|^{2p}] \right\}^{\frac{1}{2}}, \\
E \left[\sup_{s \in [0,1]} \|D_\tau^\alpha q_h^n(t, s) - D_\tau^\alpha q_h^{n-1}(t, s)\|^p \right] &\leq cE [\|D_\tau^\alpha \xi_t^n - D_\tau^\alpha \xi_t^{n-1}\|^p] \\
&\quad + c \left\{ E [\|D_\tau^\alpha \xi_t^{n-1}\|^{2p}] \right\}^{\frac{1}{2}} \left\{ E [\|\xi_t^{n-1} - \xi_t^{n-2}\|^{2p}] \right\}^{\frac{1}{2}}.
\end{aligned}$$

From the above inequalities we can deduce that

$$\sup_{n \in \mathbb{N}} \sup_{t \in [-T, T]} \sup_{\tau \in [0, 1]} E [\|D_\tau^\alpha \xi_t^n\|^p] < c_1 e^{c_2 T}, \quad (6.7)$$

$$E \left[\sup_{s \in [0,1]} \|D_\tau^\alpha O_t^n - D_\tau^\alpha O_t^{n-1}(s)\|^p \right] \leq R_{\tau, \alpha}^1(T, n) + c \int_0^t \|D_\tau^\alpha \xi_u^n - D_\tau^\alpha \xi_u^{n-1}\|^p du, \quad (6.8)$$

$$E \left[\left| \int_0^1 \|D_\tau^\alpha A_t^n - D_\tau^\alpha A_t^{n-1}(s)\|^2 ds \right|^{\frac{p}{2}} \right] \leq R_{\tau, \alpha}^2(T, n) + c \int_0^t \|D_\tau^\alpha \xi_u^n - D_\tau^\alpha \xi_u^{n-1}\|^p du, \quad (6.9)$$

where for $i = 1, 2$, there exists a constant $C(T, p)$ such that

$$R_{\tau, \alpha}^i(T, n) \leq C(T, p) \left\{ \sup_{t \in [-T, T]} \sup_{\tau \in [0, 1]} E [\|D_\tau^\alpha \xi_t^n\|^{2p}] \right\}^{1/2} \left\{ \sup_{t \in [-T, T]} E [\|\xi_t^n - \xi_t^{n-1}\|^{2p}] \right\}^{1/2}. \quad (6.10)$$

From (6.6), (6.7) and (6.10), $R_{\tau, \alpha}(T, n) = R_{\tau, \alpha}^1(T, n) + R_{\tau, \alpha}^2(T, n)$ tends to zero as n tends to infinity. On the other hand, from (6.8), (6.9) and the Gronwall inequality we have

$$\sup_{t \in [-T, T]} \sup_{\tau \in [0, 1]} E [\|D_\tau^\alpha \xi_t^n - D_\tau^\alpha \xi_t^{n-1}\|^p] \leq R_{\tau, \alpha}(T, n) e^{cT}. \quad (6.11)$$

This implies that $\{\xi_t^n, t \in [-T, T]\}$ converges uniformly in $W^{1, \infty}(X, (\mathcal{SM}(\mathbb{R}^d), \|\cdot\|_p))$ and hence by (6.2) it converges uniformly in $W^{1, \infty}(X, (X^{2m, \alpha}, \|\cdot\|_{2m, \alpha}))$. Moreover, we deduce Eq. (6.4) (respectively, the inequality (6.5)) from the corresponding

equation for $D_k \xi_t^n$ (respectively, the inequality (6.7)). In general, by induction and repeating the same argument as above, we can prove that for all $k_1, \dots, k_r \in H$, $D_{k_1 \dots k_r} \xi_t(s)$ exists for all $s \in [0, 1]$ and all $t \in \mathbb{R}$. Moreover, for any $T > 0$, $p > 1$, using the Burkholder-Davies-Gundy inequality we have

$$\sup_{t \in [-T, T]} E \left[\left\| \|D^r \xi_t(\cdot)\|_{H^{\otimes r}} \right\|_{2m, \alpha}^p \right] < \infty. \quad (6.12)$$

This finishes the proof of Theorem 6.1 concerning the smoothness of the Driver flow on the path space $P_{m_0}^{2m, \alpha}(M)$. To save the length of the paper, we omit the details of the proofs of the four inequalities listed before (6.7). The reader who is interested in the details of the proof is referred to C. Cross [3] (for the case where $h \in \mathcal{C}^1 \cap H$ is a Lipschitz Cameron-Martin vector) and Léandre [18] as well as Li [22] where the author proved that the Driver flow Φ_t is a smooth transform on the path space $P_{m_0}(M)$. \square

We will make use of the following Kolmogorov criterion for ∞ -quasi-continuous modification of a family of an M -type 2 Banach space E -valued functionals. When $E = \mathbb{R}$, it is due to Ren [28]. See also [26].

Theorem 6.2 *Let $\{X(t), t \in [-T, T]\}$ be a family of an M -type 2 Banach space E -valued functionals. Suppose that for all $p \geq 2$, $r \in \mathbb{N}$ there exist constants $c, \epsilon > 0$ and an even number β such that*

- (1) $X(t) \in W^{r, p}(X, E)$.
- (2) For all $(s, t) \in [-T, T] \times [-T, T]$, $\|X(t) - X(s)\|_E^\beta \in W^{r, p}(X)$.
- (3) For all $(s, t) \in [-T, T] \times [-T, T]$, we have

$$\left\| \|X(t) - X(s)\|_E^\beta \right\|_{W^{r, p}(X)} \leq c|t - s|^{1+\epsilon}.$$

Then there exists a version of the process $\{X(t), t \in [-T, T]\}$ which is ∞ -quasi-continuous for each $t \in [-T, T]$ and which has continuous paths.

Proof. Since E is an M -type 2 Banach space, the norm $\phi(x) = \|x\|_E$ is smooth in $E \setminus \{0\}$ and for all $k \in \mathbb{N}$ there exists M_k such that $\sup_{\|x\|_E=1} \|\nabla^k \phi(x)\| \leq M_k < \infty$.

Thus, the Tchebycheff type inequality of (r, p) -capacity for E -valued functionals holds, see e.g. [26]. Hence, for any given $r \in \mathbb{N}$, $p > 1$, $\epsilon > 0$ we have

$$C_{r, p}(\{x \in X : \|X(t) - X(s)\|_E > \epsilon\}) \leq \frac{1}{\epsilon^\beta} \left\| \|X(t) - X(s)\|_E^\beta \right\|_{r, p}.$$

Therefore, Theorem 6.2 can be proved by the same argument as used in the proof of Theorem 3.1 in [28]. \square

Theorem 6.3 For all $p \geq 2$ and $r \in \mathbb{N}$, there exist constants $c, \epsilon > 0$ such that

- (1) $\xi_t \in W^{r,p}(X^{2m,\alpha}, X^{2m,\alpha})$.
- (2) $\|\xi_t - \xi_s\|_{2m,\alpha}^\beta \in W^{r,p}(X^{2m,\alpha})$ for all $(s, t) \in [-T, T] \times [-T, T]$.
- (3) For all $(s, t) \in [-T, T] \times [-T, T]$, we have

$$\left\| \|\xi_t(\cdot) - \xi_s(\cdot)\|_{2m,\alpha}^\beta \right\|_{r,p} \leq c|t - s|^{1+\epsilon}. \quad (6.13)$$

Proof. By Theorem 6.1, for all $t \in [-T, T]$ we have $\xi_t \in W^{r,p}(X^{2m,\alpha}, X^{2m,\alpha})$. By the chain rule, for all even number $\beta > 0$ and $(s, t) \in [-T, T] \times [-T, T]$, we can prove $\|\xi_t - \xi_s\|_{2m,\alpha}^\beta \in W^{r,p}(X^{2m,\alpha})$. By Lemma 4.1 in [28], for all $(s, t) \in [-T, T] \times [-T, T]$, we have

$$\begin{aligned} \left\| \|\xi_t(\cdot) - \xi_s(\cdot)\|_{2m,\alpha}^n \right\|_{2r,p} &\leq C(n, p, r) \left\| \|\xi_t(\cdot) - \xi_s(\cdot)\|_{2m,\alpha} \right\|_{2r,4r^2p}^{2r} \\ &\quad \times \max_{0 \leq k \leq n} \left[E \|\xi_t(\cdot) - \xi_s(\cdot)\|_{2m,\alpha}^{(n-k)2rp} \right]^{\frac{1}{2rp}}. \end{aligned} \quad (6.14)$$

On the other hand, using Eq. (6.3), Eq. (6.4) and the inequalities (6.5) and (6.12), by the Hölder inequality and the Burkholder-Davies-Gundy inequality, we can verify that for all $p \geq 1$ and $r \in \mathbb{N}$,

$$\left\| \|\xi_t(\cdot) - \xi_s(\cdot)\|_{2m,\alpha} \right\|_{2p}^{2p} \leq C(p, T)|t - s|^p. \quad (6.15)$$

$$\left\| \|\xi_t(\cdot) - \xi_s(\cdot)\|_{2m,\alpha} \right\|_{r,2p}^{2p} \leq C(p, T)|t - s|^p. \quad (6.16)$$

From (6.14), (6.15) and (6.16), we deduce that for $n(p, r) = n$ large enough and for some constant $\epsilon > 0$ we have

$$\left\| \|\xi_t(\cdot) - \xi_s(\cdot)\|_{2m,\alpha}^{n(p,r)} \right\|_{2r,p} \leq c|t - s|^{1+\epsilon}.$$

The proof of Theorem 6.3 is completed. \square

Combining Theorem 6.3 with the Kolmogorov criterion (Theorem 6.2), we have the following

Theorem 6.4 For all $T > 0$, there exists a version of the Driver flow $\{\Phi_t, t \in [-T, T]\}$ on the path space $P_{m_0}^{2m,\alpha}(M)$ which is ∞ -quasi-continuous for each $t \in [-T, T]$ and which has continuous trajectory on $t \in [-T, T]$.

Proof of Theorem 1.4. By Theorem 6.3, for any $T > 0$, there exists a slim subset S_T of the path space $P_{m_0}^{2m,\alpha}(M)$ such that an ∞ -quasi-continuous modification (denoted by $\{\tilde{\Phi}_t, t \in [-T, T]\}$) of $\{\Phi_t, t \in [-T, T]\}$ can be well-defined for all $\gamma \in P_{m_0}^{2m,\alpha}(M) \setminus S_T$. Taking $T_n = 2^n$ we deduce that there exists a common slim set $S_\infty = \cup_{n \in \mathbb{N}} S_{T_n}$ such that $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$ can be well-defined for all $\gamma \in P_{m_0}^{2m,\alpha}(M) \setminus S_\infty$. Thus, $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$ can be ν -a.s. well defined on the loop space $L_{m_0}^{2m,\alpha}(M) = L_{m_0}(M) \cap P_{m_0}^{2m,\alpha}(M)$ and hence is ν -a.s. well defined on $L_{m_0}(M)$. \square

In the rest part of this paper, we fix such an ∞ -quasi-continuous version $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$ of $\{\Phi_t, t \in \mathbb{R}\}$. To end this section let us mention the following remark.

Remark 6.1 Note that the Driver flow $\{\Phi_t, t \in \mathbb{R}\}$ is $\mathcal{F}_s/\mathcal{F}_s$ measurable for all $s \in [0, 1]$, see [6] [8] [9]. Thus, the Kolmogorov criterion yields that $\{\tilde{\Phi}_t, t \in \mathbb{R}\}$ is again $\mathcal{F}_s/\mathcal{F}_s$ -measurable for all $s \in [0, 1]$. Moreover, using the same argument as used in the proof of the Sobolev norm and the capacity comparison theorems between the Wiener space and the path space via the Itô map, see [21], we can prove that for any $F \in W^{r,p}(X)$ and any subset $A \subset X$ and any $r \in \mathbb{N}$ and $p > 1$,

$$\alpha_1 \|F \circ \xi_t\|_{\frac{r}{2}, p-\epsilon} \leq \|F\|_{r,p} \leq \alpha_2 \|F \circ \xi_t\|_{2r,p+\epsilon}, \quad (6.17)$$

$$\alpha_1 C_{\frac{r}{2}, p-\epsilon}(\tilde{\xi}_t^{-1}(A)) \leq C_{r,p}(A) \leq \alpha_2 C_{2r, p+\epsilon}(\tilde{\xi}_t^{-1}(A)), \quad (6.18)$$

where $\|\cdot\|_{r,p}$ (respectively, $C_{r,p}$) denotes the (r, p) -Sobolev norm (respectively, (r, p) -capacity) on the Wiener space X , $\{\tilde{\xi}_t, t \in \mathbb{R}\}$ denotes any ∞ -quasi-continuous modification of the pull back of the Driver flow $\xi_t = I^{-1} \circ \Phi_t \circ I$, α_1 and α_2 are two constants which depend only on r, p and the upper bounds of the Riemannian curvature and the Ricci curvature as well as their higher order covariant derivatives. This yields that the flow property $\tilde{\xi}_s \circ \tilde{\xi}_t = \tilde{\xi}_{t+s}$ holds quasi-surely on X . Since $\tilde{\Phi}_t = \tilde{I} \circ \tilde{\xi}_t \circ \tilde{I}^{-1}$, we get the flow property $\tilde{\Phi}_s \circ \tilde{\Phi}_t = \tilde{\Phi}_{t+s}$ quasi-surely on $P_{m_0}(M)$. As explained in [21], the proof of the above inequalities (6.17) and (6.18) are based on the P. A. Meyer inequality on the Wiener space. However, we still do not know whether the P. A. Meyer inequality holds or not on the path space over a compact Riemannian manifold. Thus, we do not know whether or not the corresponding Sobolev norms (respectively, capacities) comparison inequalities hold on path spaces if one replace ξ_t by Φ_t , etc.

7 Cameron-Martin theorem on loop spaces

In this section we will first construct the Driver flow on the loop space through the corresponding flow on the path space. Combining this and Theorem 5.4 together with the Cruzeiro lemma we will give an alternative proof to the Cameron-Martin theorem on loop spaces established earlier by Driver[7] and Enchev-Stroock[9] by Doob's h -processes approach and the short time upper bound estimates of the gradient and the Hessian of logarithmic of the heat kernels.

Our first result in this section is the following

Theorem 7.1 Let $h \in H_0$. Then $\tilde{\Phi}_t(L_{m_0}(M)) \subset L_{m_0}(M)$. Moreover, for ν -a.s. $\gamma \in L_{m_0}(M)$, we have

$$\begin{aligned} \tilde{\Phi}_t(\gamma) &= D_h(\tilde{\Phi}_t(\gamma)), \\ \tilde{\Phi}_t(\gamma) &= \gamma. \end{aligned} \quad (7.1)$$

In view of Theorem 7.1, we regard $\tilde{\Phi}_t$ as the flow on $L_{m_0}(M)$ generated by D_h .

Proof. Since $\tilde{\Phi}_t = \Phi_t$ holds quasi-surely on $P_{m_0}(M)$, the flow equation (6.1) is verified. It remains to show that $\tilde{\Phi}_t(L_{m_0}(M) \subset L_{m_0}(M)$, i.e.,

$$\tilde{\Phi}_t(\gamma)(1) = m_0. \quad (7.2)$$

To this end, we use the same argument as in Driver [7]. Indeed, for ν -a.s. $\gamma \in L_{m_0}(M)$ and any $s < 1$, by the flow property $\tilde{\Phi}_t$ on $L_{m_0}(M)$ we have

$$\begin{aligned} d(\tilde{\Phi}_t(\gamma)(s), \tilde{\Phi}_0(\gamma)(s)) &\leq \int_0^t \left| \frac{d}{d\tau} \tilde{\Phi}_\tau(\gamma)(s) \right| d\tau \\ &= \int_0^t |U(\tilde{\Phi}_\tau(\gamma))(s)h(s)| d\tau. \end{aligned}$$

Now $U(\tilde{\Phi}_\tau(\gamma))(s)$ is isometry from \mathbb{R}^d to $T_{\tilde{\Phi}_\tau(\gamma)(s)}M$. Thus for ν -a.s. $\gamma \in L_{m_0}(M)$,

$$d(\tilde{\Phi}_t(\gamma)(s), \tilde{\Phi}_0(\gamma)(s)) \leq |h(s)|. \quad (7.3)$$

Let

$$\Sigma = \left\{ \gamma \in L_{m_0}(M) : d(\tilde{\Phi}_t(\gamma)(s), \tilde{\Phi}_0(\gamma)(s)) \leq |h(s)|, \quad \forall s < 1 \right\}.$$

Since both sides of the inequality in (7.3) are continuous, we have

$$\nu(\Sigma) = 1. \quad (7.4)$$

Taking $\gamma \in \Sigma$ and letting $s \rightarrow 1$, we have

$$\lim_{s \rightarrow 1} d(\tilde{\Phi}_t(\gamma)(s), \gamma(s)) \leq \lim_{s \rightarrow 1} |h(s)| = 0.$$

By the continuity of $s \rightarrow \tilde{\Phi}_t(s)$, we prove (7.2) for $\gamma \in \Sigma$ which differs from $L_{m_0}(M)$ up to a ν -negligible subset, cf. (7.4). \square

The following lemma is due to Ana Bela Cruzeiro [4] and is a very useful tool to study the quasi-invariance of a probability measure under the action of certain flows.

Lemma 7.2 (*Cruzeiro*) *Let $(\Omega, \mathcal{F}, \{\mathcal{F}_s\}, P)$ be a complete filtered probability space, $\Phi = \{\phi_t\}$ be a flow (i.e., a one-parameter group of measurable transformations) on Ω . Suppose that there exists the divergence $\text{div}(\Phi) \in L^1(\Omega, P)$ such that for all $f \in \mathcal{C} \subset L^\infty(\Omega, P)$, where \mathcal{C} is a dense subset of $L^\infty(\Omega, P)$, we have*

$$\left\{ \frac{d}{dt} E[f(\phi_t)] \right\} \Big|_{t=0} = E[f \text{div}(\Phi)]. \quad (7.5)$$

Moreover, assume that there exists a $\lambda > 0$ such that

$$E[e^{\lambda |\text{div}(\Phi)|}] < +\infty. \quad (7.6)$$

Then $(\phi_t)_*P$ is absolutely continuous with respect to P . Denote

$$K_t = \frac{d(\phi_t)_*P}{dP}.$$

Then

$$K_t = \exp \left(\int_0^t \operatorname{div}(\Phi)(\phi_{-s}) ds \right),$$

and for all $p > 1$ with the conjugate exponent q , i.e., $\frac{1}{p} + \frac{1}{q} = 1$, we have

$$\|K_t\|_{L^p}^p \leq E [e^{pqt|\operatorname{div}(\Phi)}].$$

Proof. See [4] [24]. □

Now we are ready to prove the Cameron-Martin theorem on the loop space.

Theorem 7.3 *For any $h \in H_o$, the pinned Wiener measure ν on $L_{m_0}(M)$ is quasi-invariant under Driver's flow $\tilde{\Phi}_t$. Let*

$$K_t = \frac{d(\tilde{\Phi}_t)_*\nu}{d\nu}.$$

Then

$$K_t(\gamma) = \exp \left(\int_0^t \delta(h)(\tilde{\Phi}_{-s}(\gamma)) ds \right), \quad \nu - \text{a.s. } \gamma \in L_{m_0}(M), \quad (7.7)$$

and for all $p > 1$ with the conjugate exponent q , i.e., $\frac{1}{p} + \frac{1}{q} = 1$, we have

$$\|K_t\|_{L^p(\nu)}^p \leq E [e^{pqt|\tilde{\delta}(h)}]. \quad (7.8)$$

Proof. Set $\Omega = L_{m_0}(M)$, $P = \nu$, $\mathcal{C} = \mathcal{FC}(L_{m_0}(M))$ (the collection of all cylindrical functionals on $L_{m_0}(M)$), ϕ_t be Driver's flow $\tilde{\Phi}_t$. Theorem 4.2 shows that the divergence $\operatorname{div}(\Phi)$ in (7.5) associated to the flow ϕ_t is just $\delta(h)$. By Theorem 5.4, the ν -exponential integrability (7.6) holds for $\delta(h)$. Hence Cruzeiro's lemma applies to $(L_{m_0}(M), \tilde{\Phi}_t, \nu)$. Thus, $(\tilde{\Phi}_t)_*\nu$ is absolutely continuous with respect to ν , i.e. the pinned Wiener measure ν is quasi-invariant on loop space $L_{m_0}(M)$ under the flow $\tilde{\Phi}_t$. Moreover, we obtain (7.7) and the L^p -inequality (7.8). □

According to Remark 1.1, our main results apply to the special case where $M = G$ is a compact connected Lie group equipped with an Ad -invariant metric and the left or the right Cartan connection. Theorem 1.5 recaptures the well-known result due to M. P. Malliavin and P. Malliavin [25] on the quasi-invariance of the pinned Wiener measure on the loop group. Indeed, our method is inspired by [25] where the authors initiated the so-called *localization method from paths to loops* based on quasi-sure analysis.

8 Stochastic anti-development of pinned Brownian motions

Since J. Bismut [2], it has been well-known that the stochastic anti-development of the pinned Brownian motion on any compact Riemannian manifold is a semimartingale up to time 1. However, the $L^p(\nu)$ -convergence ($p \geq 1$) and the Fernique type exponential integrability theorem for the stochastic anti-development of pinned Brownian motions were first proved by L. Gross on a compact Lie group, see Lemma 4.8, Remark 4.9 and Corollary 4.10 (pp.291-294) in [13]. More precisely, let $L_e(G)$ be the loop group over a compact Lie group G equipped with an Ad -invariant metric and the left Cartan (or the right Cartan) connection, e its unit element. The anti-development of the pinned Brownian motion $\{g(s), s \in [0, 1]\}$ is given by

$$b(s) = \int_0^s g^{-1}(r) \circ dg(r), \quad s \in [0, 1].$$

The Gross theorem says that there exists a small λ_0 such that for all $\lambda < \lambda_0$ we have $E_\nu \left[\exp \left(\lambda \max_{s \in [0,1]} \|b(s)\|^2 \right) \right] < +\infty$. See also Gong [12] for an alternative proof. In this section we will use Lemma 5.2 to establish the $L^p(\nu)$ -convergence and a Fernique type exponential integrability theorem for the stochastic anti-development of pinned Brownian motion on any compact Riemannian manifold with a TSS connection. Our result also sharpens the exponential exponent λ_0 appeared in the Gross-Fernique theorem. We begin with the following

Theorem 8.1 *Let G be a compact Lie group equipped with an Ad -invariant metric and the left or the right Cartan connection. Then for all*

$$\lambda < \lambda_0 = \frac{1}{2} \inf \{ \|w\|_H^2 : w \in X, \|w\|_{2m,\alpha} = 1 \},$$

we have

$$E_\nu \left[\exp (\lambda \|b\|_{2m,\alpha}^2) \right] < +\infty.$$

Moreover, for any $\lambda < \frac{1}{2}$, we have

$$E_\nu \left[\exp \left(\lambda \max_{s \in [0,1]} \|b(s)\|^2 \right) \right] < +\infty.$$

Proof. For μ -a.s. $x \in X$, since $dg_x(s) = g_x(s) \circ dx(s)$, we have

$$b(s)(x) = \int_0^s [g_x(r)]^{-1} \circ dg_x(r) = x(s) \in W^{r,p}(X, T_e G), \quad s \in [0, 1].$$

By the Donsker-Varadhan [5] refinement version of the well-known Fernique lemma, we have

$$E_\mu \left[\exp (\lambda \|b\|_{2m,\alpha}^2) \right] = E_\mu \left[\exp (\lambda \|x\|_{2m,\alpha}^2) \right] < +\infty,$$

provided that

$$\lambda < \lambda_0 = \inf\{I(w) : w \in X, \|w\|_{2m,\alpha} = 1\},$$

where

$$I(w) = \frac{1}{2}\|w\|_H^2 1_{[w \in H]} + \infty 1_{[w \notin H]}.$$

Note that the function $\|\cdot\|_{2m,\alpha}^2$ is smooth in $X^{2m,\alpha}$ in the sense of Fréchet-Gâteaux. Thus, the Wiener functional $x \rightarrow \|x\|_{2m,\alpha}^2$ belongs to $W^{\infty,\infty}(X^{2m,\alpha})$. Hence Lemma 5.2 applies to $F(x) = \|x\|_{2m,\alpha}^2$. That is to say, for any $\epsilon > 0$ and any $p > 1$ we have

$$E_\nu \left[\exp(\lambda \|b\|_{2m,\alpha}^2) \right] \leq \left\{ E_\mu \left[\exp((1+\epsilon)p\lambda \|b\|_{2m,\alpha}^2) \right] \right\}^{\frac{1}{p(1+\epsilon)}} \| \|x\|_{2m,\alpha} \|_{W^{n, \frac{n^2 p \epsilon}{1+\epsilon}}(X)}^n,$$

where $n = 2\lfloor \frac{d}{2} \rfloor + 2$. Thus, for all $\lambda < \frac{\lambda_0}{(1+\epsilon)p}$, we have $E_\nu \left[\exp(\lambda \|b\|_{2m,\alpha}^2) \right] < +\infty$. Since $\epsilon > 0$ and $p > 1$ are arbitrary, for all $\lambda < \lambda_0$ we get $E_\nu \left[\exp(\lambda \|b\|_{2m,\alpha}^2) \right] < +\infty$.

Now $\|w\|_\infty := \max_{s \in [0,1]} \|w(s)\|_\infty \leq C\|w\|_{2m,\alpha}$. Thus $E_\nu \left[\exp\left(\lambda \max_{s \in [0,1]} \|b(s)\|^2\right) \right] < +\infty$ holds provided that

$$\lambda \sup_{b \neq 0} \left[\frac{\|b\|_\infty^2}{\|b\|_{2m,\alpha}^2} \right] \leq \lambda \sup_{w \in X \setminus \{0\}} \left[\frac{\|w\|_\infty^2}{\|w\|_{2m,\alpha}^2} \right] < \lambda_0.$$

Set $X^* = X \setminus \{0\}$. Then, for all

$$\begin{aligned} \lambda &< \frac{1}{2} \inf_{w \in X^*} \left[\frac{\|w\|_H^2}{\|w\|_{2m,\alpha}^2} \right] \inf_{w \in X^*} \left[\frac{\|w\|_{2m,\alpha}^2}{\|w\|_\infty^2} \right] \\ &\leq \frac{1}{2} \inf_{w \in X^*} \left[\frac{\|w\|_H^2}{\|w\|_{2m,\alpha}^2} \cdot \frac{\|w\|_{2m,\alpha}^2}{\|w\|_\infty^2} \right] \\ &= \frac{1}{2} \inf_{w \in X^*} \left[\frac{\|w\|_H^2}{\|w\|_\infty^2} \right] = \frac{1}{2}, \end{aligned}$$

i.e., for all $\lambda < \frac{1}{2}$, we have

$$E_\nu \left[\exp\left(\lambda \max_{s \in [0,1]} \|x(s)\|^2\right) \right] < +\infty.$$

The proof of Theorem 8.1 is completed. \square

Similarly to the proof of Theorem 8.1, if we replace $b(s) = \int_0^s g^{-1}(r) \circ dg(r)$ by

$$x(s) = \int_0^s U_r^{-1}(\gamma) \circ d\gamma(r), \quad \nu - \text{a.s. } \gamma \in L_{m_0}(M),$$

then we can prove the following Fernique type exponential integrability theorem for the stochastic anti-development of pinned Brownian motions on a compact Riemannian manifold.

Theorem 8.2 *Let M be a compact Riemannian manifold equipped with a torsion-skew symmetric (TSS) connection. Then for all*

$$\lambda < \lambda_0 := \frac{1}{2} \inf\{\|w\|_H^2 : w \in X, \|w\|_{2m,\alpha} = 1\},$$

we have

$$E_\nu [\exp(\lambda \|x\|_{2m,\alpha}^2)] < +\infty.$$

Moreover, for all $\lambda < \frac{1}{2}$, we have

$$E_\nu \left[\exp \left(\lambda \max_{s \in [0,1]} \|x(s)\|^2 \right) \right] < +\infty.$$

Finally, let us prove the $L^p(\nu)$ -convergence of the stochastic anti-development of pinned Brownian motion.

Theorem 8.3 *Let M be a compact Riemannian manifold equipped with a TSS connection. Then for any $p \geq 1$, $x(s)$ converges to $x(1)$ in $L^p(\nu)$ as s tends to 1. Moreover, for any $p \geq 1$, there exists a constant C_p such that*

$$\|x(s) - x(1)\|_{L^p(\nu)} \leq C_p(1-s)^{\frac{1}{2}}.$$

Proof. Similarly to the proof of Theorem 4.1, there exist a constant C and a pair $(r, q) \in \mathbb{N} \times (1, +\infty)$ such that for $p = 2n$, $n \in \mathbb{N}$,

$$\|x(s) - x(1)\|_{L^p(\nu)}^p \leq C \| \|x(s) - x(1)\|^p \|_{W^{r,q}(X)}.$$

With respect to the Wiener measure on the Wiener space X , $x(s) - x(1)$ is a centred Gaussian variable with variance $1 - s$. Thus, there exists a constant $C(n, q)$ such that

$$E_{\mu_0} [\|x(s) - x(1)\|^{2qn}] \leq C(n, q)(1-s)^{nq}.$$

On the other hand, for all $h_1, \dots, h_r \in H$, $i = 1, \dots, d$, we have

$$\begin{aligned} \nabla_{h_1} \dots \nabla_{h_r} (x_i(s) - x_i(1))^{2n} &= 2n(2n-1) \dots (2n-r)(x_i(s) - x_i(1))^{2n-r} \\ &\quad \times (h_1^i(s) - h_1^i(1)) \dots (h_r^i(s) - h_r^i(1)). \end{aligned}$$

From which one can easily verify that

$$\|\nabla^r \|x(s) - x(1)\|^{2n}\|_q \leq C(n, r, q)(1-s)^{\frac{2n-r}{2}}.$$

Therefore, we get

$$\|x(s) - x(1)\|_{L^{2n}(\nu)} \leq C(n, r, q) \left[(1-s)^{\frac{1}{2}} + (1-s)^{\frac{2n-r}{4n}} \right].$$

Note that r and q are independent of n . Hence

$$\|x(s) - x(1)\|_{L^{2n}(\nu)} \leq C_n(1-s)^{\frac{1}{2}}.$$

This yields that $\|x(s) - x(1)\|_{L^p(\nu)} \leq C_p(1-s)^{\frac{1}{2}}$ for all $p \geq 1$. □

9 Driver's flow on loop spaces via Doob's h -theory

For the completeness of the paper, in this section we derive the pull-back Driver flow equation on loop spaces through Doob's h -processes approach, from which we then explain why and how the gradient and the Hessian estimates of the logarithmic of the heat kernel play a crucial role in the problems of the construction of Driver flow on loop space and of the quasi-invariance of the pinned Wiener measure.

Recall that with respect to the pinned Wiener measure ν on loop space $L_{m_0}(M)$, the conditional Brownian motion γ_t is given by the following Stratonovich SDE

$$d\gamma_s = U_s \circ db_s, \quad \gamma_0 = m_0,$$

where $\{b_s, s \in [0, 1]\}$ is the anti-development of $\{\gamma_s, s \in [0, 1]\}$ through the Itô map $I : X \rightarrow P_{m_0}(M)$ which can be well-defined up to a slim subset of X . Moreover, $\{b_s, s \in [0, 1]\}$ is a semi-martingale with the following Doob-Meyer decomposition

$$db_s = d\beta_s + U_s^{-1} \nabla \log p_{1-s}(\gamma_s, m_0) ds,$$

where $\{\beta_s, s \in [0, 1]\}$ is a ν -Brownian motion on $(X, \mathcal{F}_s, \mathcal{F}, \nu)$. See e.g. Bismut[2], Driver[7], Enchev-Stroock[9] and Hsu[15] [16].

Let $h \in H$ and D_h be the vector field on $L_{m_0}(M)$. Let γ^t be the Driver flow on the loop space $L_{m_0}(M)$ given by

$$\begin{aligned} \dot{\gamma}^t &= D_h(\gamma^t) = U_s^t h(s) \\ \gamma^0 &= \gamma, \end{aligned}$$

where $\{U_s^t, s \in [0, 1]\}$ is the horizontal lift of $\{\gamma_s^t, s \in [0, 1]\}$. Using the intertwining formula for the differential of the stochastic development map $I^{-1} : P_{m_0}(M) \rightarrow X$, see [6] [10] [8] [15] [24] [21], the pull-back flow $b^t = I^{-1}(\gamma^t)$ satisfies the following equation

$$\begin{aligned} \frac{\partial}{\partial t} b_s^t &= \dot{h}(s) ds + q_h(\gamma^t, s) \circ d_s b_s^t, \\ b_s^0 &= b_s, \end{aligned}$$

where

$$d_s b_s^t = d\beta_s^t + [U_s^t]^{-1} \nabla \log p_{1-s}(\gamma_s^t, m_0) ds.$$

Thus, we have

$$\left[\frac{\partial}{\partial t} - q_h(\gamma^t, s) \right] d_s b_s^t = \dot{h}_s ds,$$

whence

$$\begin{aligned}
\left[\frac{\partial}{\partial t} - q_h(\gamma^t, s) \right] \circ d_s \beta_s^t &= \dot{h}_s ds - \left[\frac{\partial}{\partial t} - q_h(\gamma^t, s) \right] ([U_s^t]^{-1} \nabla \log p_{1-s}(\gamma_s^t, m_0)) ds \\
&= \dot{h}_s ds + q_h(\gamma^t, s) ([U_s^t]^{-1} \nabla \log p_{1-s}(\gamma_s^t, m_0)) ds \\
&\quad + [U_s^t]^{-1} \left(\frac{\partial}{\partial t} U_s^t \right) [U_s^t]^{-1} \nabla \log p_{1-s}(\gamma_s^t, m_0) ds \\
&\quad - [U_s^t]^{-1} \nabla^2 \log p_{1-s}(\gamma_s^t, m_0) \left(\frac{\partial}{\partial t} \gamma_s^t \right) ds.
\end{aligned}$$

In conclusion, we have derived that $\{\beta_s^t, s \in [0, 1]\}$ (the ν -Brownian motions with drift) verify the pull-back Driver flow equation given by

$$\begin{aligned}
\left[\frac{\partial}{\partial t} - q_h(\gamma^t, s) \right] \circ d_s \beta_s^t &= \dot{h}_s ds + q_h(\gamma^t, s) ([U_s^t]^{-1} \nabla \log p_{1-s}(\gamma_s^t, m_0)) ds \\
&\quad + [U_s^t]^{-1} \left(\frac{\partial}{\partial t} U_s^t \right) [U_s^t]^{-1} \nabla \log p_{1-s}(\gamma_s^t, m_0) ds \\
&\quad - [U_s^t]^{-1} \nabla^2 \log p_{1-s}(\gamma_s^t, m_0) U_s^t h(s) ds.
\end{aligned}$$

To solve the above flow equation by standard Picard iteration or Euler iteration method, it is now clear that one has to use the gradient estimate and the Hessian estimate of the logarithmic of the heat kernel on compact Riemannian manifold. Moreover, in order to use the usual approach based on Girsanov theorem and Levy's invariance of Brownian motion under adapted rotations, we need to verify the Novikov exponential integrability condition of the drift term given in the right hand side of the flow equation for β^t . Thus, we need to use again the gradient estimate and the Hessian estimate of the logarithmic heat kernel. For these, we refer the reader to Driver[7], Enchev-Stroock [9] and Hsu [15] [16]. See also Sheu [29], Malliavin-Stroock [27], Stroock-Turetsky [31] and Hsu[17] for the short time estimates of logarithmic derivatives of the heat kernel.

To end this paper, let us mention that F. Z. Gong has informed us that using the gradient and the Hessian estimates of the logarithmic of the heat kernel, Gong and Zhi-Ming Ma can also prove the $L^p(\nu)$ -convergence and the ν -exponential integrability of the divergence functional $\delta(h)$ (for all $\lambda < \lambda_0$ for some constant λ_0 which depends on $\|h\|_H$ and possibly on the constants appeared in the gradient and the Hessian estimates of the logarithmic heat kernel) by the Doob h -processes approach. Our approach based on Airault-Malliavin-Sugita-Watanabe inequality (see Theorem 2.2 and Lemma 5.2) shows that we can get an explicit and optimal estimate for λ_0 which only depends on the upper bound of the Ricci curvature $\|Ric\|_\infty$ and $\|h\|_H$, i.e., $\lambda_0 = [(2 + \|Ric\|_\infty)\|h\|_H]^{-1}$. To help the reader to understand this strange phenomenon which has been already appeared in Section 8, we allow ourself to cite the following paragraphe due to P. Malliavin [23] (Section 5 Page 73).

“To describe a diffusion with a prescribed value at time 1, the analytic methods are developed from Doob’s theory of h -processes. To build the h -process, it is necessary to know the fundamental solution h of a parabolic equation. Since very often, the fundamental solution is approached through an S.D.E., the use of the h -theory is, in some sense, tautological. Even more, the fundamental solution h can be fully known, as for the Brownian case, and the h -theory can lead to **poor** estimates due to the apparent singularity which appears near $\tau = 1$.”

Acknowledgement. The earliest version of this paper was done in 1996 and appeared in the author’s PhD thesis [22] under the supervision of Professors Zhi-Ming Ma, P. Malliavin and A. B. Cruzeiro. The author thanks in particular Professors Shizan Fang and P. Malliavin for suggesting him to use quasi-sure analysis to study the problems in this paper. He is very grateful to Prof. Zhi-Ming Ma to introducing him into mathematical research fields and to recommend the earlier announcement [20] of Theorem 1.1 to publish in *Chinese Sc. Bull.*. Thanks are also due to Professors A. B. Cruzeiro, B. Driver, Fuzhou Gong, E. Hsu, R. Léandre, T. Lyons, J. Ren and D. W. Stroock for helpful discussions and constant encouragements.

References

- [1] AIRAULT, H., MALLIAVIN, P. (1988). Intégration géométrique sur l’espace de Wiener. *Bull. Sci. Math.* (2) **112** 3-52.
- [2] BISMUT, J. M. (1984). *Large Deviations and the Malliavin Calculus. Progress in Mathematics.* Birkhäuser.
- [3] CROSS, C. (1996). Differentials of Measure-Preserving Flows on Path Space. PhD thesis, UCSD. It can be download from <http://math.ucsd.edu/~driver/driver/thesis.htm>.
- [4] CRUZEIRO, A. B. (1983). Équations différentielles sur l’espace de Wiener et formules de Cameron-Martin non linear. *J. Funct. Anal.* **54** 206-227.
- [5] DONSKER, M. D., VARADHAN, S. R. S. (1976). Asymptotic evaluation of certain Markov process expectations for large time. III. *Comm. Math. Phys.* **29** 389-461.
- [6] DRIVER, B. K. (1992). A Cameron-Martin type quasi-invariance theorem for Brownian motion on a compact manifold. *J. Funct. Anal.* **109** 272-376.
- [7] DRIVER, B. K. (1994). A Cameron-Martin type quasi-invariance theorem for pinned Brownian motion on a compact manifold. *Trans. Amer. Math. Soc.* (1) **342** 375-394.

- [8] ENCHEV, O., STROOCK, D. W. (1996). Towards a Riemannian geometry on the path space over a Riemannian manifold. *J. Funct. Anal.* **134** 392-416.
- [9] ENCHEV, O., STROOCK, D. W. (1996). Pinned Brownian motion and its perturbations. *Adv. Math.* **119** 127-154.
- [10] FANG, S., MALLIAVIN, P. (1993). Stochastic analysis on the path space of a Riemannian manifold, I. Markovian stochastic calculus. *J. Funct. Anal.* **118** 249-274.
- [11] FERNIQUE, X. (1970). Intégrabilité des vecteurs gaussiens. *C. R. Acad. Sci. Paris.* **270** 1098-1099.
- [12] GONG, F. Z. (1995). Some Aspects of Infinite Dimensional Stochastic Analysis. PhD Thesis, Academia Sinica.
- [13] GROSS, L. (1991). Logarithmic Sobolev inequalities on loop groups. *J. Funct. Anal.* **102** 268-313.
- [14] HSU, E. (1996). Quasi-invariance of the Wiener measure on the path space over a compact Riemann manifold. *J. Funct. Anal.* **134** 417-450.
- [15] HSU, E. (1997). Integration by parts in loop spaces. *Math. Ann.* (2) **309** 331-339.
- [16] HSU, E. (1997). *Analysis on path and loop spaces. IAS/Park City Math. Series.* Vol 5. Ed. by E.P. Hsu and S.R.S. Varadhan. Amer. Math. Soc.
- [17] HSU, E. (1999). Estimates of derivatives of the heat kernels on a compact Riemannian manifold. *Proc. Amer. Math. Soc.* (12) **127** 3739-3744.
- [18] LÉANDRE, R. (2001). Stochastic Adams theorem for a general compact manifold. *Rev. Math. Phys.* (9) **13** 1095-1133.
- [19] LEDOUX, M., TALAGRAND, M. (1991) *Probability in Banach Spaces.* Springer-Verlag.
- [20] LI, X. D. (1998). Asymptotic behavior of the divergence on loop spaces over a compact Riemannian manifold. *Chinese Sci. Bull.* (4) **43** 272-274. It can be downloaded from <http://www.maths.ox.ac.uk/~lix>.
- [21] LI, X. D. Sobolev spaces and capacities theory on path spaces over a compact Riemannian manifold. To appear in *Probability Theory and Related Fields*. It can be downloaded from <http://www.maths.ox.ac.uk/~lix>.
- [22] LI, X. D. (1999). Stochastic Analysis and Geometry on Path and Loop Spaces. PhD thesis, Academia Sinica and University of Lisbon. It can be downloaded from <http://www.maths.ox.ac.uk/~lix>.

- [23] MALLIAVIN, P. (1993). Infinite dimensional analysis. *Bull. Sci. Math.* 2^e série **117** 63-90.
- [24] MALLIAVIN, P. (1997). *Stochastic Analysis*. Springer-Verlag, Berlin.
- [25] MALLIAVIN, M. P., MALLIAVIN, P. (1990). Integration on loop groups, I. Quasi invariant measures. *J. Funct. Anal.* **93** 207-237.
- [26] MALLIAVIN, P., NUALART, D. (1993). Quasi-sure analysis of stochastic flows and Banach space valued smooth functionals on the Wiener space. *J. Funct. Anal.* **112** 287-317.
- [27] MALLIAVIN, P., STROOCK, D. W. (1996) Short time behavior of the heat kernels and its logarithmic derivatives. *J. Differential Geometry.* (3) **44** 550-570.
- [28] REN, J. (1990). Analysis quasi-sûre des équations différentielles stochastiques. *Bull. Sc. Math.* 2^e série **114** 187-213.
- [29] SHEU, S. Y. (1991) Some estimates of the transition density function of a nondegenerate diffusion Markov process. *Ann. of Prob.* (2) **19** 538-561.
- [30] SHIGEKAWA, I. (1993). A quasihomomorphism on the Wiener space. *Stochastic Analysis. Proceedings of Symposia in Pure Mathematics*. Ithaca. Vol. **57** 473-486
- [31] STROOCK, D. W., TURETSKY, J. (1997) Short time behavior of logarithmic derivatives of the heat kernels. *Asian J. Math.* **1** 17-33.
- [32] SUGITA, H. (1986). Sobolev spaces of Wiener functionals and Malliavin's calculus. *J. Math. Kyoto Univ.* (1) **25** 31-48.
- [33] SUGITA, H. (1988). Positive generalized functions and potential theory over an abstract Wiener space. *Osaka J. Math.* **25** 665-696.
- [34] WATANABE, S. (1987). Analysis of Wiener functionals (Malliavin calculus) and its applications to heat kernels. *Ann. of Prob.* (1) **15** 1-39

Mathematical Institute
 University of Oxford
 24-29, St. Giles'
 Oxford, OX1 3LB, UK
 E-mail: lix@maths.ox.ac.uk