

Coalescences in Continuous-State Branching Processes

Clément Foucart ^{*} Chunhua Ma [†] Bastien Mallein [‡]

September 6, 2019

Abstract

Consider a continuous-state branching population constructed as a flow of nested subordinators. Inverting the subordinators and reversing time give rise to a flow of coalescing Markov processes with negative jumps, which correspond to the ancestral lineages of individuals in the current generation. The process of the ancestral lineage of a fixed individual is the Siegmund dual process of the continuous-state branching process. We study its semi-group, its long-term behaviour and its generator. In order to follow the coalescences in the ancestral lineages and to describe the backward genealogy of the population, we define non-exchangeable Markovian coalescent processes obtained by sampling individuals according to an independent Poisson point process over the flow. These coalescent processes are called consecutive coalescents, as only consecutive blocks can merge. They are characterized in law by finite measures on \mathbb{N} which can be thought as the offspring distributions of some inhomogeneous immortal Galton-Watson processes forward in time.

Keywords: Branching processes, coalescent processes, continuous-state branching processes, flow of subordinators, genealogy, duality.

Introduction

Random population models can be divided in two classes, those with constant finite size and those whose size is varying randomly. It is known since the 2000s that populations with constant finite size, evolving by resampling, have genealogies given by exchangeable coalescents. These processes, defined by Möhle and Sagitov [MS01], Pitman [Pit99], Sagitov [Sag99] and Schweinsberg [Sch00], are generalisations of Kingman's coalescent for which multiple coalescences of ancestral lineages are allowed. They correspond to the genealogy backward in time of so-called generalized Fleming-Viot processes. Those processes, which can be seen as scaling limits of Moran models [Mor58], were defined and studied by Donnelly and Kurtz [DK99] (via a particle system called lookdown construction) and by Bertoin and Le Gall [BLG03] (via flows of exchangeable bridges). Both constructions are similar in many aspects and are summarized via the notion of flow of partitions, see Labbé [Lab14a, Lab14b] and Foucart [Fou12]. We refer to Bertoin's book [Ber06] for a comprehensive account on exchangeable coalescents.

^{*}LAGA, UMR 7539, CNRS, Université Paris 13 & Université Paris 8, France. foucart@math.univ-paris13.fr

[†]School of Mathematical Sciences and LPMC, Nankai University, China. mach@nankai.edu.cn

[‡]LAGA, UMR 7539, Université Paris 13 & Université Paris 8, France. mallein@math.univ-paris13.fr

The main objective of this work is to study coalescent processes induced by branching processes. We briefly explain how branching concepts have been developed from the sixties to the beginning of the twenty-first century. Continuous-state branching processes (CSBPs for short) are positive Markov processes representing the size of a continuous population. They have been defined by Jiřina [Jiř58] and Lamperti [Lam67a] and are known to be scaling limits of Galton-Watson Markov chains, see Grimvall [Gri74] and Lamperti [Lam67b]. The most famous CSBP is certainly Feller’s branching diffusion

$$dX_t = \sigma\sqrt{X_t}dB_t + \beta X_t dt$$

which is the rescaled limit of binary branching processes, see Feller [Fel51] and Jiřina [Jiř69]. Feller’s CSBP is the only CSBP with continuous paths, other ones have positive jumps which represent macroscopic reproduction events in the population.

At about the same time as the rise of exchangeable coalescents, considerable research was devoted to the study of the genealogy of branching processes forward in time. Galton-Watson processes have a natural lexicographical tree’s genealogy. This representation led Aldous [Ald93] and Duquesne and Le Gall [DLG02] to study scaling limits of discrete trees and establish remarkable convergences towards the Brownian continuum tree in the Feller diffusion case and Lévy continuum trees in the case of general CSBPs. Another natural genealogy for a branching population is provided by Bertoin and Le Gall in their precursor article [BLG00] in terms of flows of subordinators. At any fixed times $s < t$, the population between time s and t is represented by a subordinator (a Lévy process with non-decreasing paths) $(X_{s,t}(x), x \geq 0)$. Individuals are ordered in such a way that ancestors from time s are the jumps locations of the subordinator and each ancestor from time s has a family at time t whose size is the size of the jump.

Both representations with trees and subordinators are future-oriented and less attention has been paid to the description of coalescences in ancestral lineages of continuous-state branching processes. We briefly review some methods that have been developed recently in order to study the genealogy backwards in time of branching processes.

When reproduction laws are stable, branching and resampling population models can be related through renormalisation by the total size and random time-change. We refer to Berestycki et al. [BBS07], Birkner et al. [BBC+05], Foucart and Hénard [FH13] and Schweinsberg [Sch03]. The connection between exchangeable coalescents and CSBPs is particular to stable laws and the study of the genealogy of a general branching process requires a different method.

One approach consists in conditioning the process to be non-extinct at a given time, sampling two or more individuals uniformly in the population and study the time of coalescence of their ancestral lineages. This program is at the heart of the works of Athreya [Ath12], Duquesne and Labbé [DL14], Harris et al. [HJR17], Johnston [Joh17], Lambert [Lam03] and Le [Le14].

Starting from a different point of view, Bi and Delmas [BD16] and Chen and Delmas [CD12] have considered stationary subcritical branching populations obtained as processes conditioned on the non-extinction. The genealogy is then studied via a Poisson representation of the population. We refer also to Evans and Ralph [ER10] for a study in the same spirit.

A third approach is to represent the backwards genealogy through point processes. Aldous and Popovic [AP05] and Popovic [Pop04] have shown how to encode the genealogy of a critical Feller diffusion with a Poisson point process on $\mathbb{R}_+ \times \mathbb{R}_+$ called Coalescent Point Process. Atoms of the coalescent point process represent times of coalescences between two “consecutive” individuals in the boundary of the Brownian tree. Such a description was further developed by Lambert and Popovic [LP13] for a Lévy continuum tree. In this general setting, multiple

coalescences are possible and the authors build a point process with multiplicities, which records both the coalescence times and the number of involved mergers in the families of the current population. Their method requires in particular to work with the height process introduced by Le Gall and Le Jan in [LGLJ98].

In the present article, we choose a different route and seek a dynamical description of the genealogy. We first observe that flows of subordinators provide a continuous branching population whose size is infinite at any time and whose ancestors are arbitrarily old. We then define and study the inverse flow, denoted by $(\hat{X}_{s,t}(x), s \leq t, x \geq 0)$, which tracks backward in time the ancestral lineage of an individual considered at any given time. In particular, the process $(\hat{X}_t(x), t \geq 0) := (\hat{X}_{0,t}(x), t \geq 0)$ is the ancestral lineage of the individual x in the population taken at time 0. This is a Markov process and we characterize its semi-group, its long-term behaviour (recurrent or transient) as well as its generator.

In a second time, we introduce new elementary non-exchangeable Markovian coalescents as simple dual objects of immortal continuous-time Galton-Watson processes. These processes are taking values in the set of partitions of \mathbb{N} whose blocks are formed with consecutive integers. We call them *consecutive coalescents* as only consecutive blocks will be allowed to merge. These coalescent processes represent the genealogy of immortal continuous-time Galton-Watson processes when time's arrow points to the past.

We will use these coalescent processes to describe the genealogy of general CSBPs. They will simplify the description given by the Coalescent Point Process as introduced in [LP13, Section 4]. Our method follows closely that of Bertoin and Le Gall for exchangeable coalescents [BLG03], [BLG05], [BLG06]. Heuristically, exchangeable bridges are replaced by subordinators and uniform random variables by atoms of a Poisson process. We shall construct random partitions by sampling individuals according to an independent Poisson process. Namely, let $(J_i^\lambda, i \geq 0)$ be the sequence of atoms (i.e jumping times) of an independent Poisson process with intensity λ and consider the random partition $C^\lambda(t)$ defined by letting integers i and j in the same equivalence class if and only if $\hat{X}_t(J_i^\lambda) = \hat{X}_t(J_j^\lambda)$. We will show that the process $(C^\lambda(t), t \geq 0)$ is a (possibly time-inhomogeneous) consecutive coalescent. We characterize its jump rates and its long-term behaviour.

We shall also show how to define the complete genealogy of individuals standing in the current generation when the so-called Grey's condition is satisfied. Loosely speaking, we will let the intensity parameter λ to infinity and describe the genealogy through coalescing consecutive intervals of $[0, \infty]$. This answers an open question in [LP13, Remark 6].

We apply then our results to the specific cases of Neveu and stable CSBPs. In the case of Neveu's CSBP (which does not fulfill Grey's condition), Bertoin and Le Gall in [BLG00] have shown that the genealogy of the CSBP, started from a fixed size (without renormalization nor time-change) is given by a Bolthausen-Sznitman coalescent. We will see that for this CSBP, the consecutive coalescents have simple explicit laws. This will enable us to recover results of Möhle [Möh15] and Möhle and Kukla [KM18] about the number of blocks in a Bolthausen-Sznitman coalescent. In the Neveu case, the process $(\hat{X}_t(x), t \geq 0)$ is related to a generalized Ornstein-Uhlenbeck process with negative jumps. We compute explicitly its Laplace exponent and recover a result due to Bertoin and Baur [BB15]. In the stable case, the consecutive coalescent is a simple deterministic time-change of a time-homogeneous consecutive coalescent whose jump rates take a simple explicit form. Last, in the critical stable case, the process $(\hat{X}_t(x), t \geq 0)$ is a positive self-similar process and we compute the Laplace exponent of its Lévy parent process.

We wish to mention that Grosjean and Huillet in [GH16] have studied a recursive balls-in-boxes model which can be seen as a consecutive coalescent in discrete time. Moreover, Johnston and Lambert [JL19+] have independently considered Poissonization techniques for studying the coalescent structure in branching processes.

The paper is organized as follows. In Section 1, we recall the definition of a continuous-state branching process and how Bochner’s subordination can be used to provide a representation of the genealogical structure associated with CSBPs. In Section 2, we investigate the inverse flow by characterizing its semi-group and studying its long-term behavior. In Section 3, we provide a complete study of the inverse flow in the case of the Feller diffusion. We recover with an elementary approach the Coalescent Point Process of Popovic [Pop04]. In Section 4, we study the coalescences in the inverse flow of a general CSBP by defining the consecutive coalescents. We describe the genealogy of the whole population standing at the current generation under Grey’s condition (recalled in Section 1). In Section 5, we investigate the infinitesimal dynamics of the inverse flow. The process of the ancestral lineage of a fixed individual is characterized by its generator. In Section 6, we focus on the stable and Neveu CSBPs for which calculations can be made explicitly. In the Appendix, we gather some elementary properties on right continuous inverse of càdlàg non-decreasing functions. We also establish some key results, which are needed in Section 4, on the discretization of subordinators through Poisson sampling.

Contents

1	Generalities on continuous-state branching processes	5
1.1	Continuous-state branching processes	5
1.2	Flows of subordinators	7
2	The inverse flow	9
3	The Feller flow	15
4	Consecutive coalescents	21
4.1	Consecutive coalescents in continuous-time Galton-Watson processes	21
4.2	Consecutive coalescents in CSBPs through Poisson sampling	26
4.3	Backward genealogy of the whole population	32
5	A martingale problem for the inverse flow	36
6	Examples	45
6.1	Feller and stable CSBPs	45
6.2	Neveu CSBP	47
A	Intermediary results	49
A.1	Right-continuous inverse	49
A.2	Discretization of subordinators	51

Notation. In the rest of the article, $\stackrel{\mathcal{L}}{=}$ denotes equality in law between random variables. Condition $\int_0^\infty f(x)dx < \infty$ means there exists $\epsilon > 0$ such that $\int_0^\epsilon f(x)dx < \infty$, and similarly $\int_A^\infty f(x)dx < \infty$ means there exists $A > 0$ such that $\int_A^\infty f(x)dx < \infty$. For any $n, m \in \mathbb{N}$ such that $n \leq m$, the integer interval between n and m is denoted by $[[n, m]]$. The set $C([0, \infty])$ is the space of continuous functions f on $[0, \infty)$, with finite limit at ∞ written $f(\infty)$. We write $f(t) = O(g(t))$ as t goes to ∞ if for a large enough t_0 , g is non-negative and $\sup_{t \geq t_0} \frac{|f(t)|}{g(t)} < \infty$.

1 Generalities on continuous-state branching processes

This section is divided in two parts. In the first one, we introduce the continuous-state branching processes as well as their fundamental properties. In the second one, we show how continuous-state branching processes can be constructed as flows of subordinators. Their main properties are also stated.

1.1 Continuous-state branching processes

We give here an overview of continuous-state branching processes and their fundamental properties. Most statements in this section can be found for instance in [Lil1, Chapter 3] or [Kyp14, Chapter 12].

Definition 1.1. A continuous-state branching process is a Feller process $(X_t, t \geq 0)$, taking values in $[0, \infty]$, with 0 and ∞ being absorbing states, whose semi-group satisfies the so-called branching property:

$$\forall x, y \geq 0, \forall t \geq 0, X_t(x + y) \stackrel{\mathcal{L}}{=} X_t(x) + \tilde{X}_t(y), \quad (1.1)$$

where $(X_t(x), t \geq 0)$ and $(\tilde{X}_t(y), t \geq 0)$ are two independent processes with the same law as $(X_t, t \geq 0)$, started respectively from x and y .

The branching and the Markov properties ensure the existence of a map $\lambda \in (0, \infty) \mapsto v_t(\lambda)$ for all $t \geq 0$, which satisfies for all $\lambda > 0$, $x \geq 0$ and $t, s \geq 0$

$$\mathbb{E}[e^{-\lambda X_t(x)}] = \exp(-xv_t(\lambda)) \text{ and } v_{s+t}(\lambda) = v_s \circ v_t(\lambda). \quad (1.2)$$

Silverstein [Sil68] shows that $t \mapsto v_t(\lambda)$ is the unique solution to the integral equation

$$\forall t \in [0, \infty), \forall \lambda \in (0, \infty) \setminus \{\rho\}, \int_{v_t(\lambda)}^\lambda \frac{dz}{\Psi(z)} = t \quad (1.3)$$

where $\rho := \inf\{z > 0; \Psi(z) \geq 0\}$ is the largest positive root of Ψ , a Lévy-Khintchine function of the form

$$\Psi(q) = \frac{\sigma^2}{2}q^2 - \beta q + \int_0^\infty (e^{-qx} - 1 + qx\mathbb{1}_{x \leq 1}) \pi(dx), \quad (1.4)$$

with $\sigma^2 \geq 0$, $\beta \in \mathbb{R}$ and π a measure on $(0, \infty)$ satisfying $\int (1 \wedge x^2)\pi(dx) < \infty$. The function Ψ is called branching mechanism and characterizes the law of the CSBP. We shall write later $\text{CSBP}(\Psi)$ for a CSBP with branching mechanism Ψ . The generator of the $\text{CSBP}(\Psi)$ acts as follows

$$\mathcal{L}f(z) = z\frac{\sigma^2}{2}f''(z) + \beta zf'(z) + z \int_0^\infty (f(z+h) - f(z) - hf'(z)\mathbb{1}_{h \leq 1}) \pi(dh) \quad (1.5)$$

on the space of twice continuously differentiable functions f with compact support. The CSBP has infinite variations if

$$\int_0^1 x\pi(dx) = \infty \quad \text{or} \quad \sigma^2 > 0. \quad (1.6)$$

An important family of branching mechanisms are those of the form

$$\Psi(q) = \frac{\sigma^2}{2}q^2 - \beta q + c_\alpha q^\alpha$$

with $\sigma^2 \geq 0$, $c_\alpha \geq 0$ and $\alpha \in (0, 2) \setminus \{1\}$. The Lévy measure π associated with such a mechanism Ψ is

$$\pi(dh) = c'_\alpha h^{-1-\alpha} dh, \quad \text{with } c'_\alpha = \frac{\alpha(\alpha-1)}{\Gamma(2-\alpha)} c_\alpha.$$

The CSBP(Ψ) is said to be supercritical, critical or subcritical if respectively $\Psi'(0^+) < 0$, $\Psi'(0^+) = 0$ or $\Psi'(0^+) > 0$. In the subcritical and critical cases, the largest root ρ is 0. In the supercritical case $\rho \in (0, \infty]$. The following theorem due to Grey [Gre74] summarizes the possible behaviors at the boundaries of a CSBP(Ψ).

Theorem 1.A (Grey, [Gre74]). *Consider $(X_t(x), t \geq 0)$ a CSBP(Ψ) started from x .*

(i) *For any $x \geq 0$,*

$$\mathbb{P}(\lim_{t \rightarrow \infty} X_t(x) = 0) = 1 - \mathbb{P}(\lim_{t \rightarrow \infty} X_t(x) = \infty) = e^{-x\rho}.$$

(ii) *For any $t > 0$, the limit $v_t(\infty) := \lim_{\lambda \rightarrow \infty} v_t(\lambda)$ exists in $(0, \infty)$ if and only if $\Psi(u) \geq 0$ for some $u \geq 0$ and*

$$\int^\infty \frac{du}{\Psi(u)} < \infty \quad (\text{condition for extinction}). \quad (1.7)$$

If (1.7) holds, then for any $t \geq 0$, $\mathbb{P}(X_t(x) = 0) = e^{-xv_t(\infty)} > 0$.

(iii) *Under condition (1.7), the following events coincide almost-surely*

$$\left\{ \lim_{t \rightarrow \infty} X_t(x) = 0 \right\} = \{ \exists t \geq 0 : X_t(x) = 0 \}.$$

(iv) *For any $t > 0$, the limit $v_t(0) := \lim_{\lambda \rightarrow 0} v_t(\lambda)$ exists in $(0, \infty)$ if and only if $\Psi(u) < 0$ for some $u \geq 0$ and*

$$\int_0^\infty \frac{du}{|\Psi(u)|} < \infty \quad (\text{condition for explosion}). \quad (1.8)$$

If (1.8) holds, then for any $t \geq 0$, $\mathbb{P}(X_t(x) = \infty) = 1 - e^{-xv_t(0)}$.

(v) *Under condition (1.8), the following events coincide*

$$\left\{ \lim_{t \rightarrow \infty} X_t(x) = \infty \right\} = \{ \exists t \geq 0 : X_t(x) = \infty \}.$$

The event $\{X_t(x) = 0 \text{ for some } t \geq 0\}$ is called extinction, and $\{X_t(x) = \infty \text{ for some } t \geq 0\}$ is called explosion. We refer to the integral conditions (1.7) and (1.8) as Grey's condition for extinction and explosion respectively. Lambert [Lam07] and Li [Li00] have studied the quasi-stationary distribution of subcritical CSBPs conditioned on the non-extinction.

Theorem 1.B (Lambert [Lam07], Li [Li00]). *In the subcritical case, under Grey's condition for extinction $\int_0^\infty \frac{du}{\Psi(u)} < \infty$, there exists a probability measure ν over $(0, \infty)$ such that for any Borelian set $A \subset (0, \infty)$*

$$\nu(A) := \lim_{t \rightarrow \infty} \mathbb{P}(X_t(x) \in A | X_t(x) > 0).$$

The Laplace transform of ν is given by

$$\int_0^\infty e^{-uz} \nu(dz) = 1 - e^{-\Psi'(0^+) \int_u^\infty \frac{dx}{\Psi(x)}} \text{ for any } u \geq 0. \quad (1.9)$$

1.2 Flows of subordinators

Observe that on the one hand, by the branching property of CSBP, the random variable $X_t(x)$ is a positive infinitely divisible random variable, parametrized by x . Therefore, for all $t \geq 0$, the process $x \mapsto X_t(x)$ is a positive Lévy process, hence a subordinator. In particular, the map $\lambda \mapsto v_t(\lambda)$ is the Laplace exponent of this (possibly killed) subordinator, and can be written as

$$v_t(\lambda) = \kappa_t + d_t \lambda + \int_{[0, \infty)} (1 - e^{-\lambda u}) \ell_t(du) \quad (1.10)$$

with $\kappa_t \geq 0$, $d_t \geq 0$ and ℓ_t a Lévy measure on \mathbb{R}_+ such that $\int_0^\infty (1 \wedge u) \ell_t(du) < \infty$. As usual, κ_t can be thought of as a mass at ∞ of ℓ_t , i.e. $\kappa_t = \ell_t(\{\infty\})$.

Remark 1.2. Note that the quantities $v_t(\infty)$ and $v_t(0)$ defined in Theorem 1.A can be rewritten, with the formula in (1.10)

$$v_t(\infty) = \kappa_t + d_t \cdot \infty + \ell_t((0, \infty)) = d_t \cdot \infty + \ell_t((0, \infty]) \quad \text{and} \quad v_t(0) = \kappa_t.$$

Therefore (1.7) holds if and only if for all $t > 0$, the measure ℓ_t is finite and $d_t = 0$. Condition (1.8) is equivalent to the positivity of κ_t for all $t > 0$.

On the other hand, the semigroup property entails that for any $s, t \geq 0$,

$$v_{t+s} = v_t \circ v_s. \quad (1.11)$$

Bochner's subordination implies that if $Y^{(t)}$ is a subordinator with Laplace exponent v_t and $Y^{(s)}$ is a subordinator with Laplace exponent v_s , then $Y^{(t)} \circ Y^{(s)}$ is a subordinator with Laplace exponent $v_t \circ v_s = v_{t+s}$. Therefore, writing \tilde{X} an independent copy of the CSBP X , we have

$$\forall x \geq 0, X_{t+s}(x) \stackrel{\mathcal{L}}{=} \tilde{X}_t(X_s(x)).$$

This last observation led Bertoin and Le Gall [BLG00] to consider representing a CSBP as a flow of subordinators, which we now define.

Definition 1.3. A flow of subordinators is a family $(X_{s,t}(x), s \leq t, x \geq 0)$ satisfying the following properties:

- (i) For every $s \leq t$, $x \mapsto X_{s,t}(x)$ is a càdlàg subordinator, with same law as $x \mapsto X_{0,t-s}(x)$.
- (ii) For every $t \in \mathbb{R}$, $(X_{r,s}, r \leq s \leq t)$ and $(X_{r,s}, t \leq r \leq s)$ are independent.
- (iii) For every $r \leq s \leq t$, $X_{r,t} = X_{s,t} \circ X_{r,s}$.
- (iv) For every $s \in \mathbb{R}$ and $x \geq 0$, we have $X_{s,s}(x) = x = \lim_{t \rightarrow s} X_{s,t}(x)$ in probability.

Remark 1.4. The convergence in (iv) also holds uniformly on compact sets by Dini's second theorem.

It was proved by Bertoin and Le Gall [BLG00] that any CSBP can be constructed as a flow of subordinators. For the sake of completeness, we prove here that CSBP and flow of subordinators are in one-to-one map.

Lemma 1.5. *Let $(X_{s,t}(x), s \leq t, x \geq 0)$ be a flow of subordinators as in Definition 1.3. There exists a branching mechanism Ψ such that for all $s \in \mathbb{R}$ and $x \geq 0$, $(X_{s,s+t}(x), t \geq 0)$ is a CSBP(Ψ) starting from x . Reciprocally, for each branching mechanism Ψ , there exists a flow of subordinators such that for all $s \in \mathbb{R}$ and $x \geq 0$, $(X_{s,s+t}(x), t \geq 0)$ is a CSBP(Ψ) starting from x .*

Proof. Let $(X_{s,t}(x), s \leq t, x \geq 0)$ be a flow of subordinators. By Definition 1.3(ii) and (iii), we have that $t \mapsto X_{s,s+t}(x)$ is a Markov process for all $x \geq 0$ and $s \in \mathbb{R}$. Moreover, Definition 1.3(iv) implies this Markov process to be continuous in probability. Since by Definition 1.3(i), $x \mapsto X_{s,s+t}(x)$ is càdlàg, the semigroup of $(X_{s,s+t}(x), t \geq 0)$ maps $C([0, \infty])$ to $C([0, \infty])$, it is therefore Feller. Finally, by Definition 1.3(i), we conclude that this Markov process is homogeneous in time, and satisfies the branching property (1.1), as

$$X_{s,s+t}(x+y) = X_{s,s+t}(x) + (X_{s,s+t}(x+y) - X_{s,s+t}(x)),$$

and $X_{s,s+t}(x+y) - X_{s,s+t}(x)$ is independent of $X_{s,s+t}(x)$ and has same law as $X_{s,s+t}(y)$. Reciprocally, by [BLG00, Proposition 1], given a branching mechanism Ψ , there exists a process $(S^{(s,t)}(a), s \leq t, a \geq 0)$ such that almost surely

- (i) for all $s \leq t$, $a \mapsto S^{(s,t)}(a)$ is a càdlàg subordinator with Laplace exponent $\lambda \mapsto v_{t-s}(\lambda)$, defined in (1.3),
- (ii) for all $t \in \mathbb{R}$, $(S^{(r,s)}, r \leq s \leq t)$ and $(S^{(r,s)}, t \leq r \leq s)$ are independent,
- (iii) for all $r \leq s \leq t$, $S^{(s,t)} \circ S^{(r,s)} = S^{(r,t)}$,
- (iv) the finite dimensional distributions of $t \mapsto S^{(s,s+t)}(a)$ are the ones of a CSBP(Ψ).

One readily observes that points (i)–(iii) imply Definition 1.3(i)–(iii). Moreover, by the fourth point, $(S^{(s,s+t)}(a), t \geq 0)$ has the law of a CSBP(Ψ) $(X_t, t \geq 0)$ starting from $X_0 = a$. As X is a Feller process, we have $\lim_{t \rightarrow 0} X_t = a$ in probability, thus (iv) yields $\lim_{t \rightarrow 0} S^{(s,s+t)}(a) = a$ in probability, completing the proof. \square

A noteworthy consequence of the above lemma is that if $(X_{s,t}(x), s \leq t, x \geq 0)$ is a flow of subordinators associated with the branching mechanism Ψ , we have that for all $s \leq t$ and $x \geq 0$,

$$\forall \lambda \in (0, \infty), \mathbb{E}(\exp(-\lambda X_{s,t}(x))) = \exp(-xv_{t-s}(\lambda)), \quad (1.12)$$

where $v_{t-s}(\lambda)$ is the function defined in (1.3). One can think of this flow of subordinators as a way to couple on the same probability space every Markov property and every branching property (1.1), for all values of t, x, y simultaneously in one process.

The flow of subordinators provides a genuine continuous-space branching population model. More precisely, the interval $[0, X_{s,t}(x)]$ can be interpreted as the set of descendants at time t of the population that was represented at time s as the interval $[0, x]$. With this interpretation, the genealogy forward in time of the population is defined as follows. If $X_{s,t}(y-) < X_{s,t}(y)$, we say that for all $z \in (X_{s,t}(y-), X_{s,t}(y)]$, the individual z at time t is a descendant of the individual y living at time s . If $X_{s,t}(y-) = X_{s,t}(y)$ (i.e. $X_{s,t}$ is continuous at y) and $y = \inf\{x > 0 : X_{s,t}(x) = z\}$, we then say that individual $z = X_{s,t}(y)$ at time t is the descendant of the individual y living at time s . One can observe that the cocycle property ensures that this construction indeed defines a genealogy. If z at time t is a descendant of y at time s , which is a descendant of x at time r , we have

$$X_{s,t}(y-) < z \leq X_{s,t}(y) \quad \text{and} \quad X_{r,s}(x-) < y \leq X_{r,s}(x).$$

By the cocycle property ($X_{r,t} = X_{s,t} \circ X_{r,s}$) and as $X_{s,t}$ is non-decreasing then

$$X_{r,t}(x-) = X_{s,t}(X_{r,s}(x-)) \leq X_{s,t}(y-) < z \quad \text{and} \quad X_{r,t}(x) = X_{s,t}(X_{r,s}(x)) \geq X_{s,t}(y) \geq z,$$

thus z at time t is a descendant of x at time r . Similar computations can be written if $X_{s,t}$ is continuous at point y and/or $X_{r,s}$ is continuous at point x .

Recall Condition (1.6) for the sample paths of the CSBP(Ψ) to have infinite variations. This condition ensures the subordinator $X_{s,t}$ to be driftless, i.e. $d_r = 0$ for all $r \geq 0$ in (1.10). As a result, under (1.6), the range $X_{s,t}([0, \infty))$ of the subordinator has zero Lebesgue measure, ensuring that almost every individual x at time t belongs to one of the infinite families of ancestors at time s . This assumption (1.6) often simplifies the interpretation of results obtained in this article. Under this assumption, we denote by $J_{s,t} := \{x \geq 0 : X_{s,t}(x) \neq X_{s,t}(x-)\}$ the set of jumps of $X_{s,t}$. By definition of the genealogy, almost surely the population at time t , indexed by \mathbb{R}_+ , can be partitioned according to their ancestor at time s by $\{(X_{s,t}(y-), X_{s,t}(y)], y \in J_{s,t}\}$.

Recall that according to Theorem 1.A-(ii), Grey's condition $\int^\infty \frac{du}{\Psi(u)} < \infty$ entails that for any $t > 0$, $\ell_t((0, \infty)) < \infty$. Under this condition, the subordinators $X_{s,t}$ are therefore compound Poisson processes. In particular, the set $J_{s,t}$ is the set of atoms of a Poisson process with intensity $v_t(\infty)$. Note that the partition $\{(X_{s,t}(y-), X_{s,t}(y)], y \in J_{s,t}\}$ consists in a family of consecutive intervals. This justifies the introduction of consecutive coalescents on \mathbb{N} in Section 4

2 The inverse flow

We start this section by a preliminary observation on the genealogy backward in time of a CSBP. Consider the Poisson point process on $\mathbb{R}_+ \times (0, \infty)$

$$\mathcal{E}_t(dx, dz) = \sum_{u \geq 0} \delta_{(a_t u, \Delta X_{-t,0}(u))} \tag{2.1}$$

with some renormalisation constant $a_t > 0$ for all $t > 0$. Recall that ρ is the largest positive root of Ψ and ν the quasi-stationary distribution (1.9) of a subcritical CSBP conditioned on the non-extinction.

Proposition 2.1. *Assume $\int^\infty \frac{du}{\Psi(u)} < \infty$ and set $a_t = 1$ if $\Psi'(0^+) < 0$, $a_t = v_t(\infty)$ if $\Psi'(0^+) \geq 0$. Then*

$$\lim_{t \rightarrow \infty} \mathcal{E}_t = \mathcal{E}_\infty \quad \text{in law, for the topology of weak convergence} \quad (2.2)$$

where \mathcal{E}_∞ is a Poisson point process with intensity respectively $\rho dx \otimes \delta_\infty(dz)$ when $\Psi'(0^+) < 0$, $dx \otimes \delta_\infty(dz)$ when $\Psi'(0^+) = 0$, and $dx \otimes \nu(dz)$ when $\Psi'(0^+) > 0$.

Remark 2.2. In the supercritical case, flows of CSBPs can be renormalized to converge almost-surely. We refer to Duquesne and Labbé [DL14], Grey [Gre74], and Foucart and Ma [FM16]. Since for any time t , $X_{-t,0}$ and $X_{0,t}$ have the same law, we could therefore renormalize in law the size of the descendants at time 0 of x from time $-t$. Typically, $\Delta X_{-t,0}(x)$ is of order exponential in the finite mean case ($|\Psi'(0^+)| < \infty$), and double exponential in the infinite mean case ($|\Psi'(0^+)| = \infty$).

Proof. Under Grey's condition, $v_t(\infty) = \ell_t((0, \infty]) < \infty$, and $x \mapsto X_{-t,0}(x)$ is a compound Poisson process with no drift, namely $d_t = 0$. Therefore, the point process \mathcal{E}_t is a Poisson point process with intensity $\frac{\ell_t(dx)}{a_t}$. From (1.10) observe additionally that for any $q \geq 0$,

$$\int_{[0, \infty]} (1 - e^{-qx}) \frac{\ell_t(dx)}{a_t} = \frac{v_t(q)}{a_t}. \quad (2.3)$$

In the supercritical case ($\Psi'(0^+) < 0$), we have $\lim_{t \rightarrow \infty} v_t(q) = \rho$ for all $q > 0$ (while $v_t(0) = 0$), $a_t = 1$ for all $t > 0$ and $\lim_{t \rightarrow \infty} \ell_t((0, \infty]) = \rho$. Therefore (2.3) shows that $\ell_t(dx)$ converges weakly towards $\rho \delta_\infty(dx)$. As a result, we conclude that \mathcal{E}_t converges in law towards a Poisson point process on $(0, \infty) \times (0, \infty]$ with intensity $\rho dx \otimes \delta_\infty(dz)$.

In the subcritical and critical cases, we have $\lim_{t \rightarrow \infty} v_t(\infty) = 0$, and we set $a_t = v_t(\infty)$. By (1.3) and (1.2), we have $\frac{d}{du} v_t(u) = \frac{\Psi(v_t(u))}{\Psi(u)}$. Therefore

$$\frac{v_t(q)}{v_t(\infty)} = \exp\left(-\int_q^\infty \frac{d}{du} \log(v_t(u)) du\right) = \exp\left(-\int_q^\infty \frac{\Psi(v_t(u))}{v_t(u) \Psi(u)} du\right).$$

Since $\lim_{t \rightarrow \infty} \frac{\Psi(v_t(u))}{v_t(u)} = \Psi'(0^+)$, by monotone convergence we obtain that $\lim_{t \rightarrow \infty} \frac{v_t(q)}{v_t(\infty)} = e^{-\Psi'(0^+) \int_q^\infty \frac{du}{\Psi(u)}}$. In the critical case ($\Psi'(0^+) = 0$), the latter limit equals 1 and we thus from (2.3) that $\frac{\ell_t(dx)}{a_t}$ converges weakly towards δ_∞ . In the subcritical case ($\Psi'(0^+) > 0$), we see that $\frac{\ell_t(dx)}{a_t}$ converges weakly towards the probability measure ν with Laplace transform (1.9). We conclude the convergence of \mathcal{E}_t to the stated limits. \square

Let us describe in details the meaning of the above convergence, for supercritical, critical and subcritical CSBPs. Observe that \mathcal{E}_t encodes information on the individuals at time $-t$ having a large family of descendants at time 0. Thus, (2.2) gives information on the origin of the earliest ancestors of the population at time 0. Depending on the sign of $\Psi'(0^+)$, we have three different behaviours:

- (i) If $\Psi'(0^+) < 0$, a unique ancestor from time $-\infty$, located at an exponential random variable with parameter ρ , which generates all individuals at time 0. This individual is the ancestor of the process.

- (ii) If $\Psi'(0^+) = 0$, then $a_t := v_t(\infty) \xrightarrow{t \rightarrow \infty} 0$ and the whole population at time 0 has a common ancestor, but the backward lineage of this ancestor converges in law as $t \rightarrow \infty$ towards ∞ .
- (iii) If $\Psi'(0^+) > 0$, then the population at time 0 is split into distinct families, each of which coming down from a different ancestor at time $-\infty$.

In the (sub)critical case, individuals from generation $-t$ with descendance at time 0 are located at distance $O(1/v_t(\infty))$ from 0. Proposition 2.1 motivates a more complete study of the ancestral lineages of individuals alive in the population at time 0. Our main aim is to provide an almost-sure description of how $(\mathcal{E}_t, t \geq 0)$ evolves and to get precise information on the sizes of the families.

We now introduce the inverse flow of the flow of subordinators $(X_{s,t}, s \leq t)$ and study some of its properties. We first define, for $s \leq t$ and $y \geq 0$

$$X_{s,t}^{-1}(y) := \inf\{x : X_{s,t}(x) > y\}.$$

The process $X_{s,t}^{-1}$ is the right-continuous inverse of the càdlàg process $X_{s,t}$. Note that the individual $X_{s,t}^{-1}(y)$ is the ancestor alive at time s of the individual y considered at time $t \geq s$. It is therefore a natural process to introduce in order to study the genealogy of a CSBP backwards in time. We call inverse flow the process $(\hat{X}_{s,t}(y), s \leq t, y \geq 0)$ defined for all $s \leq t, y \geq 0$ as follows

$$\hat{X}_{s,t}(y) = X_{-t, -s}^{-1}(y). \quad (2.4)$$

We first list some straightforward properties of inverse flows.

Lemma 2.3. *The following properties hold:*

- (i) *Almost surely, for every $s \leq t$ and $x, y > 0$, we have $\{X_{s,t}(x) > y\} = \{\hat{X}_{-t, -s}(y) < x\}$.*
- (ii) *For every $t \geq 0$, $(\hat{X}_{r,s}, r \leq s \leq t)$ and $(\hat{X}_{r,s}, t \leq r \leq s)$ are independent.*
- (iii) *Almost surely, for every $s \leq t \leq u$, $\hat{X}_{s,u} = \hat{X}_{t,u} \circ \hat{X}_{s,t}$.*
- (iv) *For all $x \geq 0$, $\hat{X}_{0,0}(x) = x = \lim_{t \rightarrow 0} \hat{X}_{0,t}(x)$ in probability.*

Remark 2.4. The convergence in (iv) also holds uniformly on compact sets.

Proof. These results are an immediate consequence of Lemma A.1, which describes well-known properties of right-continuous inverses, and the definition of flow of subordinators. More precisely, the first point is a consequence of Lemma A.1(ii), the third one of Lemma A.1(iii) and the fourth one follows from Lemma A.1(iv) and Definition 1.3(iv).

Finally, the second point follows simply from the fact that for all $a \leq b \leq t$, $\hat{X}_{a,b}$ is measurable with respect to $(X_{r,s}, -t \leq r \leq s)$. Hence, by Definition 1.3(ii), we conclude that (ii) holds. \square

We shall denote $(\hat{X}_t(y), y \geq 0, t \geq 0)$ the flow of inverse subordinators $(\hat{X}_{0,t}(y), y \geq 0, t \geq 0)$. As noted above, it tracks backwards in time the ancestral lineages of the population at time 0. Since individuals are ordered, $\hat{X}_t(y)$ can also be interpreted as the random size of the population at time $-t$ whose descendance at time 0 has size y . Observe that by Lemma 2.3(i) and Definition 1.3(i), we have

$$\forall s \leq t, \forall x, y \geq 0, \mathbb{P}(X_{s,t}(x) > y) = \mathbb{P}(\hat{X}_{s,t}(y) < x). \quad (2.5)$$

The relation (2.5) is known as Siegmund duality. We refer the reader to Siegmund [Sie76] and Clifford and Sudbury [CS85].

Theorem 2.5. *Fix $y > 0$. The process $(\hat{X}_t(y), t \geq 0)$ is a Markov process in $(0, \infty)$. Its semigroup $(Q_t, t \geq 0)$ satisfies for any bounded measurable function f and any $t \geq 0$*

$$\mathbb{E}[Q_t f(\mathfrak{e}_q)] = \mathbb{E}[f(\mathfrak{e}_{v_t(q)})] \quad \text{for all } q > 0 \quad (2.6)$$

where \mathfrak{e}_q and $\mathfrak{e}_{v_t(q)}$ are exponential random variable with parameter q and $v_t(q)$.

Observe that (2.6) characterizes the semigroup Q_t , by identification of the Laplace transforms, as it can be rewritten as: for all $q \geq 0$,

$$\int_0^\infty q e^{-qy} Q_t f(y) dy = \int_0^\infty v_t(q) e^{-v_t(q)y} f(y) dy,$$

therefore $Q_t f$ is the inverse Laplace transform of $q \mapsto \frac{v_t(q)}{q} \int_0^\infty e^{-v_t(q)y} f(y) dy$.

Proof. The cocycle property and the independence, obtained in Proposition 2.3(ii)–(iii) readily entail that $t \mapsto \hat{X}_t(y)$ has the Markov property. Moreover, if $\hat{X}_{0,t}(y) = 0$ then $X_{-t,0}(0) = y > 0$, which is impossible, as $X_{-t,0}(0)$ is the value at time t of a CSBP starting from mass 0, and 0 is an absorbing point for a CSBP. Similarly, $\hat{X}_{0,t}(y) = \infty$ yields that $\lim_{z \rightarrow \infty} X_{-t,0}(z) \leq y$, which is impossible as soon as $X_{-t,0}(z)$ is a non-null subordinator.

Finally, we now turn to the computation of the semigroup of $\hat{X}(y)$, which is obtained through the Siegmund duality. Let \mathfrak{e}_q be an independent exponential random variable with parameter q , we have

$$\mathbb{P}(\hat{X}_t(\mathfrak{e}_q) > x) = \mathbb{P}(X_{-t,0}(x) < \mathfrak{e}_q) = \mathbb{E}[e^{-qX_{-t,0}(x)}] = e^{-xv_t(q)},$$

which implies that (2.6) holds. \square

The above theorem shows that the semigroup of (\hat{X}_t) can be expressed in simple terms when applied to exponential distributions. This will motivate later on the study of the action of the flow \hat{X} on Poisson point processes.

We now observe that the Markov process $t \mapsto \hat{X}_{0,t}(y)$ can be straightforwardly extended as a Markov process on $[0, \infty]$.

Proposition 2.6 (Boundaries and Feller property). *Let $y > 0$ fixed, we denote by $(\hat{X}_t, t \geq 0)$ the Markov process $(\hat{X}_t(y), t \geq 0)$.*

(i) *The boundary 0 is an entrance boundary of $(\hat{X}_t, t \geq 0)$ if and only if $\int_0^\infty \frac{du}{\Psi(u)} < \infty$. In that case, $(Q_t, t \geq 0)$ is extended to $[0, \infty)$ by*

$$Q_t f(0) = \int_0^\infty f(u) v_t(\infty) e^{-uv_t(\infty)} du.$$

Otherwise, we set $Q_t f(0) = f(0)$.

(ii) *The boundary ∞ is an entrance boundary of $(\hat{X}_t, t \geq 0)$ if and only if $\int_0^\infty \frac{du}{|\Psi(u)|} < \infty$. In that case, $(Q_t, t \geq 0)$ is defined over $(0, \infty]$ with*

$$Q_t f(\infty) = \int_0^\infty f(u) v_t(0) e^{-uv_t(0)} du.$$

Otherwise, we set $Q_t f(\infty) = f(\infty)$.

(iii) The semigroup $(Q_t, t \geq 0)$ defined over $[0, \infty]$ is Feller.

Remark 2.7. The Markov processes $(\hat{X}_{0,t}(0), t \geq 0)$ and $(\hat{X}_{0,t}(\infty), t \geq 0)$ have the following interpretations, in terms of the CSBP

(i) The process $(\hat{X}_{0,t}(0), t \geq 0)$, starting from 0 at time 0, represents the smallest individual at generation $-t$ to have descendants at time 0. If $\int_0^\infty \frac{du}{|\Psi(u)|} < \infty$, there is extinction in finite time for the CSBP X (i.e. with positive probability, $X_{-t,0}(x) = 0$). In that case $(\hat{X}_{0,t}(0), t \geq 0)$ is a non-trivial Markov process. If $\int_0^\infty \frac{du}{|\Psi(u)|} = \infty$, there is no extinction in finite time for the CSBP, thus all individuals at time t have descendants at time 0, $(\hat{X}_t(0), t \geq 0) \equiv 0$.

(ii) The process $(\hat{X}_t(\infty), t \geq 0)$, starting from ∞ , represents the smallest individual at generation t with an infinite progeny at time 0. If $\int_0 \frac{du}{|\Psi(u)|} < \infty$, there is explosion in finite time for the CSBP X (i.e. with positive probability, $X_{-t,0}(x) = \infty$). In that case, $(\hat{X}_{0,t}(\infty), t \geq 0)$ is a non-trivial Markov process. If $\int_0 \frac{du}{|\Psi(u)|} = \infty$, there is no explosion in finite time and all individuals at time t have finitely many descendants at time 0. Thus $(\hat{X}_t(\infty), t \geq 0) \equiv \infty$ and $Q_t f(\infty) := f(\infty)$.

Proof. For any fixed time t , $(\hat{X}_t(x), x \in (0, \infty))$ is non-decreasing in x . Therefore $\lim_{x \rightarrow \infty} \hat{X}_t(x) = \hat{X}_t(\infty)$ and $\lim_{x \rightarrow 0} \hat{X}_t(x) = \hat{X}_t(0)$ exist almost-surely in $[0, \infty]$. Recall the duality relation (2.5)

$$\mathbb{P}(\hat{X}_t(y) < x) = \mathbb{P}(y < X_t(x)).$$

The first point for the boundary 0 is obtained as follows. By the duality relation, we see that

$$\mathbb{P}(\hat{X}_t(y) \geq x) = \mathbb{P}(y \geq X_t(x)).$$

By letting y to 0, we have

$$\mathbb{P}(\hat{X}_t(0) \geq x) = \mathbb{P}(X_t(x) = 0) = e^{-xv_t(\infty)}.$$

According to Theorem 1.A-(ii), $\int_0^\infty \frac{du}{|\Psi(u)|} < \infty$ is a necessary and sufficient condition for $v_t(\infty) < \infty$. It remains to justify the formula for $Q_t f(0)$. By using Theorem 2.5 and the facts that in probability, $\lim_{q \rightarrow \infty} e_q = 0$ and $\lim_{q \rightarrow \infty} v_t(q) = v_t(\infty) \in (0, \infty]$, we have for any continuous bounded function f on $[0, \infty)$,

$$Q_t f(0) = \lim_{q \rightarrow \infty} \mathbb{E}(Q_t f(e_q)) = \lim_{q \rightarrow \infty} \mathbb{E}(f(e_{v_t(q)})) = \mathbb{E}(f(e_{v_t(\infty)}))$$

by dominated convergence. We deduce the formula for $Q_t f(0)$. We now prove that the semigroup property holds at 0. By definition of $Q_t f(0)$, we have that

$$Q_{t+s} f(0) = \mathbb{E}(f(e_{v_{t+s}(\infty)})) \quad \text{and} \quad Q_t(Q_s f)(0) = \mathbb{E}(Q_s f(e_{v_t(\infty)})) = \mathbb{E}(f(e_{v_s \circ v_t(\infty)})).$$

Therefore, as $v_{t+s} = v_t \circ v_s$, we complete the proof of (i).

The proof of (ii) follows very similar lines to the proof of (i), and is based on the fact that $\int_0 \frac{du}{|\Psi(u)|} < \infty$ is a necessary and sufficient condition for $v_t(0) > 0$. The expression of $Q_t f(\infty)$ is found using that $\lim_{q \rightarrow 0} e_q = \infty$ in probability. Finally, to prove that the semigroup Q_t extended to $[0, \infty]$ is Feller, we observe that the random map $y \mapsto \hat{X}_t(y)$ jumps only on constant

stretch of $X_{-t,0}$ (being its right-continuous inverse). There is no fixed value in $(0, \infty)$ at which $X_{-t,0}$ is constant and therefore $y \in (0, \infty) \mapsto \hat{X}_t(y)$ has no fixed discontinuities. This entails that for any continuous function f over $[0, \infty]$, $Q_t f$ is continuous at any point $y \in (0, \infty)$. By definition $Q_t f(x) \xrightarrow{x \rightarrow \infty} Q_t f(\infty)$ and $Q_t f(x) \xrightarrow{x \rightarrow 0} Q_t f(0)$. The semigroup maps $C([0, \infty])$ in $C([0, \infty])$ and one only needs to show the pointwise continuity at 0 of $Q_t f$, which follows from Proposition 2.3(iv). \square

We study now the long term behaviour of $(\hat{X}_t, t \geq 0)$ in the critical and subcritical cases. By transience, we mean that $\hat{X}_t(x) \xrightarrow{t \rightarrow \infty} \infty$ a.s. for any $x \in (0, \infty)$.

Proposition 2.8. *Let Ψ be a branching mechanism. We observe that*

- (i) *if Ψ is supercritical, then \hat{X} is positive recurrent with stationary law \mathfrak{e}_ρ ;*
- (ii) *if Ψ is subcritical, then \hat{X} is transient;*
- (iii) *if Ψ is critical, then \hat{X} is transient if and only if $\int_0^{\infty} \frac{u}{\Psi(u)} du < \infty$, otherwise it is null recurrent.*

Remark 2.9. Intuitively, in the subcritical case, for any fixed $a > 0$, individuals below level a living at arbitrarily large time in the past will have no progeny at time 0. Therefore the ancestral lineage of an individual x living at time 0, goes above any fixed level a as time goes to ∞ . This explains the transience. In the critical case, large oscillations can occur when $\int_0^{\infty} \frac{x}{\Psi(x)} dx = \infty$. This latter condition is known see Duhalde et al. [DFM14] to entail that first entrance times of the CSBP have infinite mean, in such case the process $(\hat{X}_t, t \geq 0)$ is null recurrent. Note that if $\Psi(q) = cq^\alpha$ with $1 \leq \alpha \leq 2$ and $c > 0$ then $(\hat{X}_t, t \geq 0)$ is null recurrent if $\alpha = 2$ and transient if $\alpha < 2$.

Proof. We first prove (i). Let $y \in (0, \infty)$. By duality (2.5) and Theorem 1.A-(i)

$$\mathbb{P}(\hat{X}_t(y) < x) = \mathbb{P}(X_t(x) > y) \xrightarrow{t \rightarrow \infty} \mathbb{P}(\text{non-extinction}) = 1 - e^{-\rho x}.$$

Assume now Ψ subcritical or critical. For any Borelian set B and any $p > 0$, set

$$U_p(y, B) := \int_0^{\infty} e^{-pt} \mathbb{P}(\hat{X}_t(y) \in B) dt.$$

Fix $q > 0$, recall $\int_{v_t(q)}^q \frac{dx}{\Psi(x)} = t$ and $v_\infty(q) = 0$. By applying Theorem 2.5, one has

$$\begin{aligned} \mathbb{E}[U_p(\mathfrak{e}_q, B)] &= \int_0^{\infty} U_p(y, B) q e^{-qy} dy = \int_0^{\infty} \int_0^{\infty} v_t(q) e^{-pt} e^{-uv_t(q)} \mathbb{1}_B(u) du dt \\ &= \int_0^{\infty} \mathbb{1}_B(u) du \int_0^q e^{-p \int_x^q \frac{dv}{\Psi(v)}} e^{-ux} \frac{x}{\Psi(x)} dx. \end{aligned}$$

By monotone convergence

$$\lim_{p \rightarrow 0} \uparrow \int_0^{\infty} U_p(y, B) q e^{-qy} dy = \int_0^{\infty} U_0(x, B) q e^{-qx} dx = \int_0^{\infty} \mathbb{1}_B(u) du \int_0^q e^{-ux} \frac{x}{\Psi(x)} dx. \quad (2.7)$$

Set $B = (0, a)$ for $a > 0$, then

$$\int_0^\infty U_0(x, B) q e^{-qx} dx = \int_0^a du \int_0^q e^{-ux} \frac{x}{\Psi(x)} dx = \int_0^q \frac{1 - e^{-ax}}{\Psi(x)} dx.$$

In the subcritical case $\Psi'(0^+) > 0$, therefore $\int_0^q \frac{x}{\Psi(x)} dx < \infty$ and for almost every $x \in (0, \infty)$, one has $0 < U_0(x, B) < \infty$. Since for any $x \leq y$, $\hat{X}_t(x) \leq \hat{X}_t(y)$ then

$$\mathbb{P}(\hat{X}_t(x) < a) \geq \mathbb{P}(\hat{X}_t(y) < a)$$

therefore $U_0(x, B) \geq U_0(y, B)$ and then $0 < U_0(x, B) < \infty$ for all x . We may now invoke Proposition 2.2-(iv') in Gettoor [Get80], by taking the increasing sequence $B_n := (0, n)$. This entails that the process $(\hat{X}_t, t \geq 0)$ is transient. In the critical case, if $\int_0^q \frac{x}{\Psi(x)} dx < \infty$ then the process is transient. If now $\int_0^q \frac{x}{\Psi(x)} dx = \infty$ then by (2.7) for any set B with positive Lebesgue measure, $U_0(x, B) = \infty$ for all x . By Proposition 2.4-(i) in [Get80], we conclude that $(\hat{X}_t, t \geq 0)$ is recurrent. \square

3 The Feller flow

In this section, we investigate the genealogy backwards in time of Feller CSBPs. These are continuous CSBPs with quadratic branching mechanisms of the form $\Psi : q \mapsto \frac{\sigma^2}{2} q^2 - \beta q$, with $\beta \in \mathbb{R}$ and $\sigma^2 \geq 0$. For any fixed x , the Feller CSBP $(X_t(x), t \geq 0)$ with mechanism Ψ can be constructed as the solution of the stochastic differential equation

$$dX_t(x) = \sigma \sqrt{X_t(x)} dB_t + \beta X_t(x) dt, \quad X_0(x) = x$$

where $(B_t, t \geq 0)$ is a Brownian motion. We study here in detail the flow $(X_{s,t}(x), t \geq s, x \geq 0)$ of CSBPs with branching mechanism Ψ and the inverse flow $(\hat{X}_{s,t}(x), t \geq s, x \geq 0)$. Many calculations can be made explicit in this setting, see for instance Pardoux [Par08] for a study of the flow $(X_{s,t}(x), t \geq s, x \geq 0)$.

Note that Ψ is subcritical if $\beta < 0$, critical if $\beta = 0$ and supercritical if $\beta > 0$. Moreover, in the latter case we have $\rho = \frac{2\beta}{\sigma^2}$. Observe also that the differential equation (1.3) can be rewritten

$$\frac{dv_t(\lambda)}{dt} = -v_t(\lambda) \left(\frac{\sigma^2}{2} v_t(\lambda) - \beta \right), \quad \text{with } v_0(\lambda) = \lambda,$$

and it is a simple exercise to solve it into

$$v_t(\lambda) = \begin{cases} \frac{\lambda \beta e^{\beta t}}{\beta + \frac{\lambda \sigma^2}{2} (e^{\beta t} - 1)} & \text{if } \beta \neq 0 \\ \frac{\lambda}{1 + \frac{\sigma^2 \lambda t}{2}} & \text{if } \beta = 0. \end{cases} \quad (3.1)$$

Moreover, one can write $v_t(\lambda) = \int_0^\infty (1 - e^{-\lambda r}) \ell_t(dr)$, by setting

$$\ell_t(dr) = v_t(\infty)^2 e^{-\beta t} e^{-v_t(\infty) e^{-\beta t} r} dr$$

where by definition, $v_t(\infty) = \frac{2\beta}{\sigma^2(1 - e^{-\beta t})} > 0$ for $\beta \neq 0$ and $v_t(\infty) = \frac{2}{t\sigma^2}$ if $\beta = 0$. Observe that in both cases, $\frac{\ell_t}{\ell_t((0, \infty))}$ is an exponential law with parameter $\hat{\beta}_t = v_t(\infty) e^{-\beta t}$, which can be rewritten as

$$\hat{\beta}_t = \begin{cases} \frac{2\beta}{\sigma^2(e^{\beta t} - 1)} & \text{if } \beta \neq 0 \\ \frac{2}{t\sigma^2} & \text{if } \beta = 0. \end{cases}$$

Remark 3.1. We often make a distinction between $\beta \neq 0$ and $\beta = 0$, but it is worth noticing that the functions v_t and $\hat{\beta}_t$ that we defined are continuous at $\beta = 0$.

We now study the law of the inverse Feller flow $(\hat{X}_{s,t}(y), s \leq t, y \geq 0)$, in particular characterizing its marginal distributions as a process in the variable t or y .

Theorem 3.2. *The inverse flow $(\hat{X}_t(x), x \geq 0, t \geq 0)$ is characterized as follows. Setting*

$$\forall t \geq 0, \lambda \geq 0, \hat{v}_t(\lambda) = \frac{\lambda \hat{\beta}_t}{\lambda + \hat{\beta}_t e^{\beta t}},$$

we have

(i) *for any fixed $y \geq 0$, $(\hat{X}_t(y), t \geq 0)$ is a Markov process with semigroup given by*

$$\mathbb{E}[e^{-\lambda \hat{X}_t(y)}] = e^{-y \hat{v}_t(\lambda) - \frac{\sigma^2}{2} \int_0^t \hat{v}_s(\lambda) ds}.$$

(ii) *For any fixed t , $(\hat{X}_t(y), y \geq 0)$ is a subordinator with Laplace exponent \hat{v}_t started from the positive random variable $\hat{X}_t(0)$ whose Laplace transform is $\mathbb{E}[e^{-\lambda \hat{X}_t(0)}] = e^{-\frac{\sigma^2}{2} \int_0^t \hat{v}_s(\lambda) ds}$.*

Remark 3.3. The two-parameter process $(\hat{X}_t(x), t \geq 0, x \geq 0)$ is a flow of continuous-state branching processes with immigration with mechanisms $\hat{\Psi}(q) = \frac{\sigma^2}{2} q^2 + \beta q$ and linear immigration $\hat{\Phi}(q) := \frac{\sigma^2}{2} q$. In particular, $(\hat{X}_t(x) - \hat{X}_t(0), t \geq 0)$ is a Feller CSBP with branching mechanism $\hat{\Psi}$.

Proof. As $x \mapsto X_{-t,0}(x)$ is a subordinator with Laplace exponent

$$v_t(\lambda) = \hat{\beta}_t e^{\beta t} \int_0^\infty (1 - e^{-\lambda r}) \hat{\beta}_t e^{-\hat{\beta}_t r} dr,$$

we obtain that this is in fact a compound Poisson process, with jump rate $\hat{\beta}_t e^{\beta t}$ and exponential jump distribution with parameter $\hat{\beta}_t$. Therefore, writing $(N_x^{(t)}, x \geq 0)$ an homogeneous Poisson process with intensity $\hat{\beta}_t e^{\beta t}$ and $(x_i^{(t)}, i \geq 1)$ i.i.d. exponential random variables with parameter $\hat{\beta}_t$, one can rewrite $X_{-t,0}$ as

$$\forall x \geq 0, X_{-t,0}(x) = \sum_{j=1}^{N_x^{(t)}} x_j^{(t)}. \quad (3.2)$$

We set $(\tau_j^{(t)}, j \geq 1)$ the sequence of inter-arrival times of $(N_x^{(t)}, x \geq 0)$ which are i.i.d. exponential random variables with parameter $\hat{\beta}_t e^{\beta t}$, and we write $M_y^{(t)} = \sup\{n \geq 0 : \sum_{i=1}^n x_i^{(t)} \leq y\}$ for all $y \geq 0$. Since the inter-arrival times $(\tau_j^{(t)}, j \geq 1)$ are exponentially distributed, $(M_y^{(t)}, y \geq 0)$ is a Poisson process. By (3.2), $\hat{X}_{0,t}$ being the right-continuous inverse of $X_{-t,0}$, we get

$$\hat{X}_{0,t}(y) = \sum_{j=1}^{M_y^{(t)}+1} \tau_j^{(t)}. \quad (3.3)$$

Note that we have $\hat{X}_{0,t}(0) > 0$, contrarily to $X_{0,t}(0) = 0$, but that $\hat{X}_{0,t}$ is also a compound Poisson process with exponential jump rate. The construction of $X_{-t,0}$ and $\hat{X}_{0,t}$ are represented on Figure 1.




Figure 1: Inverse of compound Poisson process

Since $\tau_1^{(t)}$ is exponentially distributed with parameter $\hat{\beta}_t e^{\beta t}$, we have by (3.3)

$$\mathbb{E}\left(e^{-\lambda \hat{X}_t(0)}\right) = \mathbb{E}\left(e^{-\lambda \tau_1^{(t)}}\right) = \frac{\hat{\beta}_t}{\hat{\beta}_t + \lambda e^{-\beta t}},$$

and moreover, by straightforward Poisson computations, for all $y \geq 0$

$$\mathbb{E}\left[e^{-\lambda(\hat{X}_t(y) - \hat{X}_t(0))}\right] = \exp\left(-y \frac{\lambda \hat{\beta}_t}{\hat{\beta}_t e^{\beta t} + \lambda}\right) = e^{-y \hat{v}_t(\lambda)}.$$

It remains to verify that

$$\mathbb{E}\left[e^{-\lambda \hat{X}_t(0)}\right] = e^{-\frac{\sigma^2}{2} \int_0^t \hat{v}_s(\lambda) ds}.$$

One easily checks that

$$\hat{v}_t(\lambda) = \frac{\lambda \beta e^{-\beta t}}{\beta + \frac{\lambda}{\sigma^2} (1 - e^{-\beta t})} \text{ for all } t \geq 0$$

and in view of (3.1), the map $t \mapsto \hat{v}_t(\lambda)$ is solution to the differential equation $\frac{d\hat{v}_t(\lambda)}{dt} = -\hat{\Psi}(\hat{v}_t(\lambda))$ with $\hat{v}_0(\lambda) = \lambda$ and where $\hat{\Psi}(q) = \frac{\sigma^2}{2} q^2 + \beta q$. By change of variable, one has

$$\int_0^t \hat{v}_s(\lambda) ds = \int_{\hat{v}_t(\lambda)}^{\lambda} \frac{u}{\hat{\Psi}(u)} du = \frac{2}{\sigma^2} \log\left(\frac{\frac{\sigma^2}{2} \lambda + \beta}{\frac{\sigma^2}{2} \hat{v}_t(\lambda) + \beta}\right)$$

and thus $e^{-\frac{\sigma^2}{2} \int_0^t \hat{v}_s(\lambda) ds} = \frac{\frac{\sigma^2}{2} \hat{v}_t(\lambda) + \beta}{\frac{\sigma^2}{2} \lambda + \beta}$. Simple calculations yield that both quantities $\frac{\frac{\sigma^2}{2} \hat{v}_t(\lambda) + \beta}{\frac{\sigma^2}{2} \lambda + \beta}$ and $\frac{\hat{\beta}_t}{\hat{\beta}_t + e^{\beta t} \lambda}$ are equal to $\frac{\beta}{\beta + \lambda \frac{\sigma^2}{2} (1 - e^{-\beta t})}$. We deduce that

$$\mathbb{E}\left[e^{-\lambda \hat{X}_t(0)}\right] = \frac{\hat{\beta}_t}{\hat{\beta}_t + e^{\beta t} \lambda} = \frac{\frac{\sigma^2}{2} \hat{v}_t(\lambda) + \beta}{\frac{\sigma^2}{2} \lambda + \beta} = e^{-\frac{\sigma^2}{2} \int_0^t \hat{v}_s(\lambda) ds},$$

which concludes the proof. \square

The inverse Feller flow being itself a flow of subordinators with explicit law, many quantities can be computed explicitly, such as the most recent common ancestor of a population. Picking two individuals $x \leq y$ at time 0, the age $T_{x,y}$ of the most recent common ancestor of x and y is the first time t such that there exists an individual z at generation $-t$ that gave birth to both x and y , or more precisely

$$T_{x,y} = \inf\{t \geq 0 : \hat{X}_{0,t}(x) = \hat{X}_{0,t}(y)\}. \quad (3.4)$$

This definition of most recent common ancestor can naturally be generalized as follows: given A a subset of \mathbb{R}_+ , we set

$$T_A = \inf\{t \geq 0 : \#\{\hat{X}_{0,t}(A)\} = 1\}.$$

However, as the partition of \mathbb{R}_+ , $\cup_{z \geq 0} (\hat{X}_{0,t})^{-1}(\{z\})$ is a partition in intervals, we have

$$T_A = T_{\inf A, \sup A} \quad \text{a.s.}$$

Therefore, obtaining the law of $T_{x,y}$ will be enough to study the genealogy of the Feller flow.

Proposition 3.4. *For any $0 \leq x \leq y \leq z$, we have*

$$\forall t \geq 0, \mathbb{P}(T_{x,y} \leq t) = e^{-\hat{\beta}_t(y-x)},$$

and $T_{x,y}$ and $T_{y,z}$ are independent. In particular, we have

$$\mathbb{P}(T_{x,y} = \infty) = \begin{cases} 1 - e^{\frac{2\beta}{\sigma^2}(y-x)} & \text{if } \beta < 0 \\ 0 & \text{if } \beta \geq 0. \end{cases}$$

Among other things, this proposition proves that the population comes down from a single ancestor in critical or supercritical cases ($\beta \geq 0$), while in the subcritical case, for $\beta < 0$, the population at time 0 can be separated into families with different ancestors at time $-\infty$.

Proof. This result is a consequence of the inverse flow representation of Theorem 3.2. Indeed, for all $x \leq y$ and $\lambda \geq 0$, we have

$$\mathbb{E}[e^{-\lambda(\hat{X}_t(y) - \hat{X}_t(x))}] = e^{-(y-x)\hat{v}_t(\lambda)},$$

thus, letting $\lambda \rightarrow \infty$ we obtain $\mathbb{P}(T_{x,y} \leq t) = e^{-(y-x)\hat{v}_t(\infty)}$. Moreover, we observe that $\hat{v}_t(\infty) = \hat{\beta}_t$, proving the first equation.

By (3.4), and given that $(\hat{X}_{0,t}(y) - \hat{X}_{0,t}(x), t \geq 0)$ and $(\hat{X}_{0,t}(z) - \hat{X}_{0,t}(y), t \geq 0)$ are independent Feller CSBP with mechanism $\hat{\Psi}$, starting from $y - x$ and $z - y$ respectively, we conclude that $T_{x,y}$ and $T_{y,z}$ are independent.

To obtain $\mathbb{P}(T_{x,y} = \infty)$ we compute

$$\lim_{t \rightarrow \infty} \mathbb{P}(T_{x,y} \leq t) = \exp\left(-(y-x) \lim_{t \rightarrow \infty} \hat{\beta}_t\right) = \begin{cases} 1 & \text{if } \beta \geq 0 \\ e^{-(y-x)\frac{-2\beta}{\sigma^2}} & \text{if } \beta < 0, \end{cases}$$

concluding the proof. \square

Remark 3.5. A straightforward consequence of the above coalescent is that for any choice $\{x_1, \dots, x_n\}$ of individuals at generation 0, the coalescent tree of this family of individuals will only consist in binary merging. Indeed, for every pair (x_i, x_{i+1}) of consecutive individuals, their time of coalescence is independent from the time of coalescence of any other pair of consecutive individuals in the population, and has density with respect to the Lebesgue measure. Therefore, almost surely the first coalescing time will consist in the merging of only two neighbours.

Proposition 3.4 readily entails the representation of the genealogical tree of the population at time 0 as a functional of a Poisson point process. The following construction is reminiscent of the comb representation by Lambert and Uribe Bravo [LUB17].

Proposition 3.6. *There exists a Poisson point process N with intensity $dx \otimes \mu(dt)$ on $\mathbb{R}_+ \times (\mathbb{R}_+ \cup \{\infty\})$ where*

$$\mu(dt) = \begin{cases} \frac{2\beta^2}{\sigma^2} \frac{e^{\beta t}}{(e^{\beta t} - 1)^2} dt & \text{if } \beta > 0 \\ \frac{2}{\sigma^2 t^2} dt & \text{if } \beta = 0, \\ \frac{2\beta^2}{\sigma^2} \frac{e^{\beta t}}{(1 - e^{\beta t})^2} dt + \frac{2|\beta|}{\sigma^2} \delta_\infty & \text{if } \beta < 0 \end{cases}$$

such that almost surely, for any $0 \leq x \leq y$ and $t \geq 0$, we have

$$T_{x,y} < t \iff N([x, y] \times [t, \infty]) = 0.$$

In other words, the coalescent time of x and y is given by the position of the largest atom in the point process $N([x, y] \times \cdot)$. In particular, in the critical case ($\beta = 0$), this result recovers the Brownian coalescent point process of Popovic [Pop04, Lemma 4 and Theorem 5]. In the subcritical case ($\beta < 0$) when two individuals have no common ancestor, there are separated by an infinite atom of the point process $N([x, y] \times \cdot)$.

Proof. We observe from Proposition 3.4 that for all $x \leq y \leq z$, $T_{x,y}$ and $T_{y,z}$ are independent and $T_{x,z} \stackrel{\mathcal{L}}{=} \max(T_{x,y}, T_{y,z})$. Moreover, note by definition that $T_{x,z} \geq \max(T_{x,y}, T_{y,z})$ a.s. This yields that for all $x \leq y \leq z$,

$$T_{x,z} = \max(T_{x,y}, T_{y,z}) \quad \text{a.s.} \quad (3.5)$$

We consider the event of probability one for which the above equation is true simultaneously for all $x, y, z \in \mathbb{Q}_+$. As a result, the field $T_{x,y}$ is decreasing in x and increasing in y . Therefore, there exists a càdlàg modification of the field satisfying (3.5) simultaneously for all $x, y, z \in \mathbb{R}_+$.

As a result, we can construct a simple point process G on $\mathbb{R}_+ \times (\mathbb{R}_+ \cup \{\infty\})$ satisfying

$$T_{x,y} < t \iff G([x, y] \times [t, \infty]) = 0,$$

via the construction $G = \sum_{z \geq 0} \mathbf{1}_{\{\exists \epsilon > 0: T_{z-\epsilon, z} < T_{z, z-\epsilon}\}} \delta_{z, T_{z-\epsilon, z}}$. The point process G is simple (i.e. each atom in the point process has mass one). Moreover, setting γ_t the derivative of $t \mapsto \hat{\beta}_t$, we have

$$\begin{aligned} \mathbb{P}(G([x, y] \times [t, \infty]) = 0) &= \mathbb{P}(T_{x,y} < t) = \exp\left(-\int_t^\infty \int_x^y \gamma_s dz ds - \int_x^y \hat{\beta}_\infty ds\right) \\ &= \mathbb{P}(N([x, y] \times [t, \infty]) = 0), \end{aligned}$$

where N is a Poisson point process with intensity $dx \otimes (\gamma_t dt + \hat{\beta}_\infty \delta_\infty(dt))$. Hence, by monotone classes theorem, for all measurable relatively compact set $B \subset \mathbb{R}_+ \times (\mathbb{R}_+ \cup \{\infty\})$, we have

$$\mathbb{P}(G(B) = 0) = \mathbb{P}(N(B) = 0).$$

As a result, by [Kal02, Theorem 10.9(i)], we have $N \stackrel{\mathcal{L}}{=} G$, which concludes the proof. \square

Pitman and Yor [PY82, Sections 3 and 4], see also [DL14] for a more general setting, have shown that any flow of Feller's branching diffusions can be represented through a Poisson point process on $(0, \infty) \times \mathbb{C}$, where \mathbb{C} denotes the space of continuous paths on \mathbb{R}_+ . In our setting, this entails that the flow $(\hat{X}_t(x) - \hat{X}_t(0), t \geq 0)$ can be represented as follows: for all $t > 0$,

$$\hat{X}_t(x) = \hat{X}_t(0) + \sum_{\substack{x_i \leq x \\ i \in I}} \hat{X}_t^i$$

where $\mathcal{N} = \sum_{i \in I} \delta_{(x_i, \hat{X}^i)}$ is a PPP with intensity $dx \otimes \mathbf{n}(dX)$ and \mathbf{n} is the so-called cluster measure (see for instance [DL14]). The atoms $(\hat{X}^i, i \in I)$ can be interpreted as the ancestral lineages of the initial individuals $(x_i, i \in I)$. They are independent Feller diffusions with mechanism $\hat{\Psi}$ starting from infinitesimal masses. For any $i \in I$, denote by $\zeta_i := \inf\{t \geq 0; \hat{X}_t^i = 0\}$. The time ζ_i represents a binary coalescence time between two "consecutive" individuals. By definition of \mathbf{n} , for any $t > 0$, $\mathbf{n}(\zeta > t) = \hat{v}_t(\infty)$ and therefore $\sum_{i \in I} \delta_{(x_i, \zeta_i)}$ has the same law as N . We represent the ancestral lineages and their coalescences in Figure 2. Recall also from Remark 2.7 that $\hat{X}_t(0)$ is the first individual from generation $-t$ to have descendants at time 0.




Figure 2: Symbolic representation of ancestral lineages and their binary coalescences

In the subcritical case, $(\beta < 0)$, $(\hat{X}_t(x) - \hat{X}_t(0), x \geq 0, t \geq 0)$ is a flow of supercritical CSBPs. Following Bertoin et al. [BFM08] (see also Pardoux [Par08, Section 7]), one can define the random sequence $(x_n^*, n \geq 1)$ recursively as follows:

$$x_1^* := \inf\{x \geq 0; \hat{X}_t(x) - \hat{X}_t(0) \xrightarrow[t \rightarrow \infty]{} \infty\} \text{ and } x_{n+1}^* := \inf\{x \geq x_n^*; \hat{X}_t(x) - \hat{X}_t(0) \xrightarrow[t \rightarrow \infty]{} \infty\}.$$

The random sequence $(x_n^*, n \geq 1)$ is known as the initial prolific individuals of the flow of supercritical CSBPs $(\hat{X}_t(x) - \hat{X}_t(0), x \geq 0, t \geq 0)$ and corresponds to the jumps times of a Poisson process with intensity $-\frac{2\beta}{\sigma^2}$. Within the framework of inverse flow, the random partition of \mathbb{R}_+ : $([0, x_1^*), [x_1^*, x_2^*), \dots)$ corresponds to current families with distinct common ancestors. Note that the sequence $(x_n^*, n \geq 1)$ is also the sequence of atoms of the point process $N(\cdot \times \{\infty\})$, defined in Lemma 3.6.

We observed in this section that the law of the flow \hat{X} is explicit when X is a Feller flow. When the branching mechanism Ψ is not of the quadratic form, multiple births occur in the population. Thus, when time runs backward, coalescences of multiple lineages should arise. The

law of the inverse flow \hat{X} becomes then more involved. In the next section, we construct a simple class of Markovian coalescents which will allow us to encode easily multiple coalescences in lineages backwards in time. The law of the lineage's location $(\hat{X}_t(x), t \geq 0)$ for a fixed individual $x \geq 0$ is studied further in Section 5.

4 Consecutive coalescents

In this section we study the genealogy of branching processes both forward and backward in time, using random partitions of consecutive integers. We shall see how to define a coalescent process in this framework and that the associated coalescent theory is elementary. In a second time, we apply these results to the genealogy of a population in a continuous-state branching process sampled according to a Poisson point process with intensity λ . In a third time, by making the parameter λ increase to ∞ , we obtain a full description of the genealogical tree of individuals in a CSBP under Grey's condition.

4.1 Consecutive coalescents in continuous-time Galton-Watson processes

In this section, we construct a class of simple Markovian coalescents arising when studying the genealogy backward in time of continuous-time Galton-Watson processes. We begin by introducing the classical notation for coalescent processes on the space of partitions. For a more precise description of that framework, in the context of exchangeable coalescents, we refer to Bertoin's book [Ber06, Chapter 4], from which we borrow our definitions and notation.

Let $n \in \mathbb{N} \cup \{\infty\}$, we denote by $[n] = \{j \in \mathbb{N} : j \leq n\}$ the set of integers smaller or equal to n . We call *consecutive partition* of $[n]$ a collection C of disjoint subsets $\{C_1, C_2, \dots\}$ with consecutive integers (i.e. intervals of $[n]$), such that $\cup_{i \geq 1} C_i = [n]$. Without loss of generality, we will always assume that the subsets of the collection C are ranked in the increasing order of their elements. We denote by \mathcal{C}_n the set of consecutive partitions of $[n]$. Note that any $C \in \mathcal{C}_n$ is characterized by the ranked sequence of its blocks sizes $(\#C_1, \#C_2, \dots)$, as

$$\forall j \in \mathbb{N}, \quad C_j = \{k \in \mathbb{N} : \#C_1 + \dots + \#C_{j-1} < k \leq \#C_1 + \dots + \#C_j\}.$$

For any $i, j \in [n]$, we write $i \stackrel{C}{\sim} j$ if and only if i and j belong to the same block of C . For any $n \in \mathbb{N} \cup \{\infty\}$, we set $0_{[n]} = \{\{1\}, \{2\}, \dots, \{n\}\}$ and $1_{[n]} = \{[n]\}$. We introduce some classical operations on \mathcal{C}_n . For each $k \leq n$ and $C \in \mathcal{C}_n$, we denote by

$$C_{|[k]} = \{C_j \cap [k], j \in \mathbb{N}\},$$

the restriction of C to $[k]$ and

$$\#C_{|[k]} := \#\{j \in \mathbb{N} : C_j \cap [k] \neq \emptyset\},$$

the number of blocks of $C_{|[k]}$. Note that for any $m \leq k$, $(C_{|[k]})_{|[m]} = C_{|[m]} \in \mathcal{C}_m$.

We denote by \mathcal{C}_∞ or $\mathcal{C}_\mathbb{N}$ the set of consecutive partitions of \mathbb{N} . A consecutive partition C of \mathbb{N} has at most one block with infinite size. In such a case, C has finitely many blocks, i.e. $\#C < \infty$, and the block with infinite size is indexed by $\#C$. We define a distance on \mathcal{C}_∞ by setting

$$d(C, C') = \sup\{n \in \mathbb{N} : C_{|[n]} = C'_{|[n]}\}^{-1}.$$

Note that the metric space (\mathcal{C}_∞, d) is compact. We next introduce the coagulation operation. For any $C \in \mathcal{C}_n$ and $C' \in \mathcal{C}_{n'}$ such that $\#C \leq n'$, we define the partition $\text{Coag}(C, C')$ by

$$\text{Coag}(C, C')_j = \bigcup_{i \in C'_j} C_i \quad \text{for any } j \in \mathbb{N}.$$

As each block of $\text{Coag}(C, C')$ is the union of a consecutive sequence of consecutive blocks, one has that $\text{Coag}(C, C') \in \mathcal{C}_n$. Thus, Coag defines an internal composition law on \mathcal{C}_∞ . Observe that the blocks of $\text{Coag}(C, C')$ are coarser than those of C . Moreover for any $C, C' \in \mathcal{C}_\infty$ and $n \geq 1$

$$\text{Coag}(C, C')_{|[n]} = \text{Coag}(C_{|[n]}, C'_{|[n]}) = \text{Coag}(C_{|[n]}, C').$$

The operator Coag is therefore Lipschitz continuous with respect to d from $\mathcal{C}_\infty \times \mathcal{C}_\infty$ to \mathcal{C}_∞ and we see that it is associative. For any partition $C \in \mathcal{C}_n$, $\text{Coag}(C, 0_{[n]}) = C$ and $\text{Coag}(C, 1_{[n]}) = 1_{[n]}$.

We are interested in random consecutive partitions of the following form. Let ν be a probability distribution over $\mathbb{N} \cup \{\infty\}$ and $(Z(i), i \geq 1)$ be a random walk with step distribution ν . Set $N := \inf\{i \geq 1; Z(i) = \infty\}$ and

$$C = ([1, Z(1)], [Z(1) + 1, Z(2)], \dots, [Z(N - 1), \infty]).$$

We call C a ν -random consecutive partition. One has $N = \#C$, and conditionally on $\#C = n$, for some $n \in \mathbb{N} \cup \{\infty\}$, the blocks sizes $(\#C_j, 1 \leq j \leq n - 1)$ are i.i.d. random variables in \mathbb{N} . Let C and C' be respectively a ν -random consecutive partition and a ν' -random consecutive partition. If C and C' are independent, then

$$\#\text{Coag}(C, C')_j = \sum_{i \in C'_j} \#C_i \stackrel{\mathcal{L}}{=} \sum_{i \in C'_1} \#C_i,$$

hence $\text{Coag}(C, C')$ is a ν'' -random consecutive partition where ν'' is the law of the random variable $\sum_{i \in C'_1} \#C_i$. In view of the very particular form of a consecutive partition, it is legitimate to question whether the framework of partitions is needed. However, the use of the operator Coag enables us to encode easily multiple coalescences and to follow closely the theory of exchangeable coalescents and its terminology. This encoding simplifies the main formulas we obtain when studying the genealogy of a continuous-state branching population.

Definition 4.1. A Markov process $(C(t), t \geq 0)$ with values in $\mathcal{C}_\mathbb{N}$ is called consecutive coalescent if its semigroup is given as follows: the conditional law of $C(t + s)$ given $C(t) = C$ is the law of $\text{Coag}(C, C')$ where C' is a $\nu_{s,t}$ -random consecutive where $\nu_{s,t}$ is some probability distribution over $\mathbb{N} \cup \{\infty\}$ which may depend on t and s . A consecutive coalescent is said to be homogeneous if the law of C' depends only on s and standard if $C(0) = 0_{[\infty]}$.

We now recall further well-known material on continuous-time Galton-Watson processes. We refer to Athreya and Ney [AN04, Chapter III] for more details on these processes. Consider a finite measure μ on \mathbb{Z}_+ such that $\mu(1) = 0$. A continuous-time Galton-Watson process $(Z_t(n), t \geq 0)$ with reproduction measure μ , is a Markov process counting the number of individuals in a random population with n ancestors where all individuals behave independently, and each individual has an exponential lifetime ζ with parameter $\mu(\mathbb{Z}_+)$ and begets at its death

a random number of children with probability distribution $\mu/\mu(\mathbb{Z}_+)$. The process $(Z_t(n), t \geq 0)$ is characterized in law by μ and thus by the function

$$\psi(x) = - \sum_{k=0}^{\infty} (x^k - x)\mu(k), \quad x \in [0, 1].$$

The process $(Z_t(n), t \geq 0)$ satisfies the branching property

$$\forall n, m \geq 0, \forall t \geq 0, Z_t(n+m) \stackrel{\mathcal{L}}{=} Z_t(n) + \tilde{Z}_t(m) \quad \text{for any } n, m \in \mathbb{Z}_+ \quad (4.1)$$

where $(\tilde{Z}_t(m), t \geq 0)$ is a continuous-time Galton-Watson process independent of $(Z_t(n), t \geq 0)$, and with the same law as $(Z_t(m), t \geq 0)$. This entails that the generating function of $Z_t(n)$ for any $t \geq 0$ has the form

$$\mathbb{E}[x^{Z_t(n)}] = u_t(x)^n, \quad x \in [0, 1], n \in \mathbb{Z}_+$$

where for all $t \geq 0$, $u_t(s)$ is the solution of $\int_{u_t(s)}^s \frac{dz}{\psi(z)} = t$ for any $t \geq 0$. When μ has no mass at 0, the process is called *immortal*. Each individual has at least two children and $(Z_t(n), t \geq 0)$ is non-decreasing in time.

Following the same procedure as in Definition 1.3 but in discrete-state space, we construct a family of continuous-time branching processes by considering a flow of random walks $(Z_{s,t}(n), t \geq s, n \geq 1)$ satisfying the following properties:

- (i) for any $s \leq t$, $(Z_{s,t}(n), n \geq 0)$ is a continuous-time random walk whose jump law has support included in \mathbb{N} and generating function u_{t-s} .
- (ii) For every $t_1 < t_2 < \dots < t_p$, the random walks $(Z_{t_i, t_{i+1}}, i < p)$ are independent and satisfy

$$\forall n \geq 0, Z_{t_1, t_p}(n) = Z_{t_{p-1}, t_p} \circ \dots \circ Z_{t_1, t_2}(n).$$

- (iii) For any $n \geq 1$ and $s \in \mathbb{R}$, $(Z_{s, t+s}(n), t \geq 0)$ is an immortal homogeneous continuous-time Galton-Watson process started from n individuals.

We now construct a flow of partitions describing the genealogy of an immortal continuous-time Galton-Watson process constructed via this flow of random walks. For any $s \leq t$, set

$$\vec{C}(s, t) := ([|Z_{s,t}(i-1) + 1, Z_{s,t}(i)|], i \geq 1),$$

with $Z_{s,t}(0) = 0$. Then by (ii), for any $r < s < t$,

$$\vec{C}(r, t) = \text{Coag}(\vec{C}(s, t), \vec{C}(r, s))$$

i.e. $\vec{C}_j(r, t) = \bigcup_{i \in \vec{C}_j(r, s)} \vec{C}_i(s, t)$ and by definition $\#\vec{C}_i(s, t) = Z_{s,t}(i) - Z_{s,t}(i-1)$ for any $s \leq t$ and any $i \geq 1$. We introduce the time-reversed flow of partition by defining for all $s \leq t$,

$$C(s, t) = \vec{C}(-t, -s).$$

We sum up the main properties of C in the following proposition.

Proposition 4.2. *The stochastic flow of consecutive partitions $(C(s, t), -\infty \leq s \leq t \leq \infty)$ satisfies:*

(i) *For any $s \leq u \leq t$*

$$C(s, t) = \text{Coag}(C(s, u), C(u, t)) \text{ a.s.}$$

(ii) *If $s_1 < s_2 < \dots < s_n$, the partitions $C(s_1, s_2), \dots, C(s_{n-1}, s_n)$ are independent.*

(iii) *The random variables $(\#C_i(s, t), i \geq 1)$ are valued in \mathbb{N} and i.i.d.*

(iv) *$C(0, 0) = 0_{[\infty]}$ and $C(s, t) \rightarrow 0_{[\infty]}$ when $t - s \rightarrow 0$ and*

(v) *the random variable $C(s, t)$ has the same law as $C(0, t - s)$.*

The Markov process $(C(t), t \geq 0)$ defined by $C(t) := C(0, t)$ for any $t \geq 0$ is an homogeneous standard consecutive coalescent in the sense of Definition 4.1. Note that by (i), for any $s, t \geq 0$,

$$C(t + s) = \text{Coag}(C(t), C(t, t + s))$$

namely for any $j \geq 1$,

$$C_j(t + s) = \bigcup_{i \in C_j(t, t + s)} C_i(t), \quad (4.2)$$

so that consecutive blocks are merging as time runs.

Remark 4.3. One can readily check from (4.2) that for any $i \geq 1, s \geq 0$ and $t \geq 0$

$$\#C_i(t + s) = \sum_{m=1}^{\#C_i(t, t + s)} \#C_{m + \sum_{k=1}^{i-1} \#C_k(t, t + s)}(t). \quad (4.3)$$

Processes satisfying (4.3) have been studied in discrete time by Grosjean and Huillet [GH16].




Figure 3: Monotone labelling of an immortal Galton-Watson forest and its consecutive coalescent

By Proposition 4.2-(v), for any fixed $t \geq 0$, $C(t) \stackrel{\mathcal{L}}{=} \vec{C}(t)$. Therefore the coalescent process $(C(t), t \geq 0)$ is characterized by the reproduction measure μ of the associated continuous-time Galton-Watson process $(\# \vec{C}_1(t), t \geq 0)$. Moreover, note that by construction, for any $m \leq n$

$$(C_{|[m]}(t), t \geq 0) = ((C_{|[n]}(t))_{|[m]}, t \geq 0).$$

This consistency property ensures that the family of jump rates of $(C_{|[n]}(t), t \geq 0)$ characterizes the law of $(C(t), t \geq 0)$. In the next lemma, the coagulation rate of a consecutive coalescent restricted to $[n]$ is provided.

Lemma 4.4 (Law of the n -coalescent). *Let $n \in \mathbb{N}$ and $C \in \mathcal{C}_n$. Set $\#C = m$ and assume $C_{|[n]}(0) = C$. For any $j \in [m-1]$, consider the consecutive partitions of $[m]$*

- $C_{\text{in}}^{j,k} := (\{1\}, \dots, \{j, \dots, j+k-1\}, \dots, \{m\})$ for any $2 \leq k \leq m-j$, and attach to each $C_{\text{in}}^{j,k}$ an independent exponential clock with parameter $\mu(k)$,
- $C_{\text{out}}^j := (\{1\}, \dots, \{j, \dots, m\})$, and attach to each C_{out}^j an independent exponential clock with parameter $\bar{\mu}(m-j+1)$

where $\bar{\mu}(k) := \sum_{j=k}^{\infty} \mu(j)$ for any $k \in \mathbb{N}$. Then the process jumps from the partition $C_{|[n]}(t-)$ to $\text{Coag}(C_{|[n]}(t-), D)$ with D the partition in $\{C_{\text{in}}^{j,k}, C_{\text{out}}^j\}$ associated with the first random clock that rings.

Proof. For any $n \geq 1$, any $t \geq 0$,

$$\text{Coag}(C(t), C(t, t+s))_{|[n]} = \text{Coag}(C_{|[n]}(t), C(t, t+s)).$$

Moreover, by associativity of the operator Coag , the restricted process $(C_{|[n]}(t), t \geq 0)$ starting from C , whose number of blocks is m , has the same law as the process $(\text{Coag}(C, C_{|[m]}(t)), t \geq 0)$ where $(C_{|[m]}(t), t \geq 0)$ is the restriction at $[m]$ of the standard process started from $0_{[\infty]}$. Therefore, we only need to focus on the jump rates of the standard coalescent.

For any j , the rate at which the process jumps from $0_{[m]}$ to $C_{\text{out}}^j := (\{1\}, \dots, \{j, \dots, m\})$ is therefore given by

$$\lim_{s \rightarrow 0^+} \frac{1}{s} \mathbb{P}(\#C_j(t, t+s) \geq m-j+1).$$

Since $\#C_j(t, t+s)$ has the same law as the random variable $Z_s(1)$ where $(Z_t(1), t \geq 0)$ is a continuous-time Galton-Watson process with reproduction measure μ , then the latter limit is $\bar{\mu}(m-j+1)$.

Similarly, for any $k \leq m-j-1$, the rate at which the process jumps from $0_{[m]}$ to $C_{\text{in}}^{j,k}$ is

$$\lim_{s \rightarrow 0^+} \frac{1}{s} \mathbb{P}(\#C_j(t, t+s) = k) = \mu(k).$$

There are no simultaneous births forward in time and therefore no simultaneous coalescences. \square

By letting n and m to ∞ in Lemma 4.4, we see that the coalescences in the consecutive coalescent process C valued in \mathcal{C}_{∞} can be described in the following way: to each block j of C is associated a family $(e_{j,k}, k \geq 2)$ of exponential clocks, that ring at rate $(\mu(k), k \geq 2)$. Each time a clock $e_{j,k}$ rings, the consecutive blocks $j, j+1, \dots, j+k-1$ coalesce into one. Note

that these clocks could also be used to construct the immortal Galton–Watson process forward in time: each time the clock $e_{j,k}$ rings, the j th individual produces k children.

We now take interest in the number of blocks of a consecutive coalescent. Similarly to the continuous-state space, the dual process \hat{Z} is defined for any $n \in \mathbb{N}$ and any $t \geq 0$ by

$$\hat{Z}_t(n) := \min\{k \in \mathbb{N} : Z_{-t,0}(k) \geq n\}.$$

The process \hat{Z} is a Markov process, and for all $n \in \mathbb{N}$, $\hat{Z}_t(n)$ is the ancestor at time $-t$ of the individual n considered at time 0.

Proposition 4.5. *For any $t \geq 0$, and any $n, m \in \mathbb{N}$, $\#C_{[n]}(t) = \hat{Z}_t(n)$ and*

$$n \stackrel{C(t)}{\sim} m \iff \hat{Z}_t(n) = \hat{Z}_t(m). \quad (4.4)$$

Proof. By definition

$$C_{[n]}(t) = ([1, Z_{-t}(1)], [Z_{-t}(1) + 1, Z_{-t}(2)], \dots, [Z_{-t}(a-1) + 1, n])$$

with $a = \#C_{[n]}(t) = \min\{k \in \mathbb{N} : Z_{-t}(k) \geq n\} =: \hat{Z}_t(n)$. Consider now an integer $m \leq n$. If $\hat{Z}_t(m) = a$ then $m \in [Z_{-t}(a-1) + 1, n] = C_a(t) \cap [n]$ and $m \stackrel{C(t)}{\sim} n$. The rates of jumps in $(\#C_{[n]}(t), t \geq 0)$ are readily obtained by Lemma 4.4. \square

Remark 4.6. Consecutive coalescents can be defined for a measure μ with a mass at 0 from the relation (4.4). However the process $(C(t), t \geq 0)$ in this case is inhomogeneous in time. We mention that the process $(\hat{Z}_t(n), t \geq 0)$ is studied by Li et al. in [LPLG08].

4.2 Consecutive coalescents in CSBPs through Poisson sampling

We now explain how consecutive coalescents arise in the study of the backward genealogy of CSBPs. Loosely speaking, exchangeable bridges in the theory of exchangeable coalescents, [BLG03], are replaced by subordinators and the sequence of uniform random variables by the atoms of a Poisson process with intensity λ . The following typical random consecutive partitions form a particular class of ν -random consecutive partitions and will play a similar role as paintboxes for exchangeable coalescents.

Definition 4.7. We call (λ, ϕ) -Poisson box a random consecutive partition C obtained by setting

$$i \stackrel{C}{\sim} j \iff X^{-1}(J_i) = X^{-1}(J_j),$$

where X is a subordinator with Laplace exponent ϕ and $(J_j, j \geq 1)$ are the ranked atoms of an independent Poisson process with intensity λ .

The (λ, ϕ) -Poisson boxes will occur as typical random partitions in genealogical trees of CSBPs. More precisely, in the coalescent process describing the genealogy of individuals sampled according to a Poisson point process, the partitions will be distributed as (λ, ϕ) -Poisson boxes. The following Lemma is proved in Appendix A.2, and can be thought of as a revisiting of Pitman’s discretization of subordinators [Pit97].




Figure 4: Construction of a random Poisson-box partition C , with subordinator X and selected individuals $(J_j, j \geq 1)$, satisfying $C_{[7]} = \{\{1\}, \{2\}, \{3, 4, 5\}, \{6\}, \{7\}\}$

Lemma 4.8. *Consider a subordinator X with Laplace exponent*

$$\phi : \mu \mapsto d\mu + \int_0^\infty (1 - e^{-\mu x}) \ell(dx)$$

and $(J_k, k \geq 1)$ the atoms of an independent Poisson process with intensity λ . Let C be the (λ, ϕ) -Poisson box constructed with X and $(J_k, k \geq 1)$ and set for any $i \geq 1$, $J'_i := X^{-1}(J_k)$ for $k \in C_i$. Then

(i) C is a ν -random consecutive partition with for any $k \geq 1$

$$\nu(k) := \frac{1}{\phi(\lambda)} \int_0^\infty \frac{(\lambda x)^k}{k!} e^{-\lambda x} \ell(dx) + d\mathbf{1}_{\{k=1\}} = (-1)^{k-1} \frac{\lambda^k \phi^{(k)}(\lambda)}{k! \phi(\lambda)},$$

i.e. $\mathbb{E}(s^{\#C_1}) = 1 - \frac{\phi(\lambda(1-s))}{\phi(\lambda)}$ for all $s \in [0, 1]$.

(ii) The sequence $(J'_i, i \geq 1)$ is the atoms of a Poisson process with intensity $\phi(\lambda)$.

(iii) $(J'_i, i \geq 1)$ and C are independent.

Remark 4.9. We shall also consider killed subordinators such that $\phi(0) = \ell(\{\infty\}) = \kappa > 0$. The above Lemma can be extended to this case (see Corollary A.4). The associated (λ, ϕ) -Poisson box has finitely many blocks and the sequence $(J'_i, 1 \leq i \leq \#C)$ forms the first atoms of a Poisson process with intensity $\phi(\lambda)$. Formulas in (i) still hold true for any finite k and ν has positive mass at ∞ , namely $\nu(\infty) = \mathbb{P}(\#C_1 = \infty) = 1 - \frac{\phi(0)}{\phi(\lambda)}$.

We now construct consecutive coalescent processes related to the genealogy of the flow of subordinators $(X_{s,t}(x), s \leq t, x \geq 0)$. Denote by $(J_i^\lambda, i \geq 1)$ the sequence of atoms of an independent Poisson process with intensity λ . For any $t \geq 0$, we define $C^\lambda(t)$ as

$$i \stackrel{C^\lambda(t)}{\sim} j \text{ if and only if } \hat{X}_t(J_i^\lambda) = \hat{X}_t(J_j^\lambda). \quad (4.5)$$

The next theorem describes the law of the process $(C^\lambda(t), t \geq 0)$.

Theorem 4.10. *For any $\lambda > 0$, the partition-valued process $(C^\lambda(t), t \geq 0)$ is a consecutive coalescent. Its semigroup is Feller and its one-dimensional marginal law is characterized by*

$$\mathbb{E}[z^{\#C_1^\lambda(t)}] = 1 - \frac{v_t(\lambda(1-z))}{v_t(\lambda)} \text{ for any } z \in [0, 1]. \quad (4.6)$$

There are no simultaneous coalescences and for any $k \geq 2$, the rate at time t at which k given consecutive blocks coalesce is

$$\mu_t^\lambda(k) := \frac{\sigma^2}{2} v_t(\lambda) 1_{\{k=2\}} + v_t(\lambda)^{k-1} \int_{(0, \infty)} \frac{x^k}{k!} e^{-v_t(\lambda)x} \pi(dx). \quad (4.7)$$

In the supercritical case, by choosing for intensity $\lambda = \rho$, the process $(C^\rho(t), t \geq 0)$ becomes time-homogeneous. Corollary 4.11 is obtained by a direct application of Theorem 4.10 since $v_t(\rho) = \rho$ for any $t \geq 0$.

Corollary 4.11. *Assume Ψ supercritical and take $\lambda = \rho$. The coalescent process $(C^\rho(t), t \geq 0)$ is homogeneous in time and the coagulation rate of k given consecutive blocks is*

$$\mu^\rho(k) := \frac{\sigma^2}{2} \rho 1_{\{k=2\}} + \rho^{k-1} \int_{(0, \infty)} \frac{x^k}{k!} e^{-\rho x} \pi(dx).$$

Remark 4.12. Bertoin et al. [BFM08] have shown that in any flow of supercritical CSBPs one can embed an immortal continuous-time Galton-Watson process counting the so-called prolific individuals, whose lines of descent are infinite. The prolific individuals are located in \mathbb{R}_+ as the atoms of a Poisson process with intensity ρ at any time. Moreover, this continuous-time Galton-Watson process has reproduction measure μ^ρ . The consecutive coalescent $(C^\rho(t), t \geq 0)$ represents its genealogy backward in time.

We prove Theorem 4.10. We stress that by definition, from (4.5), $C^\lambda(t)$ is a (λ, v_t) -Poisson box and $C^\lambda(0) = 0_{[\infty]}$ since $\hat{X}_0 = \text{Id}$. Our first lemma proves that the partition-valued process $(C^\lambda(t), t \geq 0)$ is Markovian in its own filtration and is a consecutive coalescent (possibly inhomogeneous in time) in the sense of Definition 4.1.

Lemma 4.13. *For any $s, t \geq 0$*

$$C^\lambda(t+s) = \text{Coag}(C^\lambda(t), C^\lambda(t, t+s)) \quad (4.8)$$

where $C^\lambda(t, t+s)$ is a $(v_t(\lambda), v_s)$ -Poisson box which is independent of $C^\lambda(t)$.

Proof. For any $s, t \geq 0$ and all $l \geq 1$, set $J_l^\lambda(t) := \hat{X}_t(J_l^\lambda)$ for all $i \in C_l^\lambda(t)$. Let $C^\lambda(t, t+s)$ the random consecutive partition defined by

$$l \stackrel{C^\lambda(t, t+s)}{\sim} k \text{ if and only if } \hat{X}_{t, t+s}(J_l^\lambda(t)) = \hat{X}_{t, t+s}(J_k^\lambda(t)).$$

Then by the key lemma 4.8-(ii), $C^\lambda(t, t+s)$ is a $(v_t(\lambda), v_s)$ -Poisson box which is independent of $C^\lambda(t)$. Recall the cocycle property $\hat{X}_{t+s} = \hat{X}_{t, t+s} \circ \hat{X}_t$ (Theorem 2.5-i). Let $i, j \in \mathbb{N}$. Set k and l such that $i \in C_k^\lambda(t)$ and $j \in C_l^\lambda(t)$. By the cocycle property, $\hat{X}_{t, t+s}(J_k^\lambda(t)) = \hat{X}_{t, t+s}(J_l^\lambda(t))$ holds if and only if $i \stackrel{C^\lambda(t+s)}{\sim} j$ and (4.8) holds by definition of the operator Coag , see (4.2). \square

The generating function of the block's size at time t , given in (4.6), is obtained by a direct application of the key lemma 4.8 since $C^\lambda(t)$ is a (λ, v_t) -Poisson box. We now show that the semigroup satisfies the Feller property.

Lemma 4.14. *The process $(C^\lambda(t), t \geq 0)$ is Feller and for any $t \geq 0$, $C^\lambda(t, t+s) \xrightarrow{s \rightarrow 0} 0_{[\infty]}$ in probability.*

Proof. The Feller property corresponds to the continuity of the map

$$C \in \mathcal{C}_\infty \mapsto P_t^\lambda \varphi(C) := \mathbb{E}[\varphi(\text{Coag}(C, C^\lambda(t)))]$$

for any continuous function φ from \mathcal{C}_∞ to \mathbb{R}_+ . This is clear since Coag is Lipschitz continuous. We now show the weak continuity of the semigroup. By definition $J_i^\lambda(t) = \hat{X}_t(J_k)$ for any $k \in C^\lambda(t)$ and for any $i \neq j$, $J_i^\lambda(t) \neq J_j^\lambda(t)$. By Lemma 2.3-(ii) and independence between $(J_i, i \geq 1)$ and \hat{X} , we see that $(J_i^\lambda(t), i \geq 1)$ is independent of $\hat{X}_{t,t+s}$. By Lemma 2.3-(iv), since $\hat{X}_{t,t+s}(x) \xrightarrow{s \rightarrow 0} x$ uniformly on compact sets, in probability, then for any n ,

$$\mathbb{P}(\forall i \neq j \in [n], \hat{X}_{t,t+s}(J_i^\lambda(t)) \neq \hat{X}_{t,t+s}(J_j^\lambda(t))) \xrightarrow{s \rightarrow 0} 1.$$

Therefore $\mathbb{P}(d(C^\lambda(t, t+s), 0_{[\infty]}) \leq 1/n) \xrightarrow{s \rightarrow 0} 1$. □

We now seek for the coagulation rate (4.7).

Lemma 4.15. *For any $z \in (0, 1)$,*

$$\frac{1}{s} \mathbb{E}[z^{\#C_1^\lambda(t, t+s)} - z] \xrightarrow{s \rightarrow 0} \varphi_t^\lambda(z) := \frac{\Psi(v_t(\lambda)(1-z)) - (1-z)\Psi(v_t(\lambda))}{v_t(\lambda)}.$$

Proof. Let $z \in (0, 1)$, since by Lemma 4.8, the random variables $(J_l^\lambda(t), l \geq 1)$ are the atoms of an independent Poisson process with intensity $v_t(\lambda)$, then

$$\mathbb{E}(z^{\#C_1^\lambda(t, t+s)}) = 1 - \frac{v_s(v_t(\lambda)(1-z))}{v_s(v_t(\lambda))}. \quad (4.9)$$

Thus

$$\begin{aligned} \frac{1}{s} \mathbb{E}[z^{\#C_1^\lambda(t, t+s)} - z] &= \frac{1}{s} \left[\frac{(1-z)v_{t+s}(\lambda) - v_s(v_t(\lambda)(1-z))}{v_{t+s}(\lambda)} \right] \\ &= \frac{1}{s} \frac{(1-z)(v_{t+s}(\lambda) - v_t(\lambda)) + v_t(\lambda)(1-z) - v_s(v_t(\lambda)(1-z))}{v_{t+s}(\lambda)} \\ &\xrightarrow{s \rightarrow 0} \frac{\Psi(v_t(\lambda)(1-z)) - (1-z)\Psi(v_t(\lambda))}{v_t(\lambda)} =: \varphi^\lambda(t). \end{aligned}$$

The latter convergence holds since $(v_t(\lambda), t \geq 0)$ solves (1.3) and $v_{t+s} = v_s \circ v_t$. □

By letting $\theta = v_t(\lambda)$ in the next technical lemma, we see that for any $t \geq 0$, the measures μ_t^λ on \mathbb{N} defined in (4.7) have generating function φ_t^λ .

Lemma 4.16. Recall Ψ in (1.4). For any $z \in (0, 1)$ and any $\theta \geq 0$,

$$\frac{\Psi(\theta(1-z)) - (1-z)\Psi(\theta)}{\theta} = \sum_{k=2}^{\infty} (z^k - z)p_{\theta}(k)$$

with $p_{\theta}(k) = \frac{\sigma^2}{2}\theta 1_{\{k=2\}} + \int_0^{\infty} \frac{\theta^{k-1}x^k}{k!} e^{-\theta x} \pi(dx)$.

Proof. It is easy to see that $\Psi(\theta(1-z)) = \Psi_{\theta}(-\theta z) + \Psi(\theta)$, where

$$\Psi_{\theta}(u) := \Psi(u + \theta) - \Psi(\theta) = \Psi'(\theta)u + \frac{\sigma^2}{2}u^2 + \int_0^{\infty} (e^{-ux} - 1 + ux)e^{-\theta x} \pi(dx).$$

Then we have that

$$\begin{aligned} \Psi_{\theta}(-\theta z) &= -\Psi'(\theta)\theta z + \frac{\sigma^2}{2}(\theta z)^2 + \int_0^{\infty} (e^{\theta z x} - 1 - \theta z x)e^{-\theta x} \pi(dx) \\ &= -\Psi'(\theta)\theta z + \frac{\sigma^2}{2}(\theta z)^2 + z \int_0^{\infty} \sum_{k=2}^{\infty} \frac{\theta^k x^k}{k!} e^{-\theta x} \pi(dx) \\ &\quad + \theta \int_0^{\infty} \sum_{k=2}^{\infty} \frac{\theta^{k-1} x^k}{k!} (z^k - z) e^{-\theta x} \pi(dx) \\ &= -\Psi'(\theta)\theta z + \frac{\sigma^2}{2}\theta^2 z + z \int_0^{\infty} (e^{\theta x} - 1 - \theta x) e^{-\theta x} \pi(dx) \\ &\quad + \theta \sum_{k=2}^{\infty} (z^k - z) p_{\theta}(k) \\ &= -z\Psi(\theta) + \theta \sum_{k=2}^{\infty} (z^k - z) p_{\theta}(k). \end{aligned}$$

The last equality follows from the fact that

$$\Psi'(\theta) = -\beta + \sigma^2\theta + \int_0^1 x(1 - e^{-\theta x})\pi(dx) - \int_1^{\infty} x e^{-\theta x} \pi(dx).$$

Thus we have $\Psi(\theta(1-z)) = (1-z)\Psi(\theta) + \theta \sum_{k=2}^{\infty} (z^k - z)p_{\theta}(k)$ for any $z \in (0, 1)$. \square

We now explain how coalescences take place in the process $(C^{\lambda}(t), t \geq 0)$. By construction, the laws of $(C_{[n]}^{\lambda}(t), t \geq 0)$ for $n \geq 1$ are consistent and as in Lemma 4.4 the family of jump rates of $(C_{[n]}^{\lambda}(t), t \geq 0)$ characterizes the law of $(C^{\lambda}(t), t \geq 0)$. The following lemma is obtained along the same lines as Lemma 4.4 but in an inhomogeneous time setting.

Lemma 4.17. Let $n \geq 1$. The n -coalescent process $(C_{[n]}^{\lambda}(t), t \geq 0)$ has jump rates characterized by μ_t^{λ} and the coalescence events are as follows. Fix a time t . Conditionally given $\#C_{[n]}(t-) = m$, for any $j \leq m-1$, consider the consecutive partitions of $[m]$

- $C_{\text{in}}^{j,k} := (\{1\}, \dots, \{j, \dots, j+k-1\}, \dots, \{m\})$ for any $2 \leq k \leq m-j$ and attach to each C_{in}^j a random clock $\zeta_{\text{in}}^{j,k}$ with law

$$\mathbb{P}(\zeta_{\text{in}}^{j,k} > s) = \exp\left(-\int_0^s \mu_r^{\lambda}(k) dr\right).$$

- $C_{\text{out}}^j := (\{1\}, \dots, \{j, \dots, m\})$ and attach to each C_{out}^j a random clock $\zeta_{\text{out}}^{j,k}$ with law

$$\mathbb{P}(\zeta_{\text{out}}^{j,k} > s) = \exp\left(-\int_0^s \bar{\mu}_r^\lambda(m-j+1)dr\right).$$

Then the process jumps from the partition $C_{[n]}(t-)$ to $\text{Coag}(C_{[n]}(t-), D)$ with D the partition in $\{C_{\text{in}}^{j,k}, C_{\text{out}}^j\}$ associated with the first random clock that rings.

Proof of Theorem 4.10. It follows directly by combination of Lemmas 4.13–4.17. \square

We provide now some basic properties of the consecutive coalescent $(C^\lambda(t), t \geq 0)$.

Proposition 4.18. *Fix $\lambda > 0$. If Ψ is critical or supercritical then $(C^\lambda(t), t \geq 0)$ converges almost-surely towards the partition $\mathbb{1}_{\mathbb{N}}$. If Ψ is subcritical, then the process $(C^\lambda(t), t \geq 0)$ converges almost-surely towards a partition $C^\lambda(\infty)$, whose law is characterized by*

$$\mathbb{E}[z^{\#C_1^\lambda(\infty)}] = 1 - e^{-\Psi'(0^+) \int_{\lambda(1-z)}^\lambda \frac{du}{\Psi(u)}} \text{ for any } z \in (0, 1).$$

In this case, the individuals $(J_1^\lambda, J_2^\lambda, \dots)$ belong to different families with i.i.d sizes distributed as $\#C_1^\lambda(\infty)$.

Proof. Recall that for any $t \geq 0$, $\frac{d}{du}v_t(u) = \frac{\Psi(v_t(u))}{\Psi(u)}$. Therefore

$$\frac{v_t(\lambda(1-z))}{v_t(\lambda)} = \exp\left(\int_\lambda^{\lambda(1-z)} \frac{d}{du} \log(v_t(u))du\right) = \exp\left(\int_\lambda^{\lambda(1-z)} \frac{\Psi(v_t(u))}{v_t(u)} \frac{du}{\Psi(u)}\right).$$

If $\Psi'(0^+) < 0$, then $\frac{\Psi(v_t(u))}{v_t(u)} \xrightarrow{t \rightarrow \infty} \Psi(\rho)/\rho = 0$ and by monotone convergence $\frac{v_t(\lambda(1-z))}{v_t(\lambda)} \xrightarrow{t \rightarrow \infty} 1$. By Theorem 4.10, we have

$$\mathbb{E}[z^{\#C_1^\lambda(t)}] = 1 - \frac{v_t(\lambda(1-z))}{v_t(\lambda)} \xrightarrow{t \rightarrow \infty} 0.$$

The process $(\#C_1^\lambda(t), t \geq 0)$ is non-decreasing and thus converges almost-surely in $\bar{\mathbb{N}}$. Therefore $\#C_1^\lambda(t) \xrightarrow{t \rightarrow \infty} \infty$ a.s. Recall that $(C^\lambda(t), t \geq 0)$ converges almost surely towards $\mathbb{1}_{\mathbb{N}}$ if and only if $C_{[n]}^\lambda(t) = \mathbb{1}_{[n]}$ for large enough t . Since $\#C_1(t) \xrightarrow{t \rightarrow \infty} \infty$, then $\mathbb{P}(\#C_1(t) \geq n) = \mathbb{P}(T_{1,n} \leq t) \rightarrow 1$ with $T_{1,n}$ the coalescence time of the ancestral lineages of J_1^λ and J_n^λ . If $\Psi'(0^+) \geq 0$, then $\frac{\Psi(v_t(u))}{v_t(u)} \xrightarrow{t \rightarrow \infty} \Psi'(0^+)$ and by monotone convergence $\frac{v_t(\lambda(1-z))}{v_t(\lambda)} \xrightarrow{t \rightarrow \infty} e^{-\Psi'(0^+) \int_{\lambda(1-z)}^\lambda \frac{du}{\Psi(u)}}$. By Theorem 4.10, we have for any $i \geq 1$,

$$\mathbb{E}[z^{\#C_i^\lambda(t)}] = 1 - \frac{v_t(\lambda(1-z))}{v_t(\lambda)} \xrightarrow{t \rightarrow \infty} 1 - e^{-\Psi'(0^+) \int_{\lambda(1-z)}^\lambda \frac{du}{\Psi(u)}}.$$

By monotonicity, $\#C_1^\lambda(t) \xrightarrow{t \rightarrow \infty} \#C_1^\lambda(\infty)$ a.s. Therefore for large enough time t_1 , for $t \geq t_1$, $C_1^\lambda(t) = C_1^\lambda(\infty)$. Since there is no coalescence between blocks C_1^λ and C_2^λ after time t_1 , the process $(\#C_2^\lambda(t), t \geq t_1)$ is non-decreasing and converges almost-surely towards $\#C_2^\lambda(\infty)$. By induction, for any n_0 , there exists t_{n_0} such that for any $t \geq t_{n_0}$, $\#C_i^\lambda(t) = \#C_i^\lambda(\infty)$ for all $i \leq n_0$. Thus, for any $t \geq t_{n_0}$, $C_i^\lambda(t) \cap [n_0] = C_i^\lambda(\infty) \cap [n_0]$, and then $d(C^\lambda(t), C^\lambda(\infty)) \leq \frac{1}{n_0}$. \square

The following proposition is a direct consequence of the strong law of large numbers.

Proposition 4.19 (Singletons). *For any $t \geq 0$, and any $\lambda > 0$ there are infinitely many singleton blocks at time t and*

$$\frac{\#\{i \in [n]; \#C_i^\lambda(t) = 1\}}{n} \xrightarrow[n \rightarrow \infty]{} D_t^\lambda \text{ a.s.}$$

with $D_t^\lambda = \frac{\lambda}{\Psi(\lambda)} \frac{\Psi(v_t(\lambda))}{v_t(\lambda)}$. This represents the proportion of ancestors that have not been involved in coalescences by time t .

We have seen in Theorem 1.A-(iv) and Proposition 2.6 that when $\int_0 \frac{dx}{|\Psi(x)|} < \infty$, the CSBP explodes and ∞ is an entrance boundary of $(\hat{X}_t, t \geq 0)$. Recall that for any $t > 0$, $\hat{X}_t(\infty)$ is the first individual from generation t to have an infinite progeny at time 0.

Proposition 4.20 (Coming down from infinity). *For any $t > 0$, $\#C^\lambda(t) < \infty$ a.s if and only if $\int_0 \frac{dx}{|\Psi(x)|} < \infty$. Moreover*

$$v_t(0)\#C^\lambda(t) \xrightarrow[t \rightarrow 0]{} \mathfrak{e}_{1/\lambda} \text{ in law}$$

where $\mathfrak{e}_{1/\lambda}$ is an exponential random variable with parameter $1/\lambda$.

Proof. Recall that $v_t(0) > 0$ if and only if $\int_0 \frac{dx}{|\Psi(x)|} < \infty$. By Theorem 4.10, for any $i \geq 1$, $\mathbb{P}(\#C_i^\lambda(t) = \infty) = \frac{v_t(0)}{v_t(\lambda)}$ and therefore the number of blocks $\#C^\lambda(t)$ is a geometric random variable with parameter $\frac{v_t(0)}{v_t(\lambda)}$. For any fixed $x > 0$, one has

$$\mathbb{P}(v_t(0)\#C^\lambda(t) > x) = \left(1 - \frac{v_t(0)}{v_t(\lambda)}\right)^{\lfloor \frac{x}{v_t(0)} \rfloor} \xrightarrow[t \rightarrow 0]{} e^{-\frac{x}{\lambda}}. \quad \square$$

4.3 Backward genealogy of the whole population

In the previous section, we have defined some coalescent processes arising from sampling initial individuals along a Poisson process with an arbitrary intensity λ . The consecutive coalescents obtained by this procedure are only approximating the backward genealogy. They give the genealogy of a random sample of the population. The objective of this subsection is to observe that when the Grey's condition holds, one can define a consecutive coalescent matching with the complete genealogy of the population from any positive time. In all this section, assume the Grey's condition

$$\int_0^\infty \frac{dx}{\Psi(x)} < \infty.$$

Heuristically, we make $\lambda \rightarrow \infty$ in Theorem 4.10, to study the genealogy of the whole population. The limiting process would indeed characterize the genealogy of the CSBP as in this case, an everywhere dense sub-population would be sampled and its genealogy given, which is enough to deduce the genealogical relationship between any pair of individuals. However, this method cannot work directly as one would have jump rates that may explode.

Fix a time $s > 0$. The subordinator $(X_{-s,0}(x), x \geq 0)$ is a compound Poisson process with Lévy measure $\ell_s(dx)$ independent of $(X_{-t,-s}(x), x \geq 0, t \geq s)$. Let $(J_i^{v_s(\infty)}, i \geq 1)$ be the jump times of $(X_{-s,0}(x), x \geq 0)$. They are atoms of a Poisson process with intensity

$v_s(\infty) = \ell_s([0, \infty])$, independent of $(\hat{X}_{s,t}, t \geq s)$. Consider $(C(s, t), t \geq s)$ the partition-valued process defined by

$$i \stackrel{C(s,t)}{\sim} j \text{ iff } \hat{X}_{s,t}(J_i^{v_s(\infty)}) = \hat{X}_{s,t}(J_j^{v_s(\infty)}).$$

The process $(C(s, t), t > s)$ provides a dynamical description of the genealogy of initial individuals whose most recent common ancestors are found at time $s > 0$. The following theorem is a direct application of Theorem 4.10.

Theorem 4.21. *For any $s > 0$, the flow of random consecutive partitions $(C(u, t), t \geq u \geq s)$ satisfies for any $t \geq u \geq s$,*

$$C(s, t) = \text{Coag}(C(s, u), C(u, t)) \text{ a.s.} \quad (4.10)$$

where $C(u, t)$ is a Poisson box with parameters $(v_u(\infty), v_{t-u})$ independent of $C(s, u)$. Moreover for any $i \geq 1$, $t \geq s$ and $z \in (0, 1)$,

$$\mathbb{E}[z^{\#C_i(s,t)}] = 1 - \frac{v_{t-s}(v_s(\infty))(1-z)}{v_t(\infty)}$$

and the consecutive coalescent $(C(s, t), t > s)$ has coagulation rates $(\mu_t^\infty, t > s)$ with

$$\mu_t^\infty(k) := \frac{\sigma^2}{2} v_t(\infty) 1_{\{k=2\}} + v_t(\infty)^{k-1} \int_{(0,\infty)} \frac{x^k}{k!} e^{-v_t(\infty)x} \pi(dx) \text{ for any } k \geq 2. \quad (4.11)$$

We see in the next corollary that by reversing time in any block of the consecutive coalescent $(C(s, t), 0 < s \leq t)$, one obtains an inhomogeneous continuous-time Galton-Watson process. Fix an horizon time $T > 0$ and consider the consecutive partitions $C(T-t, T)$ for any $t \in [0, T[$.

Corollary 4.22. *The processes $(Z_i^T(t), 0 \leq t < T) := (\#C_i(T-t, T), 0 \leq t < T)$ are i.i.d inhomogeneous continuous-time Galton-Watson processes. For any $z \in [0, 1]$, and any $t \in [0, T[$*

$$\mathbb{E}[z^{Z_i^T(t)}] = 1 - \frac{v_t(v_{T-t}(\infty))(1-z)}{v_T(\infty)}. \quad (4.12)$$

Moreover, denoting by γ_i^T , the time of its first jump, one has for any $t \in [0, T[$

$$\mathbb{P}(\gamma_i^T > t) = \frac{\Psi(v_T(\infty))}{v_T(\infty)} \frac{v_{T-t}(\infty)}{\Psi(v_{T-t}(\infty))}.$$

Proof. The law of $Z_i^T(t)$ for fixed t is obtained by a direct application of Theorem 4.21. Only remains to show the branching property. By (4.10) for any s and t such that $0 \leq t + s < T$

$$C(T - (t + s), T) = \text{Coag}(C(T - (t + s), T - t), C(T - t, T))$$

which provides $C_i(T - (t + s), T) = \bigcup_{j \in C_i(T-t, T)} C_j(T - (t + s), T - t)$ and the branching property. \square

Remark 4.23. The process $(Z_1^T(t), 0 \leq t < T)$ corresponds to the *reduced* Galton-Watson process obtained by Duquesne and Le Gall [DLG02, Theorem 2.7.1] in the (sub)critical case. We refer also to Fekete et al. [FFK17] for an approach with stochastic differential equations. In the supercritical case, since for any $t \geq 0$, $v_{T-t}(\infty) \xrightarrow{T \rightarrow \infty} \rho$, we see in (4.12) that $(Z_1^T(t), t \geq$

0) converges, as T goes to infinity, in the finite-dimensional sense, towards a Markov process $(Z^\infty(t), t \geq 0)$ whose semigroup satisfies for any $z \in (0, 1)$

$$\mathbb{E}[z^{Z^\infty(t)}] = 1 - \frac{v_t(\rho(1-z))}{\rho}.$$

Namely, $(Z^\infty(t), t \geq 0)$ is a continuous-time Galton-Watson process, homogeneous in time, with reproduction measure μ^ρ . Heuristically, individuals from time $-t$ with descendants at time T will correspond at the limit with prolific individuals.

The coalescent process $(C(s, t), t \geq s)$ only describes coalescence in families from time $s > 0$. We define now a coalescent process from time 0 by using the flow of subordinators. Denote by $\mathcal{C}_{\mathbb{R}_+}$ the space of partitions of $(0, \infty)$ into consecutive half-closed intervals. That is to say, partitions of the form $\mathcal{C} = ((0, x_1], (x_1, x_2], \dots)$ for some non-decreasing sequence of positive real numbers $(x_i, i \geq 1)$. The space of consecutive partitions of \mathbb{N} , $(\mathcal{C}_\infty, \text{Coag})$, acts as follows on $\mathcal{C}_{\mathbb{R}_+}$: for any $\mathcal{C} \in \mathcal{C}_{\mathbb{R}_+}$ and $C \in \mathcal{C}_\infty$, for any $i \geq 1$

$$\text{Coag}(\mathcal{C}, C)_i = \bigcup_{j \in C_i} \mathcal{C}_j$$

where $\mathcal{C}_j = (x_{j-1}, x_j]$ and $x_0 = 0$. The following theorem achieves one of our goals and has to be compared with our preliminary observation in Proposition 2.1. It describes completely the genealogy backwards in time as well as the sizes of asymptotic families. For any $t > 0$, denote by J_{-t} the set of jumps of the subordinator $(X_{-t,0}(x), x \geq 0)$.

Theorem 4.24. *Define the process $(\mathcal{C}(t), t > 0)$ valued in $\mathcal{C}_{\mathbb{R}_+}$ as follows:*

$$\mathcal{C}(t) = \{(X_{-t,0}(x-), X_{-t,0}(x)], x \in J_{-t}\}.$$

The process $(\mathcal{C}(t), t > 0)$ is a time-inhomogeneous Markov process such that for any $t \geq s > 0$,

$$\mathcal{C}(t) = \text{Coag}(\mathcal{C}(s), C(s, t)) \text{ a.s.}$$

In the critical or supercritical case, $\mathcal{C}(t) \xrightarrow[t \rightarrow \infty]{} \mathbb{1}_{(0, \infty)}$ a.s. In the subcritical case, $\mathcal{C}(t) \xrightarrow[t \rightarrow \infty]{} \mathcal{C}(\infty)$ a.s and the length of a typical interval at the limit has for law the quasi-stationary distribution of the CSBP conditioned on the non-extinction:

$$\mathbb{E}[e^{-u|\mathcal{C}_1(\infty)|}] = 1 - \exp\left(-\Psi'(0^+) \int_u^\infty \frac{dv}{\Psi(v)}\right).$$

Proof. Recall that $X_{-t,0} = X_{-s,0} \circ X_{-t,-s}$ and $\hat{X}_t = \hat{X}_{s,t} \circ \hat{X}_s$. This entails that for any $x \in J_{-t} = \{J_j^{v_t(\infty)}, j \geq 1\}$,

$$(X_{-t,0}(x-), X_{-t,0}(x)) = \bigcup_{\substack{y \in (0, \infty) \\ \hat{X}_{s,t}(y) = x}} (X_{-s}(y-), X_{-s}(y)). \quad (4.13)$$

For any $t > 0$ and any $j \geq 1$, set $\mathcal{C}_j(t) = (x_{j-1}(t), x_j(t)) = (X_t(J_j^{v_t(\infty)}-), X_t(J_j^{v_t(\infty)}))$ with $x_0(t) = 0$. By definition of $C(s, t)$ and (4.13), we have

$$\mathcal{C}_i(t) = \bigcup_{j \in C_i(s, t)} \mathcal{C}_j(s).$$

Proposition 4.18 ensures that $(C(s, t), t \geq s)$ converges almost-surely as t goes to ∞ . This entails the almost-sure convergence of $(\mathcal{C}(t), t > 0)$. In the supercritical or critical case, $C(s, t) \xrightarrow{t \rightarrow \infty} 1_{\mathbb{N}}$ and then $\mathcal{C}(t) \xrightarrow{t \rightarrow \infty} \mathbb{1}_{(0, \infty)}$, where $\mathbb{1}_{(0, \infty)}$ denotes the partition of $(0, \infty)$ with only one block. In the subcritical case, Proposition 2.1 entails that for all $i \geq 1$ and $u \geq 0$,

$$\mathbb{E}[e^{-u|\mathcal{C}_i(\infty)|}] = 1 - \exp\left(-\Psi'(0^+) \int_u^\infty \frac{dv}{\Psi(v)}\right).$$

Note moreover that $\mathcal{C}(\infty) = \text{Coag}(\mathcal{C}(s), C(s, \infty))$ for any $s > 0$. \square




Figure 5: Symbolic representation of the genealogy

Remark 4.25. Bertoin and Le Gall in [BLG06, Proposition 3] have shown that in the critical case the Lévy measures $(\ell_t, t > 0)$ solve the following Smoluchowski equation

$$\frac{\partial}{\partial t} \langle f, \ell_t \rangle = v_t(\infty) \sum_{k=2}^{\infty} \mu_t^\infty(k) \int_{(0, \infty)^k} (f(x_1 + \dots + x_k) - (f(x_1) + \dots + f(x_k))) \ell_t(dx_1) \dots \ell_t(dx_k) \quad (4.14)$$

where f is a continuous function on $(0, \infty)$ with compact support and $\langle f, \ell_t \rangle = \int_{(0, \infty)} f(x) \ell_t(dx)$. The process $(\mathcal{C}(t), t > 0)$ sheds some light on this deterministic equation since $\mu_t^\infty(k)$ is the rate in $(\mathcal{C}(t), t > 0)$ at which k given consecutive intervals coagulate and by the strong law of large numbers,

$$\frac{1}{n} \sum_{i=1}^n \delta_{|\mathcal{C}_i(t)|} \xrightarrow{n \rightarrow \infty} \frac{\ell_t(dx)}{v_t(\infty)} \text{ a.s for any } t > 0,$$

where $|\mathcal{C}_i(t)|$ denotes the length of the i^{th} interval in $\mathcal{C}(t)$. We refer the reader to Iyer et al. [ILP15], [ILP18] for recent works on Equation 4.14.

When the CSBP explodes, the individuals in the current generation have finitely many ancestors. The following proposition is the analogue of Proposition 4.20.

Proposition 4.26. *Assume $\int_0^{\frac{du}{\Psi(u)}} < \infty$, then the consecutive coalescent $(\mathcal{C}(t), t > 0)$ comes down from infinity and*

$$\frac{v_t(0)}{v_t(\infty)} \# \mathcal{C}(t) \xrightarrow{t \rightarrow 0} \mathfrak{e} \text{ in law}$$

where \mathfrak{e} is a standard exponential random variable.

Proof. For any $t > 0$, the lengths of the intervals in $\mathcal{C}(t)$ are i.i.d random variables with law $\frac{\ell_t(dx)}{v_t(\infty)}$. Under the assumption, $\int_0^\infty \frac{du}{|\Psi(u)|} < \infty$, $\ell_t(\{\infty\}) = v_t(0) > 0$ and therefore the number of intervals in $\mathcal{C}(t)$ has a geometric law with parameter $\frac{v_t(0)}{v_t(\infty)}$. The convergence in law is proved by a similar calculation as in Proposition 4.20. \square

We saw in Proposition 4.5 that the number of blocks in $(C_{[n]}(t), t \geq 0)$, the coalescent process associated with a continuous-time Galton-Watson process, corresponds to the inverse flow of random walks $(\hat{Z}_t(n), t \geq 0)$ at a fixed level n . Recall that in continuous-state space the process $(\hat{X}_t(x), t \geq 0)$ can then be interpreted as the size of the ancestral population whose descendants at time 0 form a family of size x . The study of its infinitesimal dynamics is more involved than in the discrete setting and is the aim of the next section.

5 A martingale problem for the inverse flow

We investigate the extended generator $\hat{\mathcal{L}}$ of $(\hat{X}_t, t \geq 0)$. Recall that we write \mathcal{L} the generator of the CSBP with mechanism Ψ . As we consider the flow of subordinators over $[0, \infty]$, it is natural to express \mathcal{L} , given in (1.5), as follows. For all function G in C_b^2 the space of twice differentiable bounded functions with bounded derivatives, we have:

$$\mathcal{L}G(x) = \frac{\sigma^2}{2}xG''(x) + \beta xG'(x) + \int_0^\infty \pi(dh) \int_0^\infty du (G(\Delta_{h,u}(x)) - G(x) - h\mathbb{1}_{\{u \leq x\}}G'(x)\mathbb{1}_{\{h \leq 1\}}) \quad (5.1)$$

with $\Delta_{h,u}(x) := x + h\mathbb{1}_{\{x \geq u\}}$. We denote by C_0^2 the space of twice continuously differentiable functions over $[0, \infty)$ which tend to 0 at infinity and whose first derivative tends to 0 at infinity.

Theorem 5.1. *For any function F in C_0^2 , set*

$$\begin{aligned} \hat{\mathcal{L}}F(z) = & \frac{\sigma^2}{2}zF''(z) + \left(\frac{\sigma^2}{2} - \beta z\right)F'(z) \\ & + \int_0^\infty \pi(dh) \int_0^\infty du [F(\psi_{h,u}(z)) - F(z) + h\mathbb{1}_{\{h \leq 1\}}F'(z)\mathbb{1}_{\{z > u\}}] \end{aligned}$$

with

$$\psi_{h,u}(z) := z\mathbb{1}_{[0,u]}(z) + u\mathbb{1}_{[u,u+h]}(z) + (z-h)\mathbb{1}_{[u+h,\infty)}(z).$$

Then for any $y > 0$, $(\hat{X}_t(y), t \geq 0)$ solves the following well-posed martingale problem

$$(MP) \quad \left(F(\hat{X}_t(y)) - \int_0^t \hat{\mathcal{L}}F(\hat{X}_s(y))ds, t \geq 0 \right)$$

is a martingale for any function F in $\mathcal{D} := \{F \in C_0^2; F' \in L^1 \text{ and } \beta xF'(x), \frac{\sigma^2}{2}xF''(x) \xrightarrow{x \rightarrow \infty} 0\}$.

Remark 5.2. Note that $\psi_{h,u} = \Delta_{h,u}^{-1}$ is the right-continuous inverse function of $\Delta_{h,u}$, see Figure 6. For any $y \geq 0$, if individual u has at time t a progeny of size h , then $\psi_{h,u}(y)$ at time $t-$ is the infinitesimal parent of individual y at time t : if $y < u$, then y has no parent but himself, if $y \in [u, u+h]$, the parent of y is $\psi_{h,u}(y) = u$, if $y > u+h$ then its parent is $\psi_{h,u}(y) = y-h$. If $y_1 \neq y_2$ then $\psi_{h,u}(y_1) = \psi_{h,u}(y_2)$ if and only if $y_1, y_2 \in [u, u+h]$.




Figure 6: Graphs of the functions $\Delta_{h,u}$ and $\psi_{h,u}$

The proof of Theorem 5.1 is divided in four lemmas.

Lemma 5.3. *The operator $\hat{\mathcal{L}}$ is well-defined on C_b^2 .*

Proof. Let $F \in C_b^2$. We first study the integral in $\hat{\mathcal{L}}$ with respect to $\pi(dh)$ on the interval $[1, \infty)$. For any $y > 0$,

$$\begin{aligned} \left| \int_1^\infty \pi(dh) \int_0^\infty du (F(\psi_{h,u}(y)) - F(y)) \right| \\ = \left| \int_1^\infty \pi(dh) \int_0^y du (F(\psi_{h,u}(y)) - F(y)) \right| \leq y\bar{\pi}(1)2\|F\|_\infty \end{aligned}$$

where $\bar{\pi}(x) := \int_x^\infty \pi(du)$ for any $x > 0$. In order to study the integral on $[0, 1)$, note that for any $y > 0$, $u > 0$ and $h > 0$, we have

$$\psi_{h,u}(y) - y = (u - y)\mathbb{1}_{[u, u+h]}(y) - h\mathbb{1}_{[u+h, \infty)}(y) = (u - y)\mathbb{1}_{[y-h, y]}(u) - h\mathbb{1}_{[0, y-h]}(u).$$

Assume $h < 1$. We have that

$$\begin{aligned} & |F(\psi_{h,u}(y)) - F(y) + hF'(y)\mathbb{1}_{\{y>u\}}| \\ & \leq |F(\psi_{h,u}(y)) - F(y) - (\psi_{h,u}(y) - y)F'(y)| + |(\psi_{h,u}(y) - y + h\mathbb{1}_{\{y>u\}})F'(y)| \\ & \leq \frac{(\psi_{h,u}(y) - y)^2}{2}\|F''\|_\infty + |\psi_{h,u}(y) - y + h\mathbb{1}_{\{y>u\}}|\|F'\|_\infty. \end{aligned}$$

On the one hand,

$$\begin{aligned} \psi_{h,u}(y) - y + h\mathbb{1}_{\{y>u\}} &= (u - y)\mathbb{1}_{[y-h, y]}(u) - h\mathbb{1}_{[0, y-h]}(u) + h\mathbb{1}_{[0, y]}(u) \\ &= (u + h - y)\mathbb{1}_{[y-h, y]}(u) \geq 0 \end{aligned}$$

and

$$\int_0^\infty (u + h - y)\mathbb{1}_{[y-h, y]}(u)du = \left[\frac{u^2}{2} \right]_{y-h}^y + (h - y)h = \frac{h^2}{2}.$$

On the other hand

$$\frac{(\psi_{h,u}(y) - y)^2}{2} = \frac{1}{2}((u - y)^2\mathbb{1}_{[y-h, y]}(u) + h^2\mathbb{1}_{[0, y-h]}(u))$$

and

$$\int_0^\infty \frac{(\psi_{h,u}(y) - y)^2}{2} du = \int_0^\infty \frac{1}{2} ((u-y)^2 \mathbb{1}_{[y-h,y]}(u) + h^2 \mathbb{1}_{[0,y-h]}(u)) du \leq \frac{h^3}{6} + \frac{h^2}{2} y \leq \frac{h^2}{2} (y+1).$$

Thus, for any $0 \leq h < 1$

$$\int_0^\infty |F(\psi_{h,u}(y)) - F(y) + hF'(y) \mathbb{1}_{\{h < 1\}} \mathbb{1}_{\{y > u\}}| du \leq \frac{h^2}{2} (y+1) \|F''\|_\infty + \frac{h^2}{2} \|F'\|_\infty. \quad (5.2)$$

The integral with respect to $\pi(dh)$ on $(0, 1)$ in $\hat{\mathcal{L}}F$ is therefore convergent and $\hat{\mathcal{L}}$ well-defined. \square

We now follow the same method as Bertoin and Le Gall in [BLG05, Theorem 5] to show that $\hat{\mathcal{L}}$ is an extended generator, i.e. that $(F(\hat{X}_t(u)) - \int_0^t \hat{\mathcal{L}}F(\hat{X}_s(u)) ds, t \geq 0)$ is a martingale for all F in \mathcal{D} . Let g be an integrable continuous function over $[0, \infty)$ and set $G(t) = \int_0^t g(u) du$. Let $F \in \mathcal{D}$ and write $F(t) = \int_t^\infty f(x) dx$ where $f(x) := -F'(x)$. By Fubini's theorem, note that

$$\int_0^\infty \int_0^\infty g(u) f(x) \mathbb{1}_{\{x \geq u\}} du dx = \int_0^\infty g(u) F(u) du = \int_0^\infty f(x) G(x) dx.$$

Moreover, one classically has that

$$\int_0^\infty f(x) \mathbb{P}(\hat{X}_s(u) < x) dx = \mathbb{E}[F(\hat{X}_s(u))] \quad \text{and} \quad \int_0^\infty g(u) \mathbb{P}(X_s(x) > u) du = \mathbb{E}[G(X_s(x))].$$

Recall that by (2.5), we have $\mathbb{P}(\hat{X}_s(u) < x) = \mathbb{P}(X_s(x) > u)$ for all $x, u \geq 0$. Then, integrating this equality with respect to $f(x)g(u) dx du$ provides

$$\int_0^\infty du g(u) \mathbb{E}[F(\hat{X}_s(u)) - F(u)] = \int_0^\infty dx f(x) \mathbb{E}[G(X_s(x)) - G(x)]. \quad (5.3)$$

Therefore, a first step in the search for $\hat{\mathcal{L}}$ is computing the right-hand side of (5.3). Let $\hat{\mathcal{L}}^d$ and $\hat{\mathcal{L}}^c$ be, respectively, the discontinuous part and the continuous one of the generator $\hat{\mathcal{L}}$.

Lemma 5.4. *Let $\lambda > 0$ and $g(x) = e^{-\lambda x}$ for any $x \in \mathbb{R}_+$ then for any $F \in \mathcal{D}$*

$$\int_0^\infty dx f(x) \mathbb{E}[G(X_s(x)) - G(x)] = \int_0^\infty g(u) du \int_0^s dt \mathbb{E} \left[\hat{\mathcal{L}}^d F(\hat{X}_t(u)) + \hat{\mathcal{L}}^c F(\hat{X}_t(u)) \right], \quad (5.4)$$

where $f(x) = -F'(x)$ and $G(x) = \int_0^x g(u) du$.

Proof. The function $x \mapsto G(x) = \frac{1-e^{-\lambda x}}{\lambda}$ is in C_b^2 and by applying Dynkin's formula we get

$$\mathbb{E}[G(X_s(x)) - G(x)] = \int_0^s dt \mathcal{L}P_t G(x) = \int_0^s dt \mathcal{L}^c P_t G(x) + \int_0^s dt \mathcal{L}^d P_t G(x)$$

where we write for all twice derivable function H ,

$$\begin{aligned} \mathcal{L}^c H(x) &= \frac{\sigma^2}{2} x H''(x) + \beta x H'(x) \\ \mathcal{L}^d H(x) &= \int_0^\infty \pi(dh) \int_0^\infty du (H(\Delta_{h,u}(x)) - H(x) - h \mathbb{1}_{\{u \leq x\}} H'(x) \mathbb{1}_{\{h \leq 1\}}) \end{aligned}$$

which denote respectively the continuous and discontinuous parts of the generator \mathcal{L} . We start by studying the discontinuous part. For any $s \geq 0$, we can rewrite

$$\begin{aligned} & \int_0^s dt \mathcal{L}^d P_t G(x) \\ &= \int_0^s dt \int_0^\infty \pi(dh) \int_0^\infty du [P_t G(x + h \mathbb{1}_{\{u < x\}}) - P_t G(x) - h(P_t G)'(x) \mathbb{1}_{\{h < 1\}} \mathbb{1}_{\{u \leq x\}}]. \end{aligned} \quad (5.5)$$

We first compute

$$\begin{aligned} P_t G(x + h \mathbb{1}_{\{u < x\}}) - P_t G(x) &= \mathbb{E}[G(X_t(x + h \mathbb{1}_{\{u < x\}})) - G(X_t(x))] \\ &= \int_0^\infty g(v) dv (\mathbb{P}(v \leq X_t(x + h \mathbb{1}_{\{u < x\}})) - \mathbb{P}(v \leq X_t(x))) \\ &= \int_0^\infty g(v) dv (\mathbb{P}(x + h \mathbb{1}_{\{u \leq x\}} \geq \hat{X}_t(v)) - \mathbb{P}(x \geq \hat{X}_t(v))). \end{aligned}$$

By Lemma A.1(ii) and Remark 5.2, one has $\Delta_{h,u}(x) > y$ if and only if $x > \psi_{h,u}(y)$, therefore

$$P_t G(x + h \mathbb{1}_{\{u < x\}}) - P_t G(x) = \int_0^\infty g(v) dv (\mathbb{P}(\psi_{h,u}(\hat{X}_t(v)) \leq x) - \mathbb{P}(\hat{X}_t(v) \leq x)).$$

Integrating with respect to $f(x)dx$ we obtain

$$\int_0^\infty dx f(x) (P_t G(x + h \mathbb{1}_{\{u < x\}}) - P_t G(x)) = \int_0^\infty g(v) dv (\mathbb{E}[F(\psi_{h,u}(\hat{X}_t(v)))] - F(\hat{X}_t(v))). \quad (5.6)$$

We now compute the compensated part of the discontinuous generator $\hat{\mathcal{L}}^d$. By integration by parts we have

$$\int_0^\infty f(x) (P_t G)'(x) \mathbb{1}_{\{u \leq x\}} dx = -f(u) P_t G(u) - \int_u^\infty f'(x) P_t G(x) dx, \quad (5.7)$$

since $f(\infty) := \lim_{x \rightarrow \infty} f(x) = 0$ and $\lim_{x \rightarrow \infty} P_t G(x) = G(\infty) < \infty$. Moreover, we observe that

$$\begin{aligned} & \int_u^\infty f'(x) P_t G(x) dx \\ &= \int_u^\infty f'(x) \mathbb{E} \left[\int_0^\infty \mathbb{1}_{\{X_t(x) > v\}} g(v) dv \right] dx = \int_u^\infty f'(x) \mathbb{E} \left[\int_0^\infty \mathbb{1}_{\{x > \hat{X}_t(v)\}} g(v) dv \right] dx \\ &= \mathbb{E} \left[\int_0^\infty g(v) dv \int_u^\infty f'(x) \mathbb{1}_{\{x > \hat{X}_t(v)\}} dx \right] = \int_0^\infty g(v) dv (f(\infty) - \mathbb{E}[f(\hat{X}_t(v) \vee u)]) \\ &= - \int_0^\infty \mathbb{E}[f(\hat{X}_t(v) \vee u)] g(v) dv. \end{aligned}$$

Therefore (5.7) becomes

$$\begin{aligned} \int_0^\infty f(x) (P_t G)'(x) \mathbb{1}_{\{u \leq x\}} dx &= -f(u) P_t G(u) + \int_0^\infty \mathbb{E}[f(\hat{X}_t(v) \vee u)] g(v) dv \\ &= \int_0^\infty (\mathbb{E}[f(\hat{X}_t(v) \vee u)] - f(u) \mathbb{P}(\hat{X}_t(v) \leq u)) g(v) dv \\ &= \int_0^\infty \mathbb{E}[f(\hat{X}_t(v)) \mathbb{1}_{\{\hat{X}_t(v) > u\}}] g(v) dv. \end{aligned}$$

Using the above result and (5.6), (5.5) yields

$$\begin{aligned}
& \int_0^\infty dx f(x) \int_0^s dt \mathcal{L}^d P_t G(x) \\
&= \int_0^\infty dx f(x) \int_0^s dt \int_0^\infty \pi(dh) \int_0^\infty du [P_t G(x+h\mathbb{1}_{\{u < x\}}) - P_t G(x) - h(P_t G)'(x)\mathbb{1}_{\{h < 1\}}\mathbb{1}_{\{u \leq x\}}] \\
&= \int_0^s dt \int_0^\infty \pi(dh) \int_0^\infty du \int_0^\infty dx f(x) [P_t G(x+h\mathbb{1}_{\{u < x\}}) - P_t G(x) - h(P_t G)'(x)\mathbb{1}_{\{h < 1\}}\mathbb{1}_{\{u \leq x\}}]
\end{aligned} \tag{5.8}$$

$$\begin{aligned}
&= \int_0^s dt \int_0^\infty \pi(dh) \int_0^\infty du \int_0^\infty g(v) dv \Upsilon(t, h, u, v) \\
&= \int_0^\infty g(v) dv \int_0^s dt \int_0^\infty \pi(dh) \int_0^\infty du \Upsilon(t, h, u, v),
\end{aligned} \tag{5.9}$$

where

$$\Upsilon : (t, h, u, v) \mapsto \left(\mathbb{E}[F(\psi_{h,u}(\hat{X}_t(v)))] - F(\hat{X}_t(v)) - hF'(\hat{X}_t(v))\mathbb{1}_{\{\hat{X}_t(v) > u\}}\mathbb{1}_{\{h \leq 1\}} \right).$$

Above, (5.8) and (5.9) follow from applying Fubini's theorem, which we now justify. For any t and x ,

$$P_t G(x) = \frac{1 - e^{-xv_t(\lambda)}}{\lambda} \text{ and } (P_t G)''(x) = -\frac{v_t(\lambda)^2}{\lambda} e^{-xv_t(\lambda)}.$$

Since $v_t(\lambda)e^{-xv_t(\lambda)} \leq \frac{1}{x}$ then $\sup_{[x, x+h]} |(P_t G)''(z)| \leq \frac{v_t(\lambda)}{\lambda x}$, and by Taylor's inequality

$$|P_t G(x+h\mathbb{1}_{\{u < x\}}) - P_t G(x) - h(P_t G)'(x)\mathbb{1}_{\{h < 1\}}\mathbb{1}_{\{u \leq x\}}| \leq \frac{v_t(\lambda)}{\lambda x} \frac{(h \wedge 1)^2}{2} \mathbb{1}_{\{u \leq x\}}.$$

Since f is integrable then the upper bound is integrable with respect to $f(x)dx\mathbb{1}_{[0,s]}(t)dt\pi(dh)du$, which justifies the application of Fubini's theorem in (5.8).

We now explain why (5.9) holds. Recall first that $f(z) = -F'(z)$. By Theorem 2.5, for any $q > 0$ we have $\mathbb{E}[\hat{X}_t(\mathfrak{e}_q)] = \frac{1}{v_t(q)} < \infty$ therefore $\mathbb{E}[\hat{X}_t(x)] < \infty$ for a.e. x . This, with the bound (5.2) allows us to conclude that $\Upsilon(t, h, u, v)$ is integrable with respect to $g(v)dv\mathbb{1}_{[0,s]}(t)dt\pi(dh)du$.

In a second time, we deal with the continuous part of the generator \mathcal{L}^c . Applying Fubini's theorem, one has

$$\int_0^\infty dx f(x) \int_0^s \mathcal{L}^c P_t G(x) dt = \int_0^s dt \int_0^\infty dx f(x) \mathcal{L}^c P_t G(x).$$

Set $h(x) = P_t G(x)$ and $\phi(x) = f'(x)\frac{\sigma^2}{2}x + f(x)\left(\frac{\sigma^2}{2} - \beta x\right)$. Since by assumption $F \in \mathcal{D}$, then

$\lim_{x \rightarrow \infty} \phi(x) = \phi(\infty) = 0$. We now compute $\int_0^\infty dx f(x) \mathcal{L}^c P_t G(x)$. By two integrations by parts

$$\begin{aligned}
& \int_0^\infty dx f(x) \mathcal{L}^c h(x) = \int_0^\infty dx f(x) \left[\frac{\sigma^2}{2} x h''(x) + \beta x h'(x) \right] \\
& = \left[f(x) \frac{\sigma^2}{2} x h'(x) \right]_0^\infty - \int_0^\infty dx \left[f'(x) \frac{\sigma^2}{2} x + f(x) \frac{\sigma^2}{2} \right] h'(x) + \int_0^\infty dx f(x) \beta x h'(x) \\
& = \left[f(x) \frac{\sigma^2}{2} x h'(x) \right]_0^\infty - \int_0^\infty \phi(x) h'(x) dx \\
& = \left[f(x) \frac{\sigma^2}{2} x h'(x) \right]_0^\infty - \phi(\infty) h(\infty) + \int_0^\infty \phi'(x) h(x) dx \\
& = -\phi(\infty) h(\infty) + \int_0^\infty \phi'(x) \mathbb{E} \left[\int_0^\infty g(u) \mathbb{1}_{\{u \leq X_t(x)\}} du \right] dx \\
& = \int_0^\infty du g(u) \int_0^\infty dx \phi'(x) \mathbb{P}(\hat{X}_t(u) < x) \\
& = - \int_0^\infty du g(u) \mathbb{E}[\phi(\hat{X}_t(u))] = \int_0^\infty du g(u) \mathbb{E}[\hat{\mathcal{L}}^c F(\hat{X}_t(u))].
\end{aligned}$$

We can now conclude as follows. One has

$$\begin{aligned}
\int_0^\infty du g(u) \mathbb{E}[F(\hat{X}_s(u)) - F(u)] &= \int_0^\infty dx f(x) \mathbb{E}[G(X_s(x)) - G(x)] \quad (5.10) \\
&= \int_0^\infty g(v) dv \int_0^s dt \mathbb{E} \left[\hat{\mathcal{L}}^d F(\hat{X}_t(v)) + \hat{\mathcal{L}}^c F(\hat{X}_t(v)) \right]. \quad \square
\end{aligned}$$

Lemma 5.5. *For any $F \in \mathcal{D}$ and any $y \geq 0$, $\left(F(\hat{X}_t(y)) - \int_0^t \hat{\mathcal{L}} F(\hat{X}_s(y)) ds, t \geq 0 \right)$ is a martingale.*

Proof. Recall that $g(v) = e^{-\lambda v}$. We will show that (5.10) entails that for any v and any s :

$$\mathbb{E}[F(\hat{X}_s(v)) - F(v)] = \int_0^s dt \mathbb{E} \left[\hat{\mathcal{L}}^d F(\hat{X}_t(v)) + \hat{\mathcal{L}}^c F(\hat{X}_t(v)) \right]. \quad (5.11)$$

From the Feller property of $(\hat{X}_t, t \geq 0)$ and the continuity of $\hat{\mathcal{L}}F$ for any function F in \mathcal{D} , the map $v \mapsto \mathbb{E}[\hat{\mathcal{L}}F(\hat{X}_t(v))]$ is continuous. For any $a > 0$, and any $v \leq a$, $\hat{X}_t(v) \leq \hat{X}_t(a)$ a.s. therefore using the bound (5.2), we see that the function defined on $[0, a]$ by

$$\Xi(v) = \int_0^s dt \int_0^\infty \pi(dh) \int_0^\infty du \left(\mathbb{E}[F(\psi_{h,u}(\hat{X}_t(v)))] - F(\hat{X}_t(v)) + h F'(\hat{X}_t(v)) \mathbb{1}_{\{\hat{X}_t(v) > u\}} \mathbb{1}_{\{h \leq 1\}} \right)$$

is continuous, and since a is arbitrary, the mapping is continuous on $[0, \infty)$. This corresponds to the continuity of $v \mapsto \int_0^s dt \mathbb{E} \left[\hat{\mathcal{L}}^d F(\hat{X}_t(v)) \right]$. On the other hand, one can check the continuity of $v \mapsto \int_0^s dt \mathbb{E} \left[\hat{\mathcal{L}}^c F(\hat{X}_t(v)) \right]$ and by injectivity of the Laplace transform, (5.10) entails (5.11). This provides the martingale problem, as the following routine calculation shows. Let $t \geq 0$ and

$s \geq 0$. Denote by $(\mathcal{F}_s, s \geq 0)$ the natural filtration associated with $(\hat{X}_s(x), s \geq 0, x \geq 0)$,

$$\begin{aligned} & \mathbb{E} \left[F(\hat{X}_{t+s}(x)) - \int_0^{t+s} \hat{\mathcal{L}}F(\hat{X}_u(x))du \mid \mathcal{F}_s \right] \\ &= \mathbb{E} \left[F(\hat{X}_{t+s}(x)) - \int_s^{t+s} \hat{\mathcal{L}}F(\hat{X}_u(x))du \mid \mathcal{F}_s \right] - \int_0^s \hat{\mathcal{L}}F(\hat{X}_u(x))du \\ &= \mathbb{E}_{\hat{X}_s(x)} \left[F(\hat{X}_t) - \int_0^t \hat{\mathcal{L}}F(\hat{X}_u)du \right] - \int_0^s \hat{\mathcal{L}}F(\hat{X}_u(x))du \\ &= F(\hat{X}_s(x)) - \int_0^s \hat{\mathcal{L}}F(\hat{X}_u(x))du, \end{aligned}$$

where we have used (5.11) in the last equality. \square

In the following Lemma, we rewrite the generator $\hat{\mathcal{L}}$ of the one-point motion in its Courrège form. We refer to Kolokoltsov [Kol11] for a general study of generators of stochastically monotone Markov processes.

Lemma 5.6. *For any $f \in C_b^2$,*

$$\hat{\mathcal{L}}f(z) = \frac{\sigma^2}{2} z f''(z) + \int_0^z [f(z-h) - f(z) + h f'(z)] \nu(z, dh) + b(z) f'(z)$$

with

$$\nu(z, dh) := \mathbb{1}_{\{h \leq z\}} ((z-h)\pi(dh) + \bar{\pi}(h)dh)$$

and

$$b(z) := \int_0^\infty h(z \mathbb{1}_{\{h \leq 1\}} \pi(dh) - \nu(z, dh)) - \beta z + \frac{\sigma^2}{2}.$$

Moreover, the martingale problem (MP) is well-posed.

Remark 5.7. The jump measure $\nu(z, dh)$ can be compared to the jumps rate of $(\hat{Z}_t(n), t \geq 0)$ obtained in Proposition 4.5. Moreover, in the finite mean case, $\int_1^\infty h \pi(dh) < \infty$, the drift b can be rewritten as follows

$$\begin{aligned} b(z) &= \int_0^\infty h(z \pi(dh) - \nu(z, dh)) + \Psi'(0^+)z + \frac{\sigma^2}{2} \\ &= z \int_z^\infty \bar{\pi}(h)dh + \int_0^z h \bar{\pi}(h)dh + \Psi'(0^+)z + \frac{\sigma^2}{2}. \end{aligned}$$

In particular, for any $z > 0$, $b'(z) = \int_z^\infty \bar{\pi}(dh) + \Psi'(0^+)$, $b''(z) = -\bar{\pi}(z)$ and b is concave.

Proof. The continuous part $\hat{\mathcal{L}}^c$ has already the wished form, we thus focus on $\hat{\mathcal{L}}^d$. Recall $\psi_{h,u}(z) := z \mathbb{1}_{\{z \leq u\}} + (z-h) \mathbb{1}_{[u+h, \infty)}(z) + u \mathbb{1}_{[u, u+h]}(z)$. Note that

$$\begin{aligned} & \int_0^\infty \pi(dh) \int_0^\infty du [f(\psi_{h,u}(z)) - f(z) + h \mathbb{1}_{\{h \leq 1\}} f'(z) \mathbb{1}_{\{z > u\}}] \\ &= \int_0^\infty \pi(dh) \int_0^z du [(f(z-h) - f(z)) \mathbb{1}_{[u+h, \infty)}(z) + (f(u) - f(z)) \mathbb{1}_{[u, u+h]}(z) + h \mathbb{1}_{\{h \leq 1\}} f'(z)]. \end{aligned}$$

Therefore, one has

$$\begin{aligned}\hat{\mathcal{L}}^d f(z) &= \int_0^z \pi(dh) \left[(z-h)[f(z-h) - f(z)] + \int_0^z du [f(u) - f(z)] \mathbb{1}_{[u, u+h]}(z) + zh f'(z) \mathbb{1}_{h < 1} \right] \\ &\quad + \int_z^\infty \pi(dh) \left[\int_0^z du [f(u) - f(z)] \mathbb{1}_{[u, u+h]}(z) + hf'(z) \mathbb{1}_{h < 1} \right] \\ &= I + II.\end{aligned}$$

For the first integral I :

$$\begin{aligned}I &= \int_0^z \pi(dh) [(z-h)[f(z-h) - f(z)] + hf'(z) \mathbb{1}_{h < 1}] + h^2 f'(z) \mathbb{1}_{h < 1} \\ &\quad + \int_0^z \pi(dh) \int_0^z du (f(u) - f(z)) \mathbb{1}_{\{u > z-h\}},\end{aligned}$$

by Tonelli's theorem and a change of variables, one has

$$\begin{aligned}\int_0^z \pi(dh) \int_0^z du (f(u) - f(z)) \mathbb{1}_{\{u > z-h\}} &= \int_0^z (f(u) - f(z)) (\bar{\pi}(z-u) - \bar{\pi}(z)) du \\ &= \int_0^z (f(z-h) - f(z)) (\bar{\pi}(h) - \bar{\pi}(z)) dh.\end{aligned}$$

Thus

$$\begin{aligned}I &= \int_0^z [f(z-h) - f(z) + hf'(z) \mathbb{1}_{\{h \leq 1\}}] (z-h) \pi(dh) + \int_0^z (f(z-h) - f(z)) (\bar{\pi}(h) - \bar{\pi}(z)) dh \\ &\quad + \int_0^z h^2 \mathbb{1}_{\{h \leq 1\}} f'(z) \pi(dh) \\ &= \int_0^z [f(z-h) - f(z) + hf'(z) \mathbb{1}_{\{h \leq 1\}}] ((z-h) \pi(dh) + (\bar{\pi}(h) - \bar{\pi}(z)) dh) \\ &\quad + \int_0^z h^2 \mathbb{1}_{\{h \leq 1\}} \pi(dh) f'(z) - \int_0^z h \mathbb{1}_{h < 1} (\bar{\pi}(h) - \bar{\pi}(z)) dh f'(z).\end{aligned}$$

For the second integral II , the change of variables $u = z - v$ in the first integral provides

$$\begin{aligned}II &= \int_z^\infty \pi(dh) \int_0^z du (f(u) - f(z)) \mathbb{1}_{[u, u+h]}(z) + \int_z^\infty \pi(dh) zh f'(z) \mathbb{1}_{h < 1} \\ &= \bar{\pi}(z) \int_0^z dv [f(z-v) - f(z)] + z f'(z) \int_z^\infty \pi(dh) h \mathbb{1}_{h < 1}.\end{aligned}$$

Summing both expressions, $I + II$ equals to:

$$\begin{aligned}&\int_0^z [f(z-h) - f(z) + hf'(z) \mathbb{1}_{\{h \leq 1\}}] ((z-h) \pi(dh) + (\bar{\pi}(h) - \bar{\pi}(z) + \bar{\pi}(z)) dh) \\ &\quad + \left(\int_0^z h^2 \mathbb{1}_{\{h \leq 1\}} \pi(dh) - \int_0^z h \mathbb{1}_{h < 1} (\bar{\pi}(h) - \bar{\pi}(z)) dh \right. \\ &\quad \left. + z \int_z^\infty h \mathbb{1}_{h \leq 1} \pi(dh) - \bar{\pi}(z) \int_0^\infty h \mathbb{1}_{\{h < 1\}} dh \right) f'(z).\end{aligned}$$

Therefore

$$I + II = \int_0^z [f(z-h) - f(z) + hf'(z)\mathbb{1}_{\{h \leq 1\}}] \nu(z, dh) + b_1(z)f'(z)$$

with

$$\begin{aligned} b_1(z) &:= \int_0^z (h^2 \mathbb{1}_{h \leq 1} \pi(dh) - h \bar{\pi}(h) \mathbb{1}_{h \leq 1} dh) + z \int_z^\infty \pi(dh) h \mathbb{1}_{h < 1} \\ &= \int_0^\infty h \mathbb{1}_{\{h \leq 1\}} (z \pi(dh) - \nu(z, dh)), \end{aligned}$$

and we obtain

$$\begin{aligned} I + II &= \int_0^z [f(z-h) - f(z) + hf'(z)] \nu(z, dh) - \int_0^z \mathbb{1}_{\{1 < h \leq z\}} \nu(z, dh) f'(z) + b_1(z) f'(z) \\ &= \int_0^z [f(z-h) - f(z) + hf'(z)] \nu(z, dh) + b(z) f'(z). \end{aligned}$$

We now verify uniqueness of the solution to (MP) by applying Theorem 5.1 of Kolokoltsov [Kol11]. Assumptions (i) and (ii) of the theorem can be readily checked. The third assumption (iii) is that for any $z > 1$, $b(z) \leq c(1+z)$ for some $c > 0$. Let $z > 1$, one has

$$\begin{aligned} b(z) &= \int_0^1 h(z \pi(dh) - \nu(z, dh)) - \int_1^z h \nu(z, dh) - \beta z + \frac{\sigma^2}{2} \\ &= \int_0^1 (h^2 \pi(dh) + h \bar{\pi}(h) dh) - z \int_1^z h \pi(dh) + \int_1^z h^2 \pi(dh) - \int_1^z h \bar{\pi}(h) dh - \beta z + \frac{\sigma^2}{2} \\ &\leq \int_0^1 (h^2 \pi(dh) + h \bar{\pi}(h) dh) + \frac{\sigma^2}{2} - \beta z \leq c(1+z) \end{aligned}$$

where for the first inequality we use the fact that $-z \int_1^z h \pi(dh) + \int_1^z h^2 \pi(dh) \leq 0$ and we choose a large enough c for the second inequality. Finally, since \mathcal{D} contains C_c^2 , the space of twice continuously differentiable functions with compact support, [Kol11, Theorem 5.1] applies. \square

Proof of Theorem 5.1. It follows directly by combination of Lemmas 5.5 and 5.6. \square

Remark 5.8. Similar computations to the ones made in the proof of Lemma 5.4 can be done for the p -point motion $(\hat{X}_t(y_1), \dots, \hat{X}_t(y_p))$ from the duality relation

$$\mathbb{P}(\hat{X}_t(y_1) < x_1, \dots, \hat{X}_t(y_p) < x_p) = \mathbb{P}(X_t(x_1) > y_1, \dots, X_t(x_p) > y_p).$$

Consider for example the case $\sigma = \beta = 0$. For any function F in $C^2(\mathcal{D}_p)$, where we denote by $\mathcal{D}_p := \{y := (y_1, \dots, y_p) \in (0, \infty)^p, y_1 \leq y_2 \leq \dots \leq y_p\}$, we set

$$\hat{\mathcal{L}}F(y) = \int_0^\infty \pi(dh) \int_0^\infty du \left[F(\psi_{h,u}(y)) - F(y) + h \mathbb{1}_{\{h \leq 1\}} \sum_{i=1}^p \frac{\partial}{\partial y_i} F(y) \mathbb{1}_{\{y_i > u\}} \right]$$

with $\psi_{h,u}(y) = (\psi_{h,u}(y_1), \dots, \psi_{h,u}(y_p))$. Then

$$F(\hat{X}_t(y)) - \int_0^t \hat{\mathcal{L}}F(\hat{X}_s(y)) ds$$

is a local martingale, where $\hat{X}_t(y) = (\hat{X}_t(y_1), \dots, \hat{X}_t(y_p))$.

6 Examples

In this section, we apply the results obtained in the previous ones to the two following important examples: the stable CSBP and the Neveu CSBP. These CSBPs arise in many different frameworks and are known for instance to be closely related to the class of exchangeable coalescents called Beta-coalescents.

6.1 Feller and stable CSBPs

A stable CSBP is a continuous-state branching process with branching mechanism given by $\Psi(u) = c_\alpha u^\alpha - \beta u$, for some $\alpha \in (1, 2]$, $c_\alpha > 0$ and $\beta \in \mathbb{R}$. Note in particular that Feller's flow, whose inverse flow was studied in details in Section 3 is a stable CSBP with $\alpha = 2$ and $c_2 = \frac{\sigma^2}{2}$. As a direct application of Theorem 4.24, we obtain that the Markovian coalescent $(\mathcal{C}(t), t > 0)$ associated with the Feller flow has coagulation rates

$$\mu_t^\infty(k) = \frac{2\beta}{\sigma^2(1 - e^{-\beta t})} \delta_2(k).$$

In particular, in the subcritical case ($\beta < 0$), $\mathcal{C}(t)$ converges almost-surely as $t \rightarrow \infty$ towards intervals with i.i.d. exponentially distributed lengths with parameter $\hat{\rho} = 2\beta/c_2$. This corresponds to the partition of \mathbb{R}_+ into random intervals $((0, x_1^*], (x_1^*, x_2^*], \dots)$ corresponding to different ancestors at time $-\infty$ found in Section 3.

We now assume that $\Psi(u) = c_\alpha u^\alpha - \beta u$ for some $\alpha \in (1, 2)$, with $c_\alpha := \frac{\Gamma(2-\alpha)}{\alpha(\alpha-1)}$ (which corresponds to a simple time dilatation). By assumption $\alpha > 1$, Grey's condition $\int^\infty \frac{du}{\Psi(u)} < \infty$ holds. Solving the differential equation (1.3) satisfied by $v_t(\lambda)$, we have in particular that

$$v_t(\infty) = \begin{cases} c_\alpha^{-\frac{1}{\alpha-1}} \left(\frac{1 - e^{-(\alpha-1)\beta t}}{\beta} \right)^{-\frac{1}{\alpha-1}} & \text{if } \beta \neq 0 \\ (\Gamma(2-\alpha)/\alpha)^{-\frac{1}{\alpha-1}} t^{-\frac{1}{\alpha-1}} & \text{if } \beta = 0. \end{cases}$$

For the stable CSBP, the coagulation rates of its associated Markovian coalescent $(\mathcal{C}(t), t > 0)$ are given by

$$\mu_t^\infty(k) := v_t(\infty) \mu_\alpha(k),$$

with $\mu_\alpha(k) := \frac{\Gamma(k-\alpha)}{k!}$. The normalized associated probability measure is

$$\frac{\mu_t^\infty(k)}{\mu_t^\infty(\mathbb{N})} = \frac{\alpha(2-\alpha)\dots(k-1-\alpha)}{k!} \tag{6.1}$$

which is time-independent. This probability distribution corresponds to the reproduction measure of prolific individuals in supercritical stable CSBP, see Example 3 in [BFM08]. It also appears in the study of reduced α -stable trees and Beta($2-\alpha, \alpha$)-exchangeable coalescents, see respectively Duquesne and Le Gall [DLG02, page 74] and Berestycki et al. [BBS07, Section 5].

The inhomogeneous consecutive coalescent $(C(s, t), t > s)$ representing the genealogy of any stable CSBP from time $s > 0$ is obtained by a deterministic time-change of the homogeneous consecutive coalescent $(\check{C}(t), t \geq 0)$ with coagulation rates μ_α via the transformation: for any $t \geq s$,

$$C(s, t) = \check{C} \left(\int_s^t v_u(\infty) du \right).$$

Note that $\int_s^\infty v_u(\infty)du = \int_0^{v_s(\infty)} \frac{z}{\Psi(z)} dz$ which is finite if and only if Ψ is subcritical ($\beta < 0$). According to Theorem 4.24, in the subcritical case $(\mathcal{C}(t), t > 0)$ converges almost-surely as $t \rightarrow \infty$ towards a partition of intervals with i.i.d. lengths with law ν_α such that

$$\int_0^\infty e^{-uz} \nu_\alpha(dz) = 1 - e^{\beta \int_u^\infty \frac{dx}{c_\alpha x^\alpha - \beta x}} \text{ for any } u \geq 0. \quad (6.2)$$

We now turn to the martingale problem satisfied by the inverse flow of the stable CSBP. One easily computes the drift and the jump measure from Remark 5.7. By definition of \mathcal{D} , since $\beta = \sigma = 0$, we have $\mathcal{D} = \{F \in \mathcal{C}_0^2 : F' \in L^1\}$.

Proposition 6.1. *The process $(\hat{X}_t, t \geq 0)$ is characterized by the martingale problem associated with $\hat{\mathcal{L}}$, acting on \mathcal{D} , given in Lemma 5.6 with*

$$\nu(z, dh) = \left((z-h)h^{-1-\alpha} + \frac{h^{-\alpha}}{\alpha} \right) \mathbb{1}_{[0,z]}(h) dh \text{ and } b(z) = \frac{1}{\alpha(\alpha-1)(2-\alpha)} z^{2-\alpha} - \beta z.$$

In the critical case, one can identify the law of \hat{X} through some random-time change.

Proposition 6.2. *If $\beta = 0$, the process $(\hat{X}_t, t \geq 0)$ is a positive self-similar Markov process with index $a := \alpha - 1$. Namely for any $k > 0$ and any $y > 0$,*

$$(k\hat{X}_{k^{-a}t}(y), t \geq 0) \stackrel{\mathcal{L}}{=} (\hat{X}_t(ky), t \geq 0).$$

Moreover,

$$\log \hat{X}_t(x) = L_{\varphi^x(t)}$$

where $\varphi^x(t) := \inf\{s > 0; \int_0^s e^{(\alpha-1)L_u} du > t\}$ and L is a spectrally negative Lévy process started from $\log x$ with Laplace exponent

$$\kappa(q) = -d_\alpha q + \int_{-\infty}^0 (e^{qz} - 1 + q(1 - e^z)) \nu_\alpha(dz)$$

with $\nu_\alpha(dz) = (e^z(1 - e^z)^{-1-\alpha} + \frac{1}{\alpha}(1 - e^z)^{-\alpha}) e^z dz$ and $d_\alpha = \frac{1}{\alpha(\alpha-1)(2-\alpha)}$.

Proof. Recall that the critical CSBP $(X_t, t \geq 0)$ is itself selfsimilar with index $a := \alpha - 1$. See for instance Kyprianou and Pardo [KP08]. For any $k > 0$ and any $x > 0$, $kX_{k^{-a}t}(x) \stackrel{\mathcal{L}}{=} X_t(kx)$. Thus for any $y > 0$

$$\mathbb{P}(k\hat{X}_{k^{-a}t}(y) \leq x) = \mathbb{P}(X_{k^{-a}t}(x/k) \geq y) = \mathbb{P}(k^{-1}X_t(x) > y) = \mathbb{P}(\hat{X}_t(ky) \leq x).$$

By Lamperti's representation of positive self-similar Markov process, see e.g. [Kyp14, Chapter 13], $\hat{X}_t(x)$ is of the form $\exp(L_{\varphi^x(t)})$ for some Lévy process L where $t \mapsto \varphi^x(t)$ the time-change given in the statement. To identify the Laplace exponent κ of L , note that by $(\alpha - 1)$ -self-similarity, one has $\kappa(q) = x^{-q+\alpha-1} \hat{\mathcal{L}}p_q(x)$ with $p_q(x) = x^q$. The result follows from simple computations. \square

6.2 Neveu CSBP

We now turn in this section to the Neveu CSBP, with branching mechanism $\Psi(q) = q \log(q)$. Recall its Lévy-Khintchine form

$$\Psi(q) = (\gamma - 1)q + \int_0^\infty \left(e^{-qh} - 1 + qh \mathbb{1}_{\{h \leq 1\}} \right) \frac{dh}{h^2}, \text{ for any } q \geq 0$$

where $\gamma = \int_1^\infty e^{-y} y^{-2} dy$ is the Euler-Mascheroni constant. Note that Grey's condition is not satisfied by this process. Solving the differential equation (1.3) yields $v_t(\lambda) = \lambda e^{-t}$ for any $t \geq 0$ and $\lambda \in (0, \infty)$. For any fixed t , the subordinator $(X_t(x), x \geq 0)$ is stable with parameter e^{-t} . For Neveu CSBP, the consecutive coalescent process C^λ defined in Section 4 happens to be homogeneous in time, and not to depend on λ .

Proposition 6.3. *For any $\lambda > 0$, the consecutive process $(C^\lambda(t), t \geq 0)$ is an homogeneous consecutive coalescent whose coagulation rate μ is $\mu(k) = \frac{1}{k(k-1)}$ for any $k \geq 2$. The block sizes at time $t \geq 0$ have generating function $\mathbb{E}[z^{\#C_1(t)}] = 1 - (1-z)e^{-t}$ and for any $k \geq 1$*

$$\mathbb{P}(\#C_1(t) = k) = \frac{e^{-t}(2 - e^{-t}) \dots (k - 1 - e^{-t})}{k!}.$$

Proof. By Theorem 4.10, and applying the change of variable $u = v_t(\lambda)x$, we see that for any $k \geq 2$,

$$\mu_t^\lambda(k) = v_t(\lambda)^{k-1} \int_0^\infty \frac{x^k}{k!} e^{-v_t(\lambda)x} \frac{dx}{x^2} = \frac{1}{k(k-1)}$$

which does not depend on λ nor on t . Since $C(t)$ is a (λ, v_t) -Poisson box with $v_t(q) = q e^{-t}$, the other statements can be obtained by a direct application of Theorem 4.10. See also the calculations around Lemma 7 in Pitman [Pit97]. \square

Lemma 6.4. *Consider a consecutive coalescent $(C(t), t \geq 0)$ with coagulation rate $\mu(k) = \frac{1}{k(k-1)}$ for any $k \geq 2$ then, as n goes to ∞*

$$\left(\frac{\#C_{\lfloor nx \rfloor}(t)}{n e^{-t}}, t \geq 0, x \geq 0 \right) \Longrightarrow (\hat{X}_t(x), t \geq 0, x \geq 0)$$

in finite-dimensional sense in time and in the Skorokhod topology in x .

Proof. We simply prove the convergence in law of $\left(\frac{\#C_{\lfloor nx \rfloor}(t)}{n e^{-t}}, x \geq 0 \right)$ toward \hat{X}_t for a fixed value of t , with the Skorokhod topology. Then, the finite-dimensional convergence is deduce from the cocycle property of \hat{X} (Proposition 2.3) and C (Proposition 4.2). For any $t > 0$ and $n \in \mathbb{N}$, set $Z_{-t,0}(n) = \sum_{j=1}^n \#C_j(t)$. The process $(Z_{-t,0}(n), n \geq 0)$ is a random walk and from Proposition 6.3 we see that

$$\begin{aligned} \mathbb{P}(\#C_1(t) = k) &= \frac{e^{-t}(2 - e^{-t}) \dots (k - 1 - e^{-t})}{k!} \\ &= e^{-t} \frac{\Gamma(k - e^{-t})}{\Gamma(2 - e^{-t})\Gamma(k + 1)} \sim_{k \rightarrow \infty} \frac{e^{-t}}{\Gamma(2 - e^{-t})} k^{-1-e^{-t}} \end{aligned}$$

Therefore, the law of $\#C_1$ is in the domain of attraction of a stable random variable with parameter e^{-t} . Using an extension of Donsker's theorem to stable distributions, due to Prokhorov [Pro56], we obtain that

$$\left(\frac{Z_{-t,0} \left(\lfloor n^{e^{-t}} x \rfloor \right)}{n}, x \geq 0 \right) \xrightarrow[n \rightarrow \infty]{} (\tilde{X}_t(x), x \geq 0),$$

where \tilde{X}_t is a stable subordinator with Laplace exponent $\lambda \mapsto \lambda^{e^{-t}}$.

To conclude, we observe that $(\hat{X}_t(x), x \geq 0)$ has the same law as \tilde{X}_t^{-1} the right-continuous inverse of X_t and that $(\#C_{\lfloor n \rfloor}(t), n \geq 0)$ is the right continuous inverse of $(Z_{-t,0}(n), n \geq 0)$. Hence, as the map $f \mapsto f^{-1}$ is continuous for the Skorokhod topology at any non-decreasing càdlàg function f , see e.g. [Whit80, Theorem 7.2], we have convergence in law of $\left(\frac{\#C_{\lfloor nx \rfloor}(t)}{n^{e^{-t}}}, x \geq 0 \right)$ toward $(\hat{X}_t(x), x \geq 0)$. \square

Lemma 6.5 (Möhle [Möh15], Mittag-Leffler process). *The process $(\hat{X}_t, t \geq 0)$ has for generator*

$$\hat{\mathcal{L}}f(z) = z \int_0^z (f(z-h) - f(z) + hf'(z)) \frac{dh}{h^2} + ((1-\gamma)z - z \log(z))f'(z).$$

Proof. By applying Lemma 5.6, we see that $\nu(z, dh) = \mathbb{1}_{\{h \leq z\}} ((z-h)\pi(dh) + \bar{\pi}(h)dh) = \mathbb{1}_{\{h \leq z\}} \frac{z}{h^2}$. For any $z \geq 0$,

$$\begin{aligned} b(z) &= (1-\gamma)z + \int_0^\infty \frac{z}{h} (1_{\{h \leq 1\}} - 1_{\{h \leq z\}}) dh \\ &= z \int \frac{1}{h} (1_{\{h \leq 1\}} - 1_{\{h \leq z\}}) 1_{\{z \leq 1\}} dh - z \int \frac{1}{h} (1_{\{h \leq z\}} - 1_{\{h \leq 1\}}) 1_{\{z > 1\}} dh \\ &= (1-\gamma)z + z \int_z^1 \frac{dh}{h} 1_{\{z \leq 1\}} - z \int_1^z \frac{dh}{h} 1_{\{z > 1\}} = (1-\gamma)z - z \log(z). \end{aligned} \quad \square$$

Proposition 6.6 (Bertoin and Baur [BB15]). *The process $(\log \hat{X}_t, t \geq 0)$ is a generalized Ornstein-Uhlenbeck process:*

$$\log \hat{X}_t = \log(x) + L_t - \int_0^t \log \hat{X}_s ds \quad (6.3)$$

where $(L_t, t \geq 0)$ is a spectrally negative Lévy process with Laplace exponent

$$\kappa(q) = -\gamma q + \int_{-\infty}^0 (e^{qu} - 1 - qu) \frac{e^u}{(1-e^u)^2} du.$$

Proof. By injectivity of $g : x \mapsto \log(x)$, the generator of $(Y_t, t \geq 0) := (\log \hat{X}_t, t \geq 0)$ is given by $\mathcal{A}f(y) = \hat{\mathcal{L}}(f \circ g)(g^{-1}(y))$ and a computation provides

$$\mathcal{A}f(y) = \int_{-\infty}^0 (f(y+u) - f(y) - uf'(y)) \nu(du) - \gamma f'(y),$$

with $\nu(du) = \frac{e^u}{(1-e^u)^2} du$. It is well-known that the process with generator \mathcal{A} is a weak solution to the equation (6.3). See for instance, Sato and Yamazato [SY84, Theorem 3.1]. \square

The two last statements already appear in the study of the Bolthausen-Sznitman coalescent. We explain now some connections between the Neveu consecutive coalescent and the Bolthausen-Sznitman exchangeable coalescent. The following is a rephrasing of an observation made by Hénard [Hén15] and Möhle [Möh15]. Denote by $(N_t^{(n)}, t \geq 0)$ the number of blocks in a Bolthausen-Sznitman coalescent started from n blocks. Recall that $(N_t^{(n)}, t \geq 0)$ jumps from n to $n - k + 1$ at rate $\frac{n}{k(k+1)}$ for any $k \in \llbracket 2, n \rrbracket$. By Proposition 4.5, one can check that $(\#C_{\lfloor n \rfloor}(t), t \geq 0)$ loses k blocks at the same rate as $(N_t^{(n)}, t \geq 0)$. Therefore $(N_t^{(n)}, t \geq 0)$ and $(\#C_{\lfloor n \rfloor}(t), t \geq 0)$ have the same law and by Lemma 6.4, as n goes to ∞

$$\left(\frac{N_t^{(n)}}{n e^{-t}}, t \geq 0 \right) \Longrightarrow (\hat{X}_t(1), t \geq 0)$$

in the finite-dimensional sense. Such a convergence was shown, in the Skorohod topology, by Möhle in [Möh15, Theorem 1.1], Kukla and Möhle in [KM18, Theorem 2.1-(a)] with different techniques. We refer also to Bertoin and Baur [BB15, Theorem 3.1-(i)] for an almost-sure convergence. The connections between Neveu's consecutive coalescent and Bolthausen-Sznitman exchangeable one are not surprising since it is known that for any initial size x , the genealogy of i.i.d random variables sampled in $[0, x]$ is described by a Bolthausen-Sznitman coalescent, see Bertoin and Le Gall [BLG00, Theorem 4].

Several natural questions on the inverse flow and its consecutive coalescent have not been addressed here and are left for possible future works. It might be interesting for instance to look for a complete description of the two-parameter flow $(\hat{X}_t(x), t \geq 0, x \geq 0)$ in the general case, as given for the Feller flow in Section 3. Moreover, the genealogy of the total population has only been characterized under the Grey's condition. When this condition is not fulfilled the process $(\mathcal{C}(t), t \geq 0)$ cannot be described by a single consecutive coalescent on \mathbb{N} . We recall that Duquesne and Winkel [DW07] have described the genealogy forward in time of a CSBP (including those without Grey's condition) through a collection of continuous-time Galton-Watson processes. A natural question is thus to see if in a dual way, one can represent the backward genealogy of the total population with a collection of consecutive coalescents on \mathbb{N} .

Acknowledgement. C.F and B.M are partially supported by the French National Research Agency (ANR): ANR GRAAL and ANR MALIN; LABEX MME-DII and CNRS PEPS grants. C.M is supported by the NSFC of China (11871032). C.F would like to thank T. Duquesne and V. Rivero for fruitful discussions at early stages of this work.

A Intermediary results

A.1 Right-continuous inverse

In this section, we recall and compile some elementary properties on right continuous inverse of càdlàg non-decreasing functions. As multiple competing definitions of generalized inverse coexist, it can be challenging to find a single reference for the results we need. Therefore we give a short proof of these well-known facts, in order to be self-contained. Let f be a càdlàg non-decreasing function on \mathbb{R}_+ , we denote by

$$f^{-1} : y \in [0, \infty) \mapsto \inf\{x \geq 0 : f(x) > y\} \tag{A.1}$$

its right continuous inverse.

Lemma A.1. *Let $f, (f_n, n \geq 1), g$ be càdlàg non-decreasing functions on \mathbb{R}_+ .*

- (i) *The function f^{-1} is non-decreasing and càdlàg.*
- (ii) *For every $x, y \geq 0$, we have $f(x) > y \iff f^{-1}(y) < x$.*
- (iii) *We have $(f \circ g)^{-1} = g^{-1} \circ f^{-1}$.*
- (iv) *If $\lim_{n \rightarrow \infty} f_n = \text{Id}$ pointwise, then $\lim_{n \rightarrow \infty} f_n^{-1} = \text{Id}$ pointwise, with Id being the identity function on $[0, \infty)$.*

Remark A.2. Dini's theorems imply that both convergences in (iv) hold uniformly on compact sets.

Proof. Let f be a càdlàg non-decreasing function, note that for all $y < z$, we have

$$\{x \geq 0 : f(x) > z\} \subset \{x \geq 0 : f(x) > y\}.$$

Therefore $f^{-1}(y) \leq f^{-1}(z)$, which proves that f^{-1} is increasing. In particular, it has left limits at each point. We now observe that for all $y \geq 0$, as f is non-decreasing,

$$\inf_{z > y} f^{-1}(z) = \inf\{\inf\{x \geq 0 : f(x) > z\}, z > y\} = \inf\{x \geq 0 : f(x) > y\} = f^{-1}(y),$$

proving that f^{-1} is right continuous at point y , which proves (i).

Let $x, y \geq 0$, we first assume that $f^{-1}(y) < x$. Then by definition of f^{-1} , there exists $u \in [f^{-1}(y), x)$ such that $f(u) > y$. As f is non-decreasing, we deduce that $f(x) \geq f(u) > y$.

We now assume that $f^{-1}(y) \leq x$. As f is non-decreasing, we observe that $f(x) \geq f(f^{-1}(y))$. Therefore, the only thing left to prove is that

$$\forall y \geq 0, f(f^{-1}(y)) \geq y \tag{A.2}$$

We write $z = f^{-1}(y)$. By definition of $f^{-1}(y)$, for all $\epsilon > 0$, there exists $u < z + \epsilon$ such that $f(u) > y$. Then, as f is right-continuous, we have $f(z) = \inf_{u > z} f(u)$, thus for all $\eta > 0$, there exists $\epsilon > 0$ such that if $u < z + \epsilon$, then $f(u) < f(z) + \eta$. As a result, for all $\eta > 0$, there exists $u < z + \epsilon$ such that $y < f(u) < f(z) + \eta$. This inequality being true for all $\eta > 0$, we therefore conclude that $f(z) \geq y$, completing the proof (A.2). We thus deduce that $f(x) \geq y$, completing the proof of (ii).

In a third time, we note that given f and g two càdlàg non-decreasing functions on \mathbb{R}_+ , we have for all $y \geq 0$,

$$(f \circ g)^{-1}(y) = \inf\{z \geq 0 : (f \circ g)(z) > y\} = \inf\{z \geq 0 : f(g(z)) > y\}.$$

By point (ii), this can therefore be rewritten as

$$(f \circ g)^{-1}(y) = \inf\{z \geq 0 : g(z) > f^{-1}(y)\} = g^{-1} \circ f^{-1}(y),$$

proving point (iii).

We finally prove the last point. Let (f_n) be a sequence of non-decreasing càdlàg functions such that $\lim_{n \rightarrow \infty} f_n = \text{Id}$ pointwise. We prove that for all $y \geq 0$, $\lim_{n \rightarrow \infty} f_n^{-1}(y) = y$. Let $\epsilon > 0$, by point (ii), we have that

$$f_n^{-1}(y) < y + \epsilon \iff f_n(y + \epsilon) > y.$$

As $\lim_{n \rightarrow \infty} f_n(y + \epsilon) = y + \epsilon$, we conclude that for all n large enough, $f_n^{-1}(y) < y + \epsilon$. Similarly, we also have

$$f_n^{-1}(y) \geq y - \epsilon \iff f_n(y - \epsilon) \leq y$$

therefore $f_n^{-1}(y) \geq y - \epsilon$ for all n large enough by pointwise convergence of f_n at point $y - \epsilon$. This concludes the proof of (iv). \square

A.2 Discretization of subordinators

In this section, we introduce the key lemma allowing us to study the genealogical structure of CSBPs. Namely, we prove that the pullback measure of a Poisson process by a subordinator is a Poisson process decorated by i.i.d. integer-valued random variables.

Lemma A.3. *Let $\lambda \geq 0$ and $(X(x), x \geq 0)$ be a subordinator with Laplace exponent*

$$\phi : \mu \mapsto d\mu + \int (1 - e^{-\mu x}) \ell(dx).$$

We denote by N an independent Poisson point process with intensity λ , and we write $(J_j, j \geq 1)$ the positions of the atoms of N , ranked in the decreasing order. Then, setting $M = \sum_{j=1}^{\infty} \delta_{X^{-1}(J_j)}$ the image measure of N by X^{-1} , we have

$$M = \sum_{j=1}^{\infty} Z_j \delta_{J'_j},$$

where $(J'_j, j \geq 1)$ are the atoms of a Poisson point process with intensity $\phi(\lambda)$ and $(Z_j, j \geq 1)$ are i.i.d. random variables, independent of $(J'_j, j \geq 1)$, such that

$$\mathbb{P}(Z_1 = k) = \frac{1}{\phi(\lambda)} \int_0^{\infty} \frac{(\lambda x)^k}{k!} e^{-\lambda x} \ell(dx) + d\mathbf{1}_{\{k=1\}} = (-1)^{k-1} \frac{\lambda^k}{k!} \frac{\phi^{(k)}(\lambda)}{\phi(\lambda)},$$

i.e. $\mathbb{E}(z^{Z_1}) = 1 - \frac{\phi(\lambda(1-z))}{\phi(\lambda)}$ for all $z \in [0, 1]$.

Proof. The proof is based on a joint construction by the same “master” Poisson point process of the subordinator X and the Poisson point process N , in such a way that M becomes a simple functional of that master point process. To see why such a construction is possible, we write

$$\phi(\lambda) = d\lambda + \int (1 - e^{-\lambda x}) \ell(dx),$$

with $d \geq 0$ the drift and ℓ the Lévy measure of X on \mathbb{R}_+ . By the Lévy-Itô décomposition, one can write

$$\forall t \geq 0, \quad X_t = dt + \sum_{0 \leq s \leq t} x_s,$$

with $(t, x_t)_{t \geq 0}$ being the atoms of a Poisson point process with intensity $dt \otimes \ell(dx)$. The proof being slightly simpler for $d = 0$, we focus here on the case $d > 0$.

Recalling that \mathcal{D} denotes the set of càdlàg non-decreasing functions on \mathbb{R}_+ , we introduce the point process $R = \sum_{i \in I} \delta_{(t_i, x_i, N^{(i)})}$ on $\mathbb{R}_+ \times \mathbb{R}_+ \times \mathcal{D}$ with intensity $dt \otimes dx \otimes \mathcal{P}^\lambda(dN)$, with \mathcal{P}^λ being the law of a Poisson point process with intensity λ on \mathbb{R}_+ . We also set $N^{(0)}$ an independent Poisson point process with intensity λ . We then define

$$\bar{X}_t = dt + \sum_{i \in I} x_i \mathbf{1}_{\{t_i \leq t\}},$$

which is a subordinator with Laplace exponent ϕ . Then, denoting $(J_j^{(i)}, j \geq 1)$ the atoms of the Poisson point process $N^{(i)}$, we set

$$\bar{N} = \sum_{j=1}^{\infty} \delta_{\bar{X}_{J_j^{(0)}/d}} + \sum_{i \in I} \sum_{j=1}^{\infty} \delta_{\bar{X}_{t_i-} + J_j^{(i)}} \mathbf{1}_{\{J_j^{(i)} < x_i\}}.$$

Heuristically, the point process \bar{N} can be thought of as follows: \mathbb{R}_+ is divided in intervals $\cup_{i \in I} [\bar{X}_{t_i-}, \bar{X}_{t_i}]$ corresponding to jumps in the subordinator \bar{X} and the remaining space B corresponding to points with an antecedent by \bar{X} . Atoms are added to the interval $[\bar{X}_{t_i-}, \bar{X}_{t_i}]$ according to the point process $N^{(i)}$, and to the set B with the point process $N^{(0)}$. It should then be heuristically clear that \bar{N} is a Poisson point process with intensity λ independent of \bar{X} . To verify it, we compute its conditional Laplace transform against a smooth locally compact test function f . By construction, $(N^{(i)}, i \in I \cup \{0\})$ are i.i.d. Poisson point process with intensity λ , which are further independent from \bar{X} , thus

$$\begin{aligned} & \mathbb{E} \left(\exp(-\langle \bar{N}, f \rangle) \middle| \bar{X} \right) \\ &= \mathbb{E} \left(\exp \left(- \sum_{j \geq 0} f(\bar{X}_{J_j^{(0)}/d}) \right) \middle| \bar{X} \right) \prod_{i \in I} \mathbb{E} \left(\exp \left(- \sum_{j \geq 0} f(J_j^{(i)} + \bar{X}_{t_i-}) \mathbf{1}_{\{J_j^{(i)} < x_i\}} \right) \middle| \bar{X} \right) \\ &= \exp \left(-\lambda \int_0^\infty (1 - e^{-f(\bar{X}_s/d)}) ds - \lambda \sum_{i \in I} \int_{\bar{X}_{t_i-}}^{\bar{X}_{t_i}} (1 - e^{-f(\bar{X}_s)}) ds \right) \quad \text{a.s.} \\ &= \exp \left(\lambda \int (1 - e^{-f(x)}) dx \right), \end{aligned}$$

by change of variable, using that $\bar{X}'_t = d$ at all continuity points t of \bar{X} .

As a result, the couple (\bar{X}, \bar{N}) has same law as (X, N) given in the lemma. Moreover, we have immediately by construction that

$$\bar{M} := \bar{X}^{-1} * \bar{N} = \sum_{j=1}^{\infty} \delta_{J_j^{(0)}/d} + \sum_{i \in I} N^{(i)}([0, x_i]) \delta_{t_i}.$$

and computing the law of that point process is straightforward by the definition of R . Indeed, by independence, for any continuous function f with compact support, we have

$$\mathbb{E} \left(\exp(-\langle \bar{M}, f \rangle) \right) = \mathbb{E} \left(\exp \left(- \sum_{j \geq 1} f(J_j^{(0)}/d) \right) \right) \mathbb{E} \left(\exp \left(- \sum_{i \in I} N^{(i)}([0, x_i]) f(t_i) \right) \right).$$

Then, using Campbell's formula, we have both

$$\begin{aligned} \mathbb{E} \left(\exp \left(- \sum_{j \geq 1} f(J_j^{(0)}/d) \right) \right) &= \exp \left(\lambda d \int (1 - e^{-f(x)}) dx \right) \\ \mathbb{E} \left(\exp \left(- \sum_{i \in I} N^{(i)}([0, x_i]) f(t_i) \right) \right) &= \exp \left(\int (1 - e^{-N([0, x]) f(t)}) dt \ell(dx) \mathcal{P}^\lambda(dN) \right). \end{aligned}$$

But as under law \mathcal{P}^λ , $N([0, x])$ is a Poisson random variable with parameter λx , the last inequality can be written, by Fubini theorem

$$\begin{aligned} \mathbb{E} \left(\exp \left(- \sum_{i \in I} N^{(i)}([0, x_i]) f(t_i) \right) \right) &= \exp \left(\int 1 - \exp(\lambda x (e^{-f(t)} - 1)) dt \ell(dx) \right) \\ &= \exp \left(- \int \frac{\phi(\lambda(1 - e^{-f(t)}))}{\phi(\lambda)} dt \right). \end{aligned}$$

We deduce that the Laplace transform of \overline{M} is the same as the one of M given in the lemma, which concludes the proof. \square

This result can be straightforwardly extended to killed subordinators, by constructing it as a limit of non-killed subordinators. For the sake of completeness, we add a proof of the following result.

Corollary A.4. *Let $\lambda \geq 0$ and $(X(x), x \geq 0)$ be a subordinator with Laplace exponent*

$$\phi : \mu \mapsto \kappa + d\mu + \int (1 - e^{-\mu x}) \ell(dx).$$

With the same notation as in the previous lemma, we have $M = \sum_{j=1}^{\infty} Z_j' \delta_{J_j'}$, where $(J_j', j \geq 1)$ are the atoms of a Poisson point process with intensity $\phi(\lambda)$, $(Z_j', j \geq 1)$ are i.i.d. random variables, independent of $(J_j', j \geq 1)$, such that

$$\mathbb{P}(Z_1 = k) = \frac{1}{\phi(\lambda)} \int_0^\infty \frac{(\lambda x)^k}{k!} e^{-\lambda x} \ell(dx) + d \mathbf{1}_{\{k=1\}} = (-1)^{k-1} \frac{\lambda^k \phi^{(k)}(\lambda)}{k! \phi(\lambda)},$$

$$\text{and } Z_j' = \begin{cases} Z_j & \text{if } \sup_{i < j} Z_i < \infty, \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Let Y be a subordinator with Laplace exponent $\lambda \mapsto d\lambda + \int 1 - e^{-\lambda x} \ell(dx)$, and R an independent Poisson process with intensity κ . Observe that for all $r > 0$, the process defined by

$$Y_t^r = Y_t + rR_t, \quad t \geq 0,$$

is a Lévy process, and that $X = \lim_{r \rightarrow \infty} Y^r$ is a Lévy process with Laplace exponent ϕ . We set $(J_j^r, j \geq 1)$ and $(Z_j^r, j \geq 1)$ the quantities obtained by applying Lemma A.3, and $T = \inf\{t > 0 : R_t = 1\}$. Observe that for all j such that $J_j^r < T$, the quantities J_j^r and Z_j^r do not depend on r . On the contrary, past time T , as $r \rightarrow \infty$, all values J_j^r converge toward T , and the associated value of Z^r converges toward ∞ .

Explicit formulas for the law of Z^∞ are straightforward Poisson computations. \square

References

- [Ald93] David Aldous, *The continuum random tree. III*, Ann. Probab. **21** (1993), no. 1, 248–289. MR 1207226
- [AN04] Krishna Athreya and Peter Ney, *Branching processes*, Dover Publications, Inc., Mineola, NY, 2004, Reprint of the 1972 original [Springer, New York; MR0373040]. MR 2047480
- [AP05] David Aldous and Lea Popovic, *A critical branching process model for biodiversity*, Adv. in Appl. Probab. **37** (2005), no. 4, 1094–1115. MR 2193998
- [Ath12] Krishna Athreya, *Coalescence in critical and subcritical Galton-Watson branching processes*, J. Appl. Probab. **49** (2012), no. 3, 627–638. MR 3012088
- [BB15] Erich Baur and Jean Bertoin, *The fragmentation process of an infinite recursive tree and Ornstein-Uhlenbeck type processes*, Electron. J. Probab. **20** (2015), no. 98, 20. MR 3399834
- [BBC⁺05] Matthias Birkner, Jochen Blath, Marcella Capaldo, Alison Etheridge, Martin Möhle, Jason Schweinsberg, and Anton Wakolbinger, *Alpha-stable branching and beta-coalescents*, Electron. J. Probab. **10** (2005), no. 9, 303–325. MR 2120246
- [BBS07] Julien Berestycki, Nathanaël Berestycki, and Jason Schweinsberg, *Beta-coalescents and continuous stable random trees*, Ann. Probab. **35** (2007), no. 5, 1835–1887. MR 2349577
- [BD16] Hongwei Bi and Jean-François Delmas, *Total length of the genealogical tree for quadratic stationary continuous-state branching processes*, Ann. Inst. Henri Poincaré Probab. Stat. **52** (2016), no. 3, 1321–1350. MR 3531711
- [Ber06] Jean Bertoin, *Random fragmentation and coagulation processes*, Cambridge Studies in Advanced Mathematics, vol. 102, Cambridge University Press, Cambridge, 2006. MR 2253162
- [BFM08] Jean Bertoin, Joaquín Fontbona, and Servet Martínez, *On prolific individuals in a supercritical continuous-state branching process*, J. Appl. Probab. **45** (2008), no. 3, 714–726. MR 2455180
- [BLG00] Jean Bertoin and Jean-François Le Gall, *The Bolthausen-Sznitman coalescent and the genealogy of continuous-state branching processes*, Probab. Theory Related Fields **117** (2000), no. 2, 249–266. MR 1771663
- [BLG03] ———, *Stochastic flows associated with coalescent processes*, Probab. Theory Related Fields **126** (2003), no. 2, 261–288. MR 1990057
- [BLG05] ———, *Stochastic flows associated to coalescent processes. II. Stochastic differential equations*, Ann. Inst. H. Poincaré Probab. Statist. **41** (2005), no. 3, 307–333. MR 2139022 (2005m:60067)
- [BLG06] ———, *Stochastic flows associated with coalescent processes. III. Limit theorems*, Illinois J. Math. **50** (2006), no. 1-4, 147–181 (electronic). MR 2247827 (2008c:60032)

- [CD12] Yu-Ting Chen and Jean-François Delmas, *Smaller population size at the MRCA time for stationary branching processes*, Ann. Probab. **40** (2012), no. 5, 2034–2068. MR 3025710
- [CS85] Peter Clifford and Aidan Sudbury, *A sample path proof of the duality for stochastically monotone Markov processes*, Ann. Probab. **13** (1985), no. 2, 558–565. MR 781422
- [DFM14] Xan Duhalde, Clément Foucart, and Chunhua Ma, *On the hitting times of continuous-state branching processes with immigration*, Stochastic Process. Appl. **124** (2014), no. 12, 4182–4201. MR 3264444
- [DK99] Peter Donnelly and Thomas G. Kurtz, *Particle representations for measure-valued population models*, Ann. Probab. **27** (1999), no. 1, 166–205. MR 1681126 (2000f:60108)
- [DL14] Thomas Duquesne and Cyril Labbé, *On the Eve property for CSBP*, Electron. J. Probab. **19** (2014), no. 6, 31. MR 3164759
- [DLG02] Thomas Duquesne and Jean-François Le Gall, *Random trees, Lévy processes and spatial branching processes*, Astérisque (2002), no. 281, vi+147. MR 1954248
- [DW07] Thomas Duquesne and Matthias Winkel, *Growth of Lévy trees*, Probab. Theory Related Fields **139** (2007), no. 3-4, 313–371. MR 2322700
- [ER10] Steven N. Evans and Peter L. Ralph, *Dynamics of the time to the most recent common ancestor in a large branching population*, Ann. Appl. Probab. **20** (2010), no. 1, 1–25. MR 2582640
- [Fel51] William Feller, *Diffusion processes in genetics*, Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability, 1950, University of California Press, Berkeley and Los Angeles, 1951, pp. 227–246. MR 0046022
- [FFK17] Dorrotya Fekete, Joaquín Fontbona, and Andreas Kyprianou, *Skeletal stochastic differential equations for continuous-state branching process*, to appear in J. App. Probab. **56** (2019)
- [FH13] Clément Foucart and Olivier Hénard, *Stable continuous-state branching processes with immigration and Beta-Fleming-Viot processes with immigration*, Electron. J. Probab. **18** (2013), no. 23, 21. MR 3035751
- [FM16] Clément Foucart and Chunhua Ma, *Continuous-state branching processes, extremal processes and super-individuals*, Ann. Inst. H. Poincaré Probab. Statist. **55** (2019), no. 2, 1061-1086. MR 3949964.
- [Fou12] Clément Foucart, *Generalized Fleming-Viot processes with immigration via stochastic flows of partitions*, ALEA Lat. Am. J. Probab. Math. Stat. **9** (2012), no. 2, 451–472. MR 3069373
- [Get80] Ronald Gettoor, *Transience and recurrence of Markov processes*, Seminar on Probability, XIV (Paris, 1978/1979) (French), Lecture Notes in Math., vol. 784, Springer, Berlin, 1980, pp. 397–409. MR 580144

- [GH16] Nicolas Grosjean and Thierry Huillet, *On a coalescence process and its branching genealogy*, J. Appl. Probab. **53** (2016), no. 4, 1156–1165. MR 3581248
- [Gre74] David Grey, *Asymptotic behaviour of continuous time, continuous state-space branching processes*, J. Appl. Probability **11** (1974), 669–677. MR 0408016
- [Gri74] Anders Grimvall, *On the convergence of sequences of branching processes*, Ann. Probability **2** (1974), 1027–1045. MR 0362529
- [HJR17] Simon. C. Harris, Samuel. G. G. Johnston, and Matt Roberts, *The coalescent structure of continuous-time Galton-Watson trees*, arXiv e-prints, March 2017.
- [Hén15] Olivier Hénard, *The fixation line in the Λ -coalescent*, Ann. Appl. Probab. **25** (2015), no. 5, 3007–3032. MR 3375893
- [ILP15] Gautam Iyer, Nicholas Leger, and Robert L. Pego, *Limit theorems for Smoluchowski dynamics associated with critical continuous-state branching processes*, Ann. Appl. Probab. **25** (2015), no. 2, 675–713. MR 3313753
- [ILP18] ———, *Coagulation and universal scaling limits for critical galton-watson processes*, Advances in Applied Probability **50** (2018), no. 2, 504–542.
- [Jiř58] Miloslav Jiřina, *Stochastic branching processes with continuous state space*, Czechoslovak Math. J. **8** (**83**) (1958), 292–313. MR 0101554
- [Jiř69] ———, *On Feller’s branching diffusion processes*, Časopis Pěst. Mat. **94** (1969), 84–90, 107. MR 0247676
- [JL19+] Samuel Johnston and Amaury Lambert, *The coalescent structure of samples from branching processes: a unifying poissonisation approach*, In preparation, 2019+.
- [Joh17] Samuel Johnston, *Coalescence in supercritical and subcritical continuous-time Galton-Watson trees*, arXiv e-prints, September 2017.
- [Kal02] Olav Kallenberg, *Foundations of modern probability*, second ed., Probability and its Applications (New York), Springer-Verlag, New York, 2002. MR 1876169 (2002m:60002)
- [KM18] Jonas Kukla and Martin Möhle, *On the block counting process and the fixation line of the Bolthausen-Sznitman coalescent*, Stochastic Process. Appl. **128** (2018), no. 3, 939–962. MR 3758343
- [Kol11] Vassili Kolokol’tsov, *Stochastic monotonicity and duality of one-dimensional Markov processes*, Mat. Zametki **89** (2011), no. 5, 694–704. MR 2858558
- [KP08] Andreas Kyprianou and Juan-Carlos Pardo, *Continuous-state branching processes and self-similarity*, J. Appl. Probab. **45** (2008), no. 4, 1140–1160. MR 2484167
- [Kyp14] Andreas E. Kyprianou, *Fluctuations of Lévy processes with applications*, second ed., Universitext, Springer, Heidelberg, 2014, Introductory lectures. MR 3155252
- [Lab14a] Cyril Labbé, *From flows of Λ -Fleming-Viot processes to lookdown processes via flows of partitions*, Electron. J. Probab. **19** (2014), no. 55, 49. MR 3227064

- [Lab14b] ———, *Genealogy of flows of continuous-state branching processes via flows of partitions and the Eve property*, Ann. Inst. Henri Poincaré Probab. Stat. **50** (2014), no. 3, 732–769. MR 3224288
- [Lam67a] John Lamperti, *Continuous state branching processes*, Bull. Amer. Math. Soc. **73** (1967), 382–386. MR 0208685
- [Lam67b] ———, *The limit of a sequence of branching processes*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **7** (1967), 271–288. MR 0217893
- [Lam03] Amaury Lambert, *Coalescence times for the branching process*, Adv. in Appl. Probab. **35** (2003), no. 4, 1071–1089. MR 2014270
- [Lam07] ———, *Quasi-stationary distributions and the continuous-state branching process conditioned to be never extinct*, Electron. J. Probab. **12** (2007), no. 14, 420–446. MR 2299923
- [Le14] Vi Le, *Coalescence times for the Bienaymé-Galton-Watson process*, J. Appl. Probab. **51** (2014), no. 1, 209–218. MR 3189452
- [LGLJ98] Jean-Francois Le Gall and Yves Le Jan, *Branching processes in Lévy processes: the exploration process*, Ann. Probab. **26** (1998), no. 1, 213–252. MR 1617047
- [Li00] Zeng-Hu Li, *Asymptotic behaviour of continuous time and state branching processes*, J. Austral. Math. Soc. Ser. A **68** (2000), no. 1, 68–84. MR 1727226
- [Li11] Zenghu Li, *Measure-valued branching Markov processes*, Probability and its Applications (New York), Springer, Heidelberg, 2011. MR 2760602
- [LP13] Amaury Lambert and Lea Popovic, *The coalescent point process of branching trees*, Ann. Appl. Probab. **23** (2013), no. 1, 99–144. MR 3059232
- [LPLG08] Yangrong Li, Anthony G. Pakes, Jia Li, and Anhui Gu, *The limit behavior of dual Markov branching processes*, J. Appl. Probab. **45** (2008), no. 1, 176–189. MR 2409319
- [LUB17] Amaury Lambert and Gerónimo Uribe Bravo, *The comb representation of compact ultrametric spaces*, p-Adic Numbers, Ultrametric Analysis and Applications **9** (2017), no. 1, 22–38.
- [Möh15] Martin Möhle, *The Mittag-Leffler process and a scaling limit for the block counting process of the Bolthausen-Sznitman coalescent*, ALEA Lat. Am. J. Probab. Math. Stat. **12** (2015), no. 1, 35–53. MR 3333734
- [Mor58] Patrick Moran, *Random processes in genetics*, Proc. Cambridge Philos. Soc. **54** (1958), 60–71. MR 0127989
- [MS01] Martin Möhle and Serik Sagitov, *A classification of coalescent processes for haploid exchangeable population models*, Ann. Probab. **29** (2001), no. 4, 1547–1562. MR 1880231
- [Par08] Étienne Pardoux, *Continuous branching processes: the discrete hidden in the continuous*, ARIMA Rev. Afr. Rech. Inform. Math. Appl. **9** (2008), 211–229. MR 2507503

- [Pit97] Jim Pitman, *Partition structures derived from Brownian motion and stable subordinators*, Bernoulli **3** (1997), no. 1, 79–96. MR 1466546
- [Pit99] ———, *Coalescents with multiple collisions*, Ann. Probab. **27** (1999), no. 4, 1870–1902. MR 1742892 (2001h:60016)
- [Pop04] Lea Popovic, *Asymptotic genealogy of a critical branching process*, Ann. Appl. Probab. **14** (2004), no. 4, 2120–2148. MR 2100386
- [Pro56] Yu. V. Prokhorov, *Convergence of random processes and limit theorems in probability theory*, Teor. Veroyatnost. i Primenen. **1** (1956), 177–238. MR 0084896
- [PY82] Jim Pitman and Marc Yor, *A decomposition of Bessel bridges*, Z. Wahrsch. Verw. Gebiete **59** (1982), no. 4, 425–457. MR 656509
- [Sag99] Serik Sagitov, *The general coalescent with asynchronous mergers of ancestral lines*, J. Appl. Probab. **36** (1999), no. 4, 1116–1125. MR 1742154
- [Sch00] Jason Schweinsberg, *Coalescents with simultaneous multiple collisions*, Electron. J. Probab. **5** (2000), Paper no. 12, 50. MR 1781024
- [Sch03] ———, *Coalescent processes obtained from supercritical Galton-Watson processes*, Stochastic Process. Appl. **106** (2003), no. 1, 107–139. MR 1983046
- [Sie76] David Siegmund, *The equivalence of absorbing and reflecting barrier problems for stochastically monotone Markov processes*, Ann. Probability **4** (1976), no. 6, 914–924. MR 0431386
- [Sil68] Martin Silverstein, *A new approach to local times*, J. Math. Mech. **17** (1967/1968), 1023–1054. MR 0226734
- [SY84] Ken-iti Sato and Makoto Yamazato, *Operator-self-decomposable distributions as limit distributions of processes of Ornstein-Uhlenbeck type*, Stochastic Process. Appl. **17** (1984), no. 1, 73–100. MR 738769
- [Whit80] Ward Whitt, *Some useful functions for functional limit theorems*, Math. Oper. Res. **5** (1980), no. 1, 67–85. MR 561155