

SOLVING THE BOLTZMANN EQUATION IN $N \log_2 N^*$

FRANCIS FILBET[†], CLÉMENT MOUHOT[‡], AND LORENZO PARESCHI[§]

Abstract. In [C. Mouhot and L. Pareschi, “Fast algorithms for computing the Boltzmann collision operator,” *Math. Comp.*, to appear; C. Mouhot and L. Pareschi, *C. R. Math. Acad. Sci. Paris*, 339 (2004), pp. 71–76], fast deterministic algorithms based on spectral methods were derived for the Boltzmann collision operator for a class of interactions including the *hard spheres model* in three dimensions. These algorithms are implemented for the solution of the Boltzmann equation in two and three dimensions, first for homogeneous solutions, then for general nonhomogeneous solutions. The results are compared to explicit solutions, when available, and to Monte Carlo methods. In particular, the computational cost and accuracy are compared to those of Monte Carlo methods as well as to those of previous spectral methods. Finally, for inhomogeneous solutions, we take advantage of the great computational efficiency of the method to show an oscillation phenomenon of the entropy functional in the trend to equilibrium, which was suggested in the work [L. Desvillettes and C. Villani, *Invent. Math.*, 159 (2005), pp. 245–316].

Key words. Boltzmann equation, spectral methods, fast algorithms, entropy

AMS subject classifications. 65T50, 68Q25, 74S25, 76P05

DOI. 10.1137/050625175

1. Introduction. The construction of approximate methods of a solution for the Boltzmann equation has a long history and can be traced back to Hilbert, Chapmann, and Enskog [13] at the beginning of the last century. The mathematical difficulties related to the Boltzmann equation make it extremely difficult, if not impossible, to determine analytic solutions in most physically relevant situations. Only in recent years, starting in the 1970s with the pioneering works by Chorin [14] and Sod [46], has the problem been tackled numerically with particular care given to accuracy and computational cost. Even nowadays the deterministic numerical solution of the Boltzmann equation still represents a challenge for scientific computing.

Most of the difficulties are due to the multidimensional structure of the collisional integral, since the integration runs on a highly dimensional unflat manifold. In addition the numerical integration requires great care since the collision integral is at the basis of the macroscopic properties of the equation. Further difficulties are represented by the presence of stiffness, such as the case of small mean free path [24] or the case of large velocities [19].

For such reasons realistic numerical simulations are based on Monte Carlo techniques. The most famous examples are the direct simulation Monte Carlo (DSMC) methods by Bird [3] and Nanbu [36]. These methods guarantee efficiency and preservation of the main physical properties. However, avoiding statistical fluctuations in the results becomes extremely expensive in the presence of nonstationary flows or when close to continuum regimes.

*Received by the editors February 24, 2005; accepted for publication (in revised form) January 5, 2006; published electronically June 30, 2006. This work was supported by the European network HYKE, funded by the EC as contract HPRN-CT-2002-00282.

<http://www.siam.org/journals/sisc/28-3/62517.html>

[†]MIP, Université Paul Sabatier 118, route de Narbonne, 31062 Toulouse Cedex 04, France (filbet@mip.ups-tlse.fr).

[‡]CEREMADE, Université Paris IX, Place du Mal de Lattre de Tassigny, 75775 Paris Cedex 16, France (cmouhot@ceremade.dauphine.fr).

[§]Dipartimento di Matematica, Università di Ferrara, Via Machiavelli 35, 44100 Ferrara, Italy (lorenzo.pareschi@unife.it).

Among deterministic approximations, perhaps the most popular method is represented by the so-called discrete velocity models (DVM) of the Boltzmann equation. These methods [29, 45, 6, 10, 39] are based on a Cartesian grid in velocity and on a discrete collision mechanism on the points of the grid that preserves the main physical properties. Unfortunately DVM are not competitive with Monte Carlo methods in terms of computational cost, and their accuracy seems to be less than first order [37, 38, 39]. In this work we are interested in high-order deterministic methods and therefore we shall not discuss algorithms based on DVM. We refer the reader to the work in preparation [34].

More recently a new class of numerical methods based on the use of spectral techniques in the velocity space has been developed. The methods were first derived in [40], inspired from spectral methods in fluid mechanics [11] and by previous works on the use of Fourier transform techniques for the Boltzmann equation (see [5], for instance). The numerical methods are based on approximating the distribution function by a periodic function in the phase space and on its representation by Fourier series. The resulting Fourier–Galerkin approximation can be evaluated with a computational cost of $O(n^2)$ (where n is the total number of discretization parameters in velocity), which is lower than that of previous deterministic methods (but still larger than that of Monte Carlo methods).

The new methods were further developed in [41, 43], where evolution equations for the Fourier modes were explicitly derived and spectral accuracy of the methods was proven. Strictly speaking these methods are not conservative, since they preserve mass, whereas momentum and energy are approximated with spectral accuracy. This trade-off between accuracy and conservation seems to be an unavoidable compromise in the development of numerical schemes for the Boltzmann equation (with the noticeable exception of [28]).

We recall here that the spectral method has been applied also to nonhomogeneous situations [20, 22]; to the Landau equation [19, 42], where fast algorithms can be readily derived; and to the case of granular gases [35, 21]. For a recent introduction to numerical methods for the Boltzmann equation and related kinetic equations we refer the reader to [15]. Finally, let us mention that Bobylev and Rjasanow [7, 8] have also constructed fast algorithms based on a Fourier transform approximation of the distribution function.

In [33, 32] a fast spectral method was proposed for a class of particle interactions, including pseudo-Maxwell molecules (in dimension 2) and hard spheres (in dimension 3), on the basis of the previous spectral method together with a suitable semidiscretization of the collision operator. This method permits us to reduce the computational cost from $O(n^2)$ to $O(n \log_2 n)$ without losing the spectral accuracy, and thus making the method competitive with Monte Carlo. The principles and basic features of this new method will be presented in the following sections.

The rest of the paper is organized as follows. Section 2 is devoted to a short introduction on the Boltzmann equation and its physical properties. In section 3 we explain the principles of the different spectral algorithms used to compute the collision operator. Several numerical results and comparisons to exact solutions as well as to Monte Carlo methods are given in section 4. An application to a challenging nonhomogeneous test case is finally given in section 5. Some final considerations close the paper in the last section.

2. The Boltzmann equation. The Boltzmann equation describes the behavior of a dilute gas of particles when the only interactions taken into account are binary

elastic collisions. It reads as $x \in \Omega$, $v \in \mathbb{R}^d$, where $\Omega \subset \mathbb{R}^d$ is the spatial domain ($d \geq 2$)

$$\frac{\partial f}{\partial t} + v \cdot \nabla_x f = Q(f, f),$$

where $f(t, x, v)$ is the time-dependent particles distribution function in the phase space. The Boltzmann collision operator Q is a quadratic operator local in (t, x) . The time and position act only as parameters in Q and therefore will be omitted in its description

$$(2.1) \quad Q(f, f)(v) = \int_{v_* \in \mathbb{R}^d} \int_{\sigma \in \mathbb{S}^{d-1}} B(|v - v_*|, \cos \theta) (f'_* f' - f_* f) \, d\sigma \, dv_*.$$

In (2.1) we used the shorthand $f = f(v)$, $f_* = f(v_*)$, $f' = f(v')$, $f'_* = f(v'_*)$. The velocities of the colliding pairs (v, v_*) and (v', v'_*) are related by

$$v' = v - \frac{1}{2}((v - v_*) - |v - v_*| \omega), \quad v'_* = v - \frac{1}{2}((v - v_*) + |v - v_*| \omega).$$

The collision kernel B is a nonnegative function which by physical arguments of invariance depends only on $|v - v_*|$ and $\cos \theta = \hat{g} \cdot \omega$ (where $\hat{g} = (v - v_*)/|v - v_*|$). In this work we are concerned with *short-range interaction* models. More precisely we assume that B is locally integrable. This assumption is satisfied by the *hard spheres (HS) model*, which in dimension $d = 3$ is written as

$$(2.2) \quad B(|v - v_*|, \cos \theta) = |v - v_*|$$

and is known as *Grad's angular cutoff assumption* when it is (artificially) extended to interactions deriving from power-law potentials. As an important benchmark model for the numerical simulation we therefore introduce the so-called *variable hard spheres (VHS) model*, which is written as

$$(2.3) \quad B(|v - v_*|, \cos \theta) = C_\gamma |v - v_*|^\gamma,$$

for some $\gamma \in [0, 1]$ and a constant $C_\gamma > 0$.

For this class of model, one can split the collision operator into

$$Q(f, f) = Q^+(f, f) - L(f) f,$$

with

$$(2.4) \quad Q^+(f, f) = \int_{\mathbb{R}^d} \int_{S^{d-1}} B(|v - v_*|, \cos \theta) f' f'_* \, d\sigma \, dv_*,$$

$$(2.5) \quad L(f) = \int_{\mathbb{R}^d} \int_{S^{d-1}} B(|v - v_*|, \cos \theta) f_* \, d\sigma \, dv_*.$$

Boltzmann's collision operator has the fundamental properties of conserving mass, momentum, and energy

$$\int_{v \in \mathbb{R}^d} Q(f, f) \phi(v) \, dv = 0, \quad \phi(v) = 1, v, |v|^2,$$

and satisfies the well-known Boltzmann's H theorem

$$-\frac{d}{dt} \int_{v \in \mathbb{R}^d} f \log f \, dv = - \int_{v \in \mathbb{R}^d} Q(f, f) \log(f) \, dv \geq 0.$$

The functional $-\int f \log f$ is the *entropy* of the solution. Boltzmann's H theorem implies that any equilibrium distribution function, i.e., any function which is a maximum of the entropy, has the form of a locally Maxwellian distribution

$$(2.6) \quad M(\rho, u, T)(v) = \frac{\rho}{(2\pi T)^{d/2}} \exp\left(-\frac{|u-v|^2}{2T}\right),$$

where ρ , u , T are the *density*, *mean velocity*, and *temperature* of the gas, defined, respectively, by

$$(2.7) \quad \rho = \int_{v \in \mathbb{R}^d} f(v) \, dv, \quad u = \frac{1}{\rho} \int_{v \in \mathbb{R}^d} v f(v) \, dv, \quad T = \frac{1}{d\rho} \int_{v \in \mathbb{R}^d} |u-v|^2 f(v) \, dv.$$

For further details on the physical background and derivation of the Boltzmann equation we refer to [13] and [49].

3. The spectral methods. In this section we shall explain the principles of the algorithms for computing the collision integral for a fixed value of the spatial variable x . Indeed, it is well known that one can reduce to this case by some splitting strategy (see [41, 20], for example).

3.1. A general framework. We consider the spatially homogeneous Boltzmann equation written in the general form

$$(3.1) \quad \frac{\partial f}{\partial t} = Q(f, f),$$

where Q is given by

$$(3.2) \quad Q(f, f) = \int_{\{(y,z) \in \mathcal{C}\}} \mathcal{B}(y, z) (f' f'_* - f_* f) \, dy \, dz, \quad v \in \mathbb{R}^d,$$

with

$$v' = v + \Theta'(y, z), \quad v'_* = v + \Theta'_*(y, z), \quad v_* = v + \Theta_*(y, z).$$

In the equations above, \mathcal{C} is some given unbounded domain, and Θ , Θ' , Θ'_* are suitable functions, to be defined later. This general framework emphasizes the translation invariance property of the collision operator, which is crucial for the spectral methods. We will explain more precisely in the next paragraphs some changes of variables which allow us to reduce the classical operator (2.1) to the form (3.2).

A problem associated with deterministic methods which use a fixed discretization in the velocity domain is that the velocity space is approximated by a finite region. Physically the domain for the velocity is \mathbb{R}^d , and the property of having compact support is not preserved by the collision operator. In general the collision process spreads the support by a factor of $\sqrt{2}$ in the elastic case (see [44, 31] and also [30] for similar properties in the inelastic case). As a consequence, for the continuous equation in time, the function f is immediately positive in the whole domain \mathbb{R}^d . Thus, at the numerical level, some nonphysical condition has to be imposed to keep the support of the function in velocity uniformly bounded. In order to do this there are two main strategies.

- One can remove the physical binary collisions that will lead outside the bounded velocity domain, which means a possible increase of the number of local invariants (that is, the functions φ such that $(\varphi'_* + \varphi' - \varphi_* - \varphi)$ is zero everywhere on the domain). If this is done properly (i.e., without removing too many collisions), the scheme remains conservative (and without spurious invariants). However, this truncation breaks down the convolution-like structure of the collision operator, which requires the translation invariance in velocity. Indeed, the modified collision kernel depends on v through the boundary conditions. This truncation is the starting point of most schemes based on DVM in a bounded domain.
- One can add some nonphysical binary collisions by periodizing the function and the collision operator. This implies the loss of some local invariants (some nonphysical collisions are added). Thus the scheme is no longer conservative, except for the mass if the periodization is done carefully. In this way the structural properties of the collision operator are maintained and thus they can be exploited to derive fast algorithms. This periodization is the basis of the spectral methods.

Therefore, we consider the space homogeneous Boltzmann equation in a bounded domain in velocity $\mathcal{D}_T = [T; T]^d$ ($0 < T < \infty$). We need to truncate the integration in y and z , since periodization would yield an infinite result if not. Thus we set y and z to belong to some truncated domain $\mathcal{C}_R \subset \mathcal{C}$ (the parameter R refers to its size and will be defined later). For a compactly supported function with support included in B_S , the ball centered at 0 with radius $S > 0$, one has to prescribe suitable relations (depending on the precise change of variable and truncation chosen) between S, R, T in order to retain all possible collisions and at the same time prevent intersections of the regions, where f is different from zero (dealiasing condition). Then the *truncated* collision operator reads

$$(3.3) \quad Q^R(f, f) = \int_{\mathcal{C}_R} \mathcal{B}(y, z) (f'_* f' - f_* f) dy dz$$

for $v \in \mathcal{D}_T$ (the expression for $v \in \mathbb{R}^d$ is deduced by periodization). By making some changes of variable on v , one can easily prove for the two choices of variables y, z in the next subsections that for any function φ periodic on \mathcal{D}_T the following weak form is satisfied:

$$(3.4) \quad \int_{\mathcal{D}_T} Q^R(f, f) \varphi(v) dv = \frac{1}{4} \int_{\mathcal{D}_T} \int_{\mathcal{C}_R} \mathcal{B}(y, z) f_* f (\varphi'_* + \varphi' - \varphi_* - \varphi) dy dz dv.$$

Now we use the representation Q^R to derive spectral methods. Hereafter we use just one index to denote the d -dimensional sums with respect to the vector $k = (k_1, \dots, k_d) \in \mathbb{Z}^d$, and hence we set

$$\sum_{k=-N}^N := \sum_{k_1, \dots, k_d=-N}^N .$$

The approximate function f_N is represented as the truncated Fourier series

$$(3.5) \quad f_N(v) = \sum_{k=-N}^N \hat{f}_k e^{i\frac{\pi}{T}k \cdot v},$$

$$\hat{f}_k = \frac{1}{(2T)^d} \int_{\mathcal{D}_T} f(v) e^{-i\frac{\pi}{T}k \cdot v} dv.$$

In a Fourier–Galerkin method the fundamental unknowns are the coefficients \hat{f}_k , $k = -N, \dots, N$. We obtain a set of ODEs for the coefficients \hat{f}_k by requiring that the residual of (3.3) be orthogonal to all trigonometric polynomials of degree less than N . Hence for $k = -N, \dots, N$

$$(3.6) \quad \int_{\mathcal{D}_T} \left(\frac{\partial f_N}{\partial t} - Q^R(f_N, f_N) \right) e^{-i\frac{\pi}{T}k \cdot v} dv = 0.$$

By substituting expression (3.5) in (3.4) we get

$$Q^R(f_N, f_N) = Q^{R,+}(f_N, f_N) - L^R(f_N) f_N,$$

with

$$(3.7) \quad L^R(f_N) f_N = \sum_{l=-N}^N \sum_{m=-N}^N \beta(m, m) \hat{f}_l \hat{f}_m e^{i\frac{\pi}{T}(l+m) \cdot v},$$

$$(3.8) \quad Q^{R,+}(f_N, f_N) = \sum_{l=-N}^N \sum_{m=-N}^N \beta(l, m) \hat{f}_l \hat{f}_m e^{i\frac{\pi}{T}(l+m) \cdot v},$$

where

$$(3.9) \quad \beta(l, m) = \int_{\mathcal{C}_R} \mathcal{B}(y, z) e^{i\frac{\pi}{T}(l \cdot \Theta'(y, z) + m \cdot \Theta'_*(y, z))} dy dz.$$

The *spectral equation* is the projection of the collision equation in \mathbb{P}_N , the $(2N + 1)^d$ -dimensional vector space of trigonometric polynomials of degree at most N in each direction, i.e.,

$$\frac{\partial f_N}{\partial t} = \mathcal{P}_N Q^R(f_N, f_N),$$

where \mathcal{P}_N denotes the orthogonal projection on \mathbb{P}_N in $L^2(\mathcal{D}_T)$. A straightforward computation leads to the following set of ODEs on the Fourier coefficients

$$(3.10) \quad \frac{\partial \hat{f}_k}{\partial t} = \sum_{\substack{l+m=k \\ l, m=-N}}^N \hat{\beta}(l, m) \hat{f}_l \hat{f}_m,$$

where $\hat{\beta}(l, m)$ are the so-called *kernel modes*, given by

$$\hat{\beta}(l, m) = \beta(l, m) - \beta(m, m),$$

with the initial condition

$$(3.11) \quad \hat{f}_k(0) = \frac{1}{(2T)^d} \int_{\mathcal{D}_T} f_0(v) e^{-i\frac{\pi}{T}k \cdot v} dv.$$

3.2. Classical spectral methods. In the classical spectral method [41], a simple change of variables in (2.1) permits us to write

$$(3.12) \quad Q(f, f) = \int_{\mathbb{R}^d} \int_{S^{d-1}} \mathcal{B}^c(g, \omega) (f(v')f(v'_*) - f(v)f(v_*)) \, d\omega \, dg,$$

with $g = v - v_* \in \mathbb{R}^d$, $\omega \in \mathbb{S}^{d-1}$, and

$$(3.13) \quad \begin{cases} v' = v - \frac{1}{2}(g - |g|\omega) =: v + \Theta'(g, \omega), \\ v'_* = v - \frac{1}{2}(g + |g|\omega) =: v + \Theta_*(g, \omega), \\ v_* = v + g =: v + \Theta_*(g, \omega). \end{cases}$$

Finally, \mathcal{B}^c is defined by

$$(3.14) \quad \mathcal{B}^c(g, \omega) = 2^{d-1} (1 - (\hat{g} \cdot \omega))^{d/2-1} B(|g|, 2(\hat{g} \cdot \omega)^2 - 1).$$

The Boltzmann operator (3.12) is now written in the form (3.2) with $(y, z) = (g, \omega) \in \mathbb{R}^d \times S^{d-1} =: \mathcal{C}$. Moreover, from the conservation of the momentum $v'_* + v' = v_* + v$ and the energy $|v'_*|^2 + |v'|^2 = |v_*|^2 + |v|^2$, we get the following results [40], assuming $\text{supp } f \subset B_S$:

- We have $\text{supp } Q(f, f) \subset B_{\sqrt{2}S}$;
- the collision operator is then given by

$$Q(f, f)(v) = \int_{B_{2S}} \int_{S^{d-1}} B(|g|, \cos \theta) (f(v')f(v'_*) - f(v_*)f(v)) \, d\omega \, dg,$$

with $v', v'_*, v_* \in B_{(2+\sqrt{2})R}$.

As a consequence of this result, in order to write a spectral approximation to (3.1) we consider the distribution function f restricted on $[-T, T]^d$ ($0 < T < +\infty$), assuming $f(v) = 0$ on $[-T, T]^d \setminus B_S$, and extend it by periodicity to a periodic function on $[-T, T]^d$. We truncate the domain for $(y, z) = (g, \omega)$ as $\mathcal{C}_R = B_R \times \mathbb{S}^{d-1}$ for $R > 0$ (thus defining Q^R). Following the previous discussion on the dealiasing condition, we take $R = 2S$, and the shortest period can be restricted to $[-T, T]^d$, with $T \geq (3 + \sqrt{2})S/2$ (for a more detailed discussion see [41]).

Then we apply the spectral algorithm (3.7), (3.8) and get the *kernel modes* $\beta^c(l, m)$

$$(3.15) \quad \beta^c(l, m) = \int_{B_R} \int_{S^{d-1}} B(|g|, \cos \theta) e^{-i \frac{\pi}{T} (g \cdot \frac{(l+m)}{2} - i|g|\omega \cdot \frac{(m-l)}{2})} \, d\omega \, dg.$$

We refer to [41, 22] for the explicit computation of Fourier coefficients $\beta^c(l, m)$ in the VHS case, where B is given by (2.3). Now the evaluation of the right-hand side of (3.10) requires exactly $O(N^{2d})$ operations. We emphasize that the usual cost for a DVM method based on N^d parameters for f in the velocity space is $O(N^{2d}M)$, where M is the numbers of angle discretizations.

3.3. Fast spectral methods. Here we shall approximate the collision operator starting from a representation which conserves more symmetries of the collision operator when one truncates it in a bounded domain. This representation was used

in [7, 26] to derive finite differences schemes and is close to the classical Carleman representation (cf. [12]). The basic identity we shall need is (for $u \in \mathbb{R}^d$)

$$(3.16) \quad \frac{1}{2} \int_{S^{d-1}} F(|u|\sigma - u) d\sigma = \frac{1}{|u|^{d-2}} \int_{\mathbb{R}^d} \delta(2y \cdot u + |y|^2) F(y) dy.$$

Using (3.16) the collision operator (2.1) can be written as

$$(3.17) \quad Q(f, f)(v) = 2^{d-1} \int_{x \in \mathbb{R}^d} \int_{y \in \mathbb{R}^d} \mathcal{B}^f(y, z) \delta(y \cdot z) \times (f(v + z)f(v + y) - f(v + y + z)f(v)) dy dz,$$

with

$$\mathcal{B}^f(y, z) = 2^{d-1} B \left(|y + z|, -\frac{y \cdot (y + z)}{|y||y + z|} \right) |y + z|^{-(d-2)}.$$

Thus the collision operator is now written in the form (3.2), with $(y, z) \in \mathbb{R}^d \times \mathbb{R}^d =: \mathcal{C}$, $\mathcal{B}(y, z) = \mathcal{B}^f(y, z) \delta(y \cdot z)$, and $v'_* = v + z =: v + \Theta'_*(y, z)$, $v' = v + y =: v + \Theta'(y, z)$, $v_* = v + y + z =: v + \Theta_*(y, z)$.

Now we consider the bounded domain $\mathcal{D}_T = [T, T]^d$ ($0 < T < \infty$) for the distribution f , and the bounded domain $B_R \times B_R$ for (y, z) (for some $R > 0$). If f has support included in B_S , $S > 0$, geometrical arguments similar to the one for the classical spectral methods (see [41, 33, 32]) show that we can take $R = 2S$ and $T \geq (1 + 3\sqrt{2})S/2$ to get all collisions and prevent intersections of the regions, where f is different from zero. The (truncated) operator now reads

$$(3.18) \quad Q^R(f, f)(v) = \int_{y \in B_R} \int_{z \in B_R} \mathcal{B}^f(y, z) \delta(y \cdot z) (f(v + z)f(v + y) - f(v + y + z)f(v)) dy dz$$

for $v \in \mathcal{D}_T$. This representation of the collision kernel yields better decoupling properties between the arguments of the operator. From now on we can apply the spectral algorithm (3.7), (3.8) to this collision operator, and the corresponding kernel modes are given by

$$\beta^f(l, m) = \int_{y \in B_R} \int_{z \in B_R} \tilde{B}(y, z) \delta(y \cdot z) e^{i\frac{\pi}{T}(l \cdot y + m \cdot z)} dy dz.$$

In what follows we shall focus on β^f , and one easily checks that $\beta^f(l, m)$ depends only on $|l|$, $|m|$, and $|l \cdot m|$.

Remark 3.1. Note that the classical spectral method originates the following form of the kernel modes in the y, z notation:

$$\beta^c(l, m) = \int_{y \in B_R} \int_{z \in B_R} \mathcal{B}^f(y, z) \delta(y \cdot z) \chi_{|y+z| \leq R} e^{i\frac{\pi}{T}(l \cdot y + m \cdot z)} dy dz.$$

One can notice that the condition $|y + z|^2 = |y|^2 + |z|^2 \leq R^2$ couples the modulus of y and z such that the ball is not completely covered (for instance, if y and z are orthogonal both with modulus R , the condition is not satisfied, since $|y + z| = \sqrt{2R}$). This explains the better decoupling properties between the arguments of the collision operator of this representation.

4. Fast algorithms. The search for fast deterministic algorithms for the collision operator, i.e., algorithms with a cost lower than $O(N^{2d+\epsilon})$ (with typically $\epsilon = 1$ for DVM, or $\epsilon = 0$ for the classical spectral method), consists mainly of identifying some convolution structure in the operator (see, for example, [8, 42]). The aim is to approximate each $\beta^f(l, m)$ by a sum

$$(4.1) \quad \beta^f(l, m) = \sum_{p=1}^A \alpha_p(l) \alpha'_p(m).$$

This gives a sum of A discrete convolutions, and thus the algorithm can be computed in $O(AN^d \log_2 N)$ operations by means of standard fast Fourier transform (FFT) techniques [11]. To this purpose we shall use a further approximated collision operator, where the number of possible directions of collision is reduced to a finite set. We start from representation (3.18) and write y and z in spherical coordinates

$$Q^R(f, f)(v) = \frac{1}{4} \int_{e \in S^{d-1}} \int_{e' \in S^{d-1}} \delta(e \cdot e') \, de \, de' \\ \times \left[\int_{-R}^R \int_{-R}^R \rho^{d-2} (\rho')^{d-2} \mathcal{B}^f(\rho, \rho') (f(v + \rho' e') f(v + \rho e) - f(v + \rho' e' + \rho e) f(v)) \, d\rho \, d\rho' \right]$$

(note that thanks to the orthogonality condition imposed by the Dirac mass on y and z , \mathcal{B}^f depends only on the modulus of y and z). Let us denote by \mathcal{A} a discrete set of orthogonal couples of unit vectors (e, e') , which is even; i.e., $(e, e') \in \mathcal{A}$ implies that $(-e, e')$, $(e, -e')$, and $(-e, -e')$ belong to \mathcal{A} (this property on the set \mathcal{A} is required to preserve the conservation properties of the operator). Now we define $Q^{R, \mathcal{A}}$ to be

$$Q^{R, \mathcal{A}}(f, f)(v) = \frac{1}{4} \int_{(e, e') \in \mathcal{A}} d\mathcal{A} \\ \times \left[\int_{-R}^R \int_{-R}^R \rho^{d-2} (\rho')^{d-2} \mathcal{B}^f(\rho, \rho') (f(v + \rho' e') f(v + \rho e) - f(v + \rho' e' + \rho e) f(v)) \, d\rho \, d\rho' \right],$$

where $d\mathcal{A}$ denotes a discrete measure on \mathcal{A} , which is also even in the sense that $d\mathcal{A}(e, e') = d\mathcal{A}(-e, e') = d\mathcal{A}(e, -e') = d\mathcal{A}(-e, -e')$. It is easy to check that $Q^{R, \mathcal{A}}$ has the same conservation properties as Q^R . We make the decoupling assumption that

$$(4.2) \quad \forall y \perp z, \quad \mathcal{B}^f(y, z) = a(|y|) b(|z|).$$

This assumption is obviously satisfied if \mathcal{B}^f is constant. This is the case of Maxwellian molecules in dimension $d = 2$ and hard spheres in dimension $d = 3$ (the most relevant kernel for applications). Extensions to more general interactions are discussed in [33].

Let us describe the method in dimension $d = 3$ with \mathcal{B}^f satisfying the decoupling assumption (4.2) (see [33] for other dimensions). First we change to spherical coordinates

$$\beta^f(l, m) = \frac{1}{4} \int_{e \in S^2} \int_{e' \in S^2} \delta(e \cdot e') \\ \times \left[\int_{-R}^R |\rho| a(\rho) e^{i \frac{\pi}{2} \rho(l \cdot e)} \, d\rho \right] \left[\int_{-R}^R |\rho'| b(\rho') e^{i \frac{\pi}{2} \rho'(m \cdot e')} \, d\rho' \right] \, de \, de'$$

and then we integrate first e' on the intersection of the unit sphere with the plane e^\perp

$$\beta^f(l, m) = \frac{1}{4} \int_{e \in S^2} \phi_{R,a}^3(l \cdot e) \left[\int_{e' \in S^2 \cap e^\perp} \phi_{R,b}^3(m \cdot e') de' \right] de,$$

where

$$\phi_{R,a}^3(s) = \int_{-R}^R |\rho| a(\rho) e^{i \frac{\pi}{T} \rho s} d\rho, \quad \phi_{R,b}^3(s) = \int_{-R}^R |\rho| b(\rho) e^{i \frac{\pi}{T} \rho s} d\rho.$$

Thus we get the following decoupling formula with two degrees of freedom:

$$\beta_R(l, m) = \int_{e \in S_+^2} \phi_{R,a}^3(l \cdot e) \psi_{R,b}^3(\Pi_{e^\perp}(m)) de,$$

where S_+^2 denotes the half-sphere and

$$\psi_{R,b}^3(\Pi_{e^\perp}(m)) = \int_0^\pi \phi_{R,b}^3(|\Pi_{e^\perp}(m)| \cos \theta) d\theta,$$

and Π_{e^\perp} is the orthogonal projection on the plane e^\perp . In the particular case where $\mathcal{B}^f = 1$ (HS model), we can compute the functions ϕ_R^3 and ψ_R^3

$$\phi_R^3(s) = R^2 (2 \operatorname{Sinc}(Rs) - \operatorname{Sinc}^2(Rs/2)), \quad \psi_R^3(s) = \int_0^\pi \phi_R^3(s \cos \theta) d\theta.$$

Now the function $e \mapsto \phi_{R,a}^3(l \cdot e) \psi_{R,b}^3(\Pi_{e^\perp}(m))$ is periodic on S_+^2 . Taking a spherical parametrization (θ, φ) of $e \in S_+^2$ and taking for the set \mathcal{A} uniform grids of respective size M_1 and M_2 for θ and φ we get

$$\beta^f(l, m) \simeq \frac{\pi^2}{M_1 M_2} \sum_{p,q=0}^{M_1, M_2} \alpha_{p,q}(m) \alpha'_{p,q}(l),$$

where

$$\alpha_{p,q}(l) = \phi_{R,a}^3(l \cdot e_{\theta_p, \varphi_q}), \quad \alpha'_{p,q}(m) = \psi_{R,b}^3\left(\Pi_{e_{\theta_p, \varphi_q}^\perp}(m)\right)$$

and $(\theta_p, \varphi_q) = (p\pi/M_1, q\pi/M_2)$.

We consider this expansion with $M = M_1 = M_2$ to avoid anisotropy in the computational grid. By using the FFT algorithm, the computational cost of the algorithm is then $O(M^2 N^3 \log N)$ (compared to $O(M^2 N^6)$ of a direct discretization on the grid for a DVM method, and to $O(N^6)$ of the classical spectral method).

Let us finally mention that the mathematical analysis of the fast algorithm in [33] provides the following results:

- It is spectrally accurate according to the parameters N and M ;
- the error on the conservation laws of momentum and energy is spectrally small according to the parameter N , and no additional error (according to the speed-up parameter M) is made.

These two properties were the main motivations for the development of the method of [33] described above to obtain the decomposition (4.1). Other advantages of this particular decomposition is that it does not introduce instability in the equation (see [33, Theorem 3.1], for instance) and it is naturally adaptive (as it is based on the rectangular quadrature rule for approximating integrals of periodic functions). Finally, another advantage of the proposed method is that it is still easy to implement since it is based only on FFT.




FIG. 1. Two-dimensional homogeneous case: evolution of the numerical L^1 and L^∞ relative error of $f(t, v)$.

5. Numerical results in the homogeneous case. In this section we will present several numerical results for the space homogeneous equation which show the improvement of the fast spectral algorithms with respect to the classical spectral methods and how they compare with Monte Carlo methods. The time discretization is performed by suitable Runge–Kutta methods.

5.1. Spatially homogeneous Maxwell molecules in dimension 2.

Comparison to exact solutions. We consider two-dimensional pseudo-Maxwell molecules (i.e., the VHS model with $\gamma = 0$). In this case we have an exact solution given by

$$f(t, v) = \frac{\exp(-v^2/2S)}{2\pi S^2} \left[2S - 1 + \frac{1 - S}{2S} v^2 \right],$$

with $S = 1 - \exp(-t/8)/2$, which corresponds to the well-known exact solution [4]. This test is performed to check spectral accuracy, by comparing the error at a given time, when using $n_v = 8, 16,$ and 32 Fourier modes for each coordinate. We present the results obtained by the classical spectral method and the fast spectral method with different numbers of discrete angles.

Figure 1 shows the relative $L^\infty, L^1,$ and L^2 norms of the difference between the numerical and exact solutions as functions of time. These errors are computed according to the formula

$$\mathcal{E}_p = \left(\frac{\sum_{i=-N}^N |f_i(t) - f(v_i, t)|^p}{\sum_{i=-N}^N |f(v_i, t)|^p} \right)^{1/p},$$

with $i = (i_1, i_2)$ and $N = n_v/2$ for $p = 1$ and $p = 2$. A similar expression is used for the L^∞ error. Note that the error increases initially and then decreases almost monotonically in time. After a long time the error starts increasing again. This effect is due to aliasing. Indeed, for a fixed computational domain, when the number of Fourier modes increases, the effect of aliasing becomes dominant over the error due to the spectral approximation. For this reason, the size of the domain is chosen in order to minimize the aliasing error. A trade-off should be obtained between aliasing and spectral error, which means that the size of the domain should be increased when

TABLE 1

Comparison of the L^1 error in two dimensions between the classical spectral method and the fast spectral method with different numbers of discrete angles and with a second-order Runge–Kutta time discretization at time $T_{end} = 1$.

Number of points	Classical spectral	Fast spectral with $M = 4$	Fast spectral with $M = 6$	Fast spectral with $M = 8$
8	0.02013	0.02778	0.02129	0.02112
16	0.00204	0.00329	0.00238	0.00224
32	1.405E-5	2.228E-5	1.861E-5	1.772E-5

increasing the number of Fourier modes. Roughly speaking, the period should be chosen in such a way that the two contributions of the error are of the same order of magnitude. In this test, the radius of the ball, which defines the computational domain, is $T = 4$ for $n_v = 8$, $T = 5$ for $n_v = 16$, and $T = 7$ for $n_v = 32$. We refer to [41] for a more detailed discussion about aliasing.

Concerning the comparison between the classical and fast spectral methods, we observe that for a fixed value of n_v , the numerical error of the classical spectral method and of the fast algorithm is of the same order. Moreover, the influence of the number of discrete angles is very weak. Indeed, with only $M = 4$, the results are quite similar even for large n_v and as expected the number of discrete angles does not affect the variations of energy, which are of the same order of magnitude as the numerical error (note that there is no variation for the momentum since in the special case of even solutions, it is preserved at 0 by the spectral scheme). In Table 1, we give a quantitative comparison of the numerical error \mathcal{E}_1 at time $T_{end} = 1$. We can also observe the spectral accuracy for the classical and fast methods: the order of accuracy is about 3 between 8 and 16 grid points, whereas it becomes 7 between 16 to 32 points.

Efficiency and accuracy. Now we still consider two-dimensional pseudo-Maxwell molecules (i.e., $\gamma = 0$) with the initial datum

$$f(0, v) = \frac{1}{4\pi} \left[\exp\left(-\frac{|v - v_0|^2}{2}\right) + \exp\left(-\frac{|v + v_0|^2}{2}\right) \right], \quad v \in \mathbb{R}^2,$$

where $v_0 = (1, 2)$. In this case, we do not know the exact solution but we want to study the influence of the number of discrete angles on a nonisotropic solution. Thus, this test is used to check the energy conservation and the evolution of high-order moments of the solution.

The time step is chosen small enough to reduce the influence of the time discretization, i.e., $\Delta t = 0.025$. Moreover, the computational domain is taken large enough with respect to the number of grid points in order to reduce the aliasing error due to the periodization of the solution. Simulations are performed with $n_v = 16, 32$, and 64 points.

In Figure 2 the relaxation of the entropy and the temperature components for the fast and classical spectral methods is shown. The energy is conserved by the continuous collision operator, but when using the spectral method the total energy can change with time, which is a good indicator of the accuracy of the numerical solution. Indeed, the total discrete energy is not exactly conserved over time, but if the aliasing error is small, it is conserved within spectral accuracy; typically here the variations are about 10^{-4} when $n_v = 32$. Moreover, we observe that the number of discrete angles does not significantly affect the transient regime. For instance, with only four angles




FIG. 2. Two-dimensional homogeneous case: relaxation of the entropy and the temperature components for the fast and classical spectral methods with respect to the number of modes per direction n_v and the length box T .




FIG. 3. Two-dimensional homogeneous case: time evolution of the variations of high-order normalized moments M_4 , M_5 , and M_6 of $f(t, v)$ for the fast and classical spectral methods with respect to the number of modes per direction n_v and the length box T .

on the half-sphere, the relaxation of entropy and temperature components is very close to the numerical solution obtained by the classical spectral method. Finally, we plot in Figure 3 the time evolution of high-order moments of $f_N(t, v)$ given in discrete form by

$$\mathcal{M}_k(t) = \Delta v^2 \sum_{l=-N}^N |v_l|^k f_N(t, v_l).$$

High-order moments give information on the accuracy of the approximate distribution function tail. Once again, we observe that the number of angles does not affect the results even if the solution is nonisotropic.

To conclude, we observe that in dimension $d = 2$, the fast algorithm is really efficient in terms of accuracy and computational cost compared to the classical spectral method. In Table 2 we report the computational times of the methods which show a speed-up of the fast solver independently of the number of points used in our tests and with a maximum speed-up reached for $N = 64$, where the fast methods with $M = 4$ is more than 17 times faster than the classical method.

5.2. Spatially homogeneous hard spheres in dimension 3. In this section we consider the three-dimensional HS molecules model. The initial condition is chosen

TABLE 2

Comparison of the computational time in two dimensions between the classical spectral method and the fast spectral method with different numbers of discrete angles and with a second-order Runge–Kutta time discretization.

Number of points	Classical spectral	Fast spectral with $M = 4$	Fast spectral with $M = 6$	Fast spectral with $M = 8$
16	2 sec. 40	1 sec. 15	1 sec. 70	2 sec. 30
32	38 sec. 01	5 sec. 55	8 sec. 47	11 sec. 10
64	616 sec.	35 sec. 50	54 sec. 66	71 sec. 27

TABLE 3

Comparison of the computational time in three dimensions between the classical spectral method and the fast spectral method with different numbers of discrete angles and with a second-order Runge–Kutta time discretization.

Number of points	Classical spectral	Fast spectral with $M = 4$	Fast spectral with $M = 6$	Fast spectral with $M = 8$
16	1 min. 14 sec.	3 min. 31 sec.	7 min. 45 sec.	13 min. 44 sec.
32	118 min. 02 sec.	50 min. 31 sec.	105 min. 19 sec.	186 min. 18 sec.
64	125 h 54 min.	8 h 45 min. 22 sec.	21 h 39 min.	35 h 01 min. 28 sec.

as the sum of two Gaussians

$$f(v, 0) = \frac{1}{2(2\pi\sigma^2)} \left[\exp\left(-\frac{|v - v_0|^2}{2\sigma^2}\right) + \exp\left(-\frac{|v + v_0|^2}{2\sigma^2}\right) \right],$$

with $\sigma = 1$ and $v_0 = (2, 1, 0)$. The final time of the simulation is $T_{end} = 3$ and corresponds approximately to the time for which the steady state of the solution is reached. The time step is $\Delta t = 0.1$ and the length box is taken as $T = 12$ when $n_v = 16$ and as $T = 15$ when $n_v = 32$.

This test is used to check the evolution of moments and particularly the stress tensor $P_{i,j}$, $i, j = 1, \dots, 3$ defined as

$$P_{i,j} = \int_{\mathbb{R}^3} f(v)(v_i - u_i)(v_j - u_j) dv, \quad (i, j) \in \{1, 2, 3\}^2,$$

where $(u_i)_i$ are the components of the mean velocity. As in the previous case, we compare the classical and fast methods in terms of computational time (see Table 3) and accuracy. In Figure 4, we propose the evolution of the temperature for the two methods using 32 grid points in each direction. This solution is also compared with the solution obtained from the Monte Carlo method. The discrete temperatures agree well in this case and the efficiency of the fast algorithm is verified since the computational time is highly reduced using only $M_1 = M_2 = 4$ discrete angles without affecting the accuracy of the distribution function. We remark that in dimension $d = 3$ the speed-up of the methods becomes really evident for large values of N . Again for $N = 64$ and $M = 4$ the fast method is more than 14 times faster.

Comparison with Monte Carlo. Finally, we compare the results obtained with our spectral method with those obtained by a Monte Carlo scheme. We use a standard version of Monte Carlo, which may be referred to as the Nanbu–Babovsky scheme [36, 2].

In the case of Monte Carlo methods, the moments are computed by using unbiased estimators averaged over several runs. The number of runs N_{runs} , number of particles N_p , and the time step Δt , have been chosen in such a way to balance the time discretization error with statistical fluctuations.




FIG. 4. Three-dimensional homogeneous case: comparison between the fast and classical spectral methods and the Monte Carlo method for the temperature components relaxation.

The first run consists of taking a large number of particles and making time averaging in order to minimize the fluctuation errors: $N_{\text{runs}} = 10^3$, $N_p = 10^4$, and $\Delta t = 0.01$. The total computational time for computing the evolution of moments in this case is 113 min. 23 sec. On the other hand we perform a second run, where we use more averaging to minimize fluctuation errors: $N_{\text{runs}} = 5 \cdot 10^3$, $N_p = 5 \cdot 10^3$, and $\Delta t = 0.01$. The computational time is now 100 min.

We remark that the computations have been obtained by using the HS model (VHS with $\gamma = 1$), which is the most realistic. In this case, the computational time of the Monte Carlo methods becomes larger than in the case when we consider pseudo-Maxwell molecules (VHS with $\gamma = 0$) for which the collision kernel is constant and no rejection is needed. For the spectral method the computational cost is independent of the collision kernel.

From the comparison, it is obvious that, for three-dimensional computations, the greater cost of the classical spectral scheme (with the same number of degrees of freedom) is compensated by a much greater accuracy, allowing for better results with the same computational cost. Moreover, with the fast algorithm, the spectral scheme really becomes competitive in terms of computational time since the accuracy is not affected when we use fewer discrete angles (for instance, $M = 4$). In Figure 5, we compare the accuracy on the evolution of high-order moments with the different methods: the fourth-order moments are very close, but the results obtained with the Monte Carlo methods are affected by fluctuations on the tail of the distribution function, which are difficult to remove (see the evolution of the eighth-order moment). Note that for two- and three-dimensional pseudo-Maxwell molecules, comparisons had also been performed between Monte Carlo methods and the classical spectral method in [41, section 6.3].

5.3. Stability of spectral methods with respect to nonsmooth data. In

this subsection we perform some numerical simulations in order to study the behavior of the spectral methods when applied to nonsmooth data. We consider the distribution

$$f_0(v) = \begin{cases} 1 & \text{if } |v|^2 \leq 1, \\ 0 & \text{else} \end{cases}$$

and use a mollified initial datum, which suitably approximates moments. We perform two numerical simulations using the fast spectral method with 32^2 and 64^2 grid points and plot the evolution of the entropy and the fourth-order moment in Figure 6. Even




FIG. 5. Three-dimensional homogeneous case: time evolution of the kinetic entropy H and high-order moments M_4 , M_6 , and M_8 of $f(t, v)$ for the fast and classical spectral methods and the Monte Carlo methods.




FIG. 6. Time evolution of the kinetic entropy and the fourth-order moment with respect to the number of grid points 32^2 and 64^2 .

for this discontinuous initial datum, we observe that for the two configurations the numerical entropy is decreasing and both numerical solutions converge to the same steady state. Moreover, we plot the evolution of the distribution function with respect to time in Figure 7: the fast spectral method is very stable for this numerical test even if spurious oscillations are first generated; the distribution becomes smooth and converges to an approximated Maxwellian. As expected from a Fourier-Galerkin method, the accuracy degenerates in the discontinuity region. However, surprisingly the method seems to remain stable.




FIG. 7. Time evolution of the distribution function for 64×64 grid points.

Finally, let us mention that we have also performed some numerical tests when the initial datum approximates a Dirac distribution. The spectral method is still stable even if spurious oscillations are generated, but this problem is inherent to gridded methods and we refer to [23] for a rescaling method, which follows the variation of the distribution function and allows us to treat concentrated distributions.

6. Application to the nonhomogeneous case. In [20, 22], we performed several numerical simulations to compare the spectral scheme with Monte Carlo methods and showed that when we are interested in the transient regime, the deterministic method becomes very efficient. Obviously, the fast spectral method will still improve the computational cost.

In what follows, since we will be interested in the study of the trend to a global equilibrium state of the kinetic equation, we avoid the use of a splitting method by solving the whole nonhomogeneous equation in time by a second-order Runge–Kutta method. Clearly the spectral methods apply straightforwardly to the collision operator also in this situation. The transport part is treated by the positive flux conservative method (see [18, 21, 20] for further details).

6.1. Definition of the problem. We consider the full Boltzmann equation in dimension $d = 2$ on the torus

$$\frac{\partial f}{\partial t} + v \cdot \nabla_x f = Q(f, f), \quad x \in [0, L]^2, \quad v \in \mathbb{R}^2,$$

with periodic boundary conditions in x . We first introduce the hydrodynamical fields associated with a kinetic distribution $f(t, x, v)$. These are the $(d + 2)$ scalar fields of density ρ (scalar), mean velocity u (vector valued), and temperature T (scalar) defined by the formulas (2.7). Whenever $f(t, x, v)$ is a smooth solution to the Boltzmann equation with periodic boundary conditions, one has the global conservation laws for

mass, momentum, and energy

$$\begin{aligned}\frac{d}{dt} \int_{[0,L]^2 \times \mathbb{R}^2} f(t, x, v) \, dx \, dv &= 0, \\ \frac{d}{dt} \int_{[0,L]^2 \times \mathbb{R}^2} f(t, x, v) v \, dx \, dv &= 0, \\ \frac{d}{dt} \int_{[0,L]^2 \times \mathbb{R}^2} f(t, x, v) \frac{|v|^2}{2} \, dx \, dv &= 0.\end{aligned}$$

Therefore, without loss of generality we shall impose

$$\int_{[0,L]^2 \times \mathbb{R}^2} f(t, x, v) \, dx \, dv = 1, \quad \int_{[0,L]^2 \times \mathbb{R}^2} f(t, x, v) v \, dx \, dv = 0,$$

and

$$\int_{[0,L]^2 \times \mathbb{R}^2} f(t, x, v) \frac{|v|^2}{2} \, dx \, dv = 1.$$

These conservation laws are then enough to uniquely determine the stationary state of the Boltzmann equation: the normalized global Maxwellian distribution

$$(6.1) \quad M_g(v) = \frac{1}{2\pi} \exp\left(-\frac{|v|^2}{2}\right).$$

We shall use the following terminology: a velocity distribution of the form (6.1) will be called a *Maxwellian distribution*, whereas a distribution of the form

$$(6.2) \quad M_l(x, v) = \frac{\rho(x)}{2\pi T(x)} \exp\left(-\frac{|v - u(x)|^2}{2T(x)}\right)$$

will be called a *local Maxwellian distribution* (in the sense that the constants ρ , u , and T appearing there depend on the position x). We also define the notion of *relative local entropy* H_l , the entropy relative to the local Maxwellian, and the *relative global entropy* H_g , the entropy relative to the global Maxwellian distribution, respectively, by

$$H_l(t) = \int f \log\left(\frac{f}{M_l}\right) \, dx \, dv, \quad H_g(t) = \int f \log\left(\frac{f}{M_g}\right) \, dx \, dv.$$

Our goal here is to investigate numerically the long-time behavior of the solution f . If f is any reasonable solution of the Boltzmann equation, satisfying certain a priori bounds of compactness (in particular, ensuring that no kinetic energy is allowed to leak at large velocities), then it is possible to prove that f does indeed converge to the global Maxwellian distribution M_g as t goes to $+\infty$. Of course, obtaining these a priori bounds is extremely difficult; as a matter of fact, they have been established only in the spatially homogeneous situation (which means that the distribution function does not depend on the position variable x ; see the survey in [49]) or in a close-to-equilibrium setting (see in particular [25] for the torus), and it still constitutes a famous open problem for spatially inhomogeneous initial data far from equilibrium. More recently, Desvillettes and Villani [16] and Strain and Guo [47] were interested in the study of rates of convergence for the full Boltzmann equation. Roughly speaking,

in [16] it was proved that if the solution to the Boltzmann equation is smooth enough and satisfies bounds from below of the form

$$\forall t \geq 0, x \in [0, L]^2, v \in \mathbb{R}^2, \quad f(t, x, v) \geq K_0 e^{-A_0 |v|^{q_0}} \quad (A_0, K_0 > 0, q_0 \geq 2),$$

(although this bound can be shown to be a consequence of the regularity bounds; see [31]), then (with constructive bounds)

$$\|f(t) - M_g\| = O(t^{-\infty}),$$

which means that the solution converges almost exponentially fast to the global equilibrium (namely, with polynomial rate $O(t^{-r})$ with r as large as desired).

The solution f to the Boltzmann equation satisfies the formula of additivity of the entropy: the entropy can be decomposed into the sum of a purely hydrodynamic part and (by contrast) of a purely kinetic part. In terms of the H functional, one can write

$$H_g(t) = H_l(t) + \int_0^L \rho_l(t, x) \log \left(\frac{\rho_l(t, x)}{T_l(t, x)} \right) dx.$$

In fact, we can also show that

$$H_l(t) \leq H_g(t) \quad \forall t \geq 0.$$

Moreover, the Csiszár–Kullback–Pinsker inequality asserts that (when the total mass of the solution is normalized to 1)

$$H(f|M) \geq \frac{1}{2} \|f - M\|_{L^1}^2.$$

In other words, controlling the speed of convergence of the entropy to its equilibrium value is enough to control the speed of convergence of the solution to its equilibrium in a very strong sense.

Moreover, in [16] Desvillettes and Villani conjectured that time oscillations should occur on the evolution of the relative local entropy. In fact their proof does not rule out the possibility that the entropy production undergoes important oscillations in time, and actually most of the technical work is caused by this possibility.

6.2. Description and interpretation of the results. Here we performed simulations on the full Boltzmann equation in a simplified geometry (one dimension of space, two dimensions of velocity, periodic boundary conditions, fixed Knudsen number) with the fast spectral method to observe the evolution of the entropy and to check numerically if such oscillations occur. Clearly this test is challenging for a numerical method due to the high accuracy required to capture such oscillating behavior.

Then we considered an initial datum as a perturbation of the global equilibrium M_g

$$(6.3) \quad f_0(x, v) = \frac{1}{2\pi} (1 + A_0 \cos(k_0 x)) \exp(-|v|^2/2), \quad x \in [0, L], \quad v \in \mathbb{R}^2,$$

for some constants $A_0 > 0$ and $k_0 = 2\pi/L$.

In Figures 8 and 9, we are indeed able to observe oscillations in the entropy production and in the hydrodynamic entropy. The strength of the oscillations depends a lot on the length L of the domain, which is consistent with the fact that




FIG. 8. Influence of the length box: relative entropy with respect to the local Maxwellian $H_I(t)$ using $64 \times 64 \times 64$ for $L = \pi, 2\pi, 3\pi,$ and 4π with $A_0 = 0.1$ in (6.3).




FIG. 9. Influence of the length box: relative entropy with respect to the local Maxwellian $H_I(t)$ using $64 \times 64 \times 64$ for $L = 2\pi, 3\pi,$ and 4π with $A_0 = 0.1$ in (6.4).

such oscillations are never observed in the spatially homogeneous case ($L = 0$). The superimposed curves yield the time evolution of the total H functional and of its kinetic part, respectively. In all cases, a local Maxwellian distribution is chosen for the initial datum; the first plot corresponds to $L = 1$ and the second one to $L = 4$. Some slight oscillations can be seen in the case $L = 1$, but what is most striking is that after a short while, the kinetic entropy is very close to the total entropy, which is an indication that the solution evolves basically in a spatially homogeneous way (contrary to the intuition of the hydrodynamic regime). On the contrary, in the case $L = 4$, the oscillations are much more important in frequency and amplitude (note that this is a logarithmic plot): the solution “hesitates” between states in which it

TABLE 4

Influence of the length box: damping rate and oscillation frequency for the relative entropy with respect to the local Maxwellian $H_1(t)$ using $64 \times 64 \times 64$ with $A_0 = 0.1$ in (6.3).

Length box	Oscillation frequency ω	ω/L	Damping rate α	$-\alpha L^2$
$L = \pi/2$	01.10	0.701	-6.521	16.04
$L = \pi$	02.25	0.716	-2.202	21.71
$L = 2\pi$	04.50	0.716	-0.641	25.26
$L = 3\pi$	06.61	0.701	-0.285	25.31
$L = 4\pi$	08.78	0.699	-0.160	25.27
$L = 8\pi$	17.57	0.699	-0.040	25.35

is very close to hydrodynamic and states in which it is not at all. Further note that the equilibration is much more rapid when the box is small and that the convergence seems to be exponential.

It is in fact possible to give a simple interpretation of these oscillations thanks to the work [17]. Since this effect is observed near the global equilibrium one can replace the Boltzmann collision operator with the linearized Boltzmann collision operator (moreover, oscillation is effectively observed for the linearized Boltzmann collision operator as well). Then it is straightforward that the computations above correspond to observing the time evolution of one Fourier mode in x (here with frequency k_0 and amplitude A_0). Hence by an obvious rescaling, this evolution is given by the semigroup $\tilde{T}_{k_0/L}(t)$, where $\tilde{T}_k(t)$ is defined in [17] (this is the semigroup for the k th Fourier mode in x for the linearized equation). An asymptotic study of the spectrum of its infinitesimal generator for small frequencies k was done in [17]. The dominant term in terms of long-time behavior (i.e., the one with the lower rate of decrease) is given by the $(d+2)$ “hydrodynamical eigenvalues.” Moreover, explicit computations are available for the expansions of these eigenvalues according to $\varepsilon = |k|$ near $k = 0$.

At first order in ε , the eigenvalues vanish, except for two of them, which are purely imaginary. They are given by

$$\begin{cases} I_1 = i\varepsilon \sqrt{1 + 2/d} + O(\varepsilon^2), \\ I_2 = -i\varepsilon \sqrt{1 + 2/d} + O(\varepsilon^2), \\ I_3 = \dots = I_{d+2} = 0. \end{cases}$$

Therefore for $|k_0|/L \ll 1$ (realized, for instance, when k_0 is fixed and L is large enough), this analysis gives us the dominant imaginary term in the eigenvalues. In this regime, one should thus observe oscillations with frequency $\sqrt{1 + 2/d} |k_0|/L$. Thus, the period of oscillations should be given by $2\pi(1 + 2/d)^{-1/2} L/|k_0|$, which can be checked with the numerical simulations. Indeed, in Table 4 we give the ratio of the period of oscillations ω with the length box L . The numerical results agree well with the analytical computations $\omega/L \simeq 1/\sqrt{2}$.

We also observe that the damping rate is related to the length box and is proportional to $1/L^2$ when L becomes large (see Table 4; $\alpha L^2 \simeq \text{constant}$). This is coherent with the fact that no real value occurs in the hydrodynamical eigenvalues until the second order in $\varepsilon = |k|$. The coefficients for the order 2 in the expansion are computed in [17]; they are purely real and can be expressed simply in terms of the dimension d , the viscosity coefficient η , and the heat conductivity λ of the gas (indeed, these coefficients are related to the Navier–Stokes limit of the Boltzmann

equation). Namely, they are given by

$$\begin{cases} R_1 = R_2 = -\frac{\lambda}{d+2} - \frac{\eta}{2}, \\ R_3 = \dots = R_{d+1} = -\frac{\eta d}{2(d-1)}, \\ R_{d+2} = -\frac{\lambda d}{d+2}. \end{cases}$$

Therefore for $|k_0|/L \ll 1$, the damping rate is given by the minimum among these values.

To conclude these tests, we performed a final numerical experiment to evaluate the robustness of the theory of the trend to equilibrium. We have chosen an initial datum which is far from the equilibrium

(6.4)

$$f_0(x, v) = \frac{1}{2\pi v_{th}^2} (1 + A_0 \cos(k_0 x)) \left[\exp\left(-\frac{|v - v_0|^2}{2v_{th}^2}\right) + \exp\left(-\frac{|v + v_0|^2}{2v_{th}^2}\right) \right],$$

with $v_0 = (1/2, 1/2)$ and $v_{th} = \sqrt{3}/2$. We present the time evolution of the relative entropies H_l and H_g in log scale and observe that initially the entropy is strongly decreasing, and when the distribution function becomes close to a local equilibrium, some oscillations appear with the good frequency $\omega/L = 1/\sqrt{2}$ and damping rate α (see Figure 9).

Remarks.

1. Now numerical methods for the Boltzmann equation, such as the one presented in this paper, become able to provide very accurate simulations of the transient regime towards equilibrium with reasonable cost, even in the inhomogeneous case. This could be used to explore *numerically* the spectrum of the linearized Boltzmann collision operator (in the homogeneous case), or, more interestingly, the spectrum of the linearized Boltzmann collision operator together with the transport term in the inhomogeneous case. For instance, the exponential rate of convergence is directly readable on the figures above and provides a numerical estimation of the *spectral gap* (that is, the real part of the first nonzero eigenvalue) of this operator (it has been known since Ukai [48] that this operator has a spectral gap in the torus; see also [13]). Moreover, by a frequency analysis of the curve of the time evolution of the relative entropy or the L^1 distance to the equilibrium, it could be possible also to describe other eigenvalues; as long as they have a different imaginary part, it should be possible (in principle) to extract from the frequency analysis the curve corresponding to their contribution in the evolution semigroup, and then to compute their real part which corresponds to the exponential rate of decay of this curve.

2. A recent work [27] gave a detailed pointwise study of the Green function for the linearized Boltzmann equation in the domain $x \in \Omega = \mathbb{R}$. This particular study shows that the long-time behavior is governed by “fluid-like waves” (corresponding to the waves of the linearized Euler and Navier–Stokes equations) whose amplitude decreases polynomially, whereas the amplitude of the “kinetic part” of the Green function decreases exponentially. We think it is likely that this study could be extended to the torus, where the amplitude of both the fluid and the kinetic parts of the Green function should decrease exponentially. Moreover, the rate of decay of the kinetic part should not depend on the size of the box, whereas the rate of decay of the

fluid part should. Hence for a box small enough, the long-time behavior should be governed by the kinetic part of the Green function (that is, like the spatially homogeneous Boltzmann equation), whereas for a box big enough, the long-time behavior should be governed by the fluid-like waves. This is precisely what we observe numerically, and thus this theoretical study could provide a rigorous proof of the numerical observations above, at least in the linearized regime.

7. Conclusions. In this paper we have introduced and deeply tested a class of new fast algorithms for the computation of the Boltzmann collision operator. These methods allow us to reduce the computational cost from $O(n^2)$ to $O(n \log_2 n)$. We give computational evidence of the great performance of the schemes which can provide a dramatic speed-up in computing time of deterministic schemes by making them competitive with Monte Carlo methods, where higher accuracy is required. A first numerical application to a nontrivial problem in the space nonhomogeneous case confirms the strong computing potential of the new schemes.

Other methods such as singular value decomposition, fast multipole methods [9], separated representations in high-dimensional problems (see, for instance, [1]), wavelets, etc. could have been used to search for a decomposition of the form (4.1). However, to our knowledge it is not known at present how to obtain the properties described above for our decomposition with these methods.

Acknowledgment. We would like to thank Cédric Villani for suggesting the numerical study of possible oscillations in the relaxation to equilibrium.

REFERENCES

- [1] B. ALPERT, G. BEYLKIN, R. COIFMAN, AND V. ROKHLIN, *Wavelet-like bases for the fast solution of second-kind integral equations*, SIAM J. Sci. Comput., 14 (1993), pp. 159–184.
- [2] H. BABOVSKY, *On a simulation scheme for the Boltzmann equation*, Math. Methods Appl. Sci., 8 (1986), pp. 223–233.
- [3] G. A. BIRD, *Molecular Gas Dynamics*, Clarendon Press, Oxford, UK, 1994.
- [4] A. V. BOBYLEV, *Exact solutions of the Boltzmann equation*, Dokl. Akad. Nauk. SSSR, 225 (1975), pp. 1296–1299 (in Russian).
- [5] A. V. BOBYLEV, *The theory of the nonlinear spatially uniform Boltzmann equation for Maxwell molecules*, Math. Phys. Reviews, 7 (1988), pp. 111–233.
- [6] A. V. BOBYLEV, A. PALCZEWSKI, AND J. SCHNEIDER, *On approximation of the Boltzmann equation by discrete velocity models*, C. R. Acad. Sci. Paris Sér. I Math., 320 (1995), pp. 639–644.
- [7] A. V. BOBYLEV AND S. RJASANOW, *Difference scheme for the Boltzmann equation based on the fast Fourier transform*, Eur. J. Mech. B Fluids, 16 (1997), pp. 293–306.
- [8] A. V. BOBYLEV AND S. RJASANOW, *Fast deterministic method of solving the Boltzmann equation for hard spheres*, Eur. J. Mech. B Fluids, 18 (1999), pp. 869–887.
- [9] O. BOKANOWSKI AND M. LEMOU, *Fast multipole method for multivariable integrals*, SIAM J. Numer. Anal., 42 (2005), pp. 2098–2117.
- [10] C. BUET, *A discrete velocity scheme for the Boltzmann operator of rarefied gas dynamics*, Transp. Theory Statist. Phys., 25 (1996), pp. 33–60.
- [11] C. CANUTO, M. Y. HUSSAINI, A. QUARTERONI, AND T. A. ZANG, *Spectral Methods in Fluid Dynamics*, Springer Ser. Comput. Phys., Springer-Verlag, New York, 1988.
- [12] T. CARLEMAN, *Sur la théorie de l'équation intégrodifférentielle de Boltzmann*, Acta Math., 60 (1933), pp. 91–146.
- [13] C. CERCIGNANI, R. ILLNER, AND M. PULVIRENTI, *The Mathematical Theory of Dilute Gases*, Appl. Math. Sci. 106, Springer-Verlag, New York, 1994.
- [14] A. J. CHORIN, *Numerical solution of Boltzmann's equation*, Comm. Pure Appl. Math., 25 (1972), pp. 171–186.
- [15] P. DEGOND, L. PARESCHI, AND G. RUSSO, EDS., *Modeling and Computational Methods for Kinetic Equations*, Model. Simul. Sci. Eng. Technol. Birkhäuser Boston, Boston, MA, 2004.

- [16] L. DESVILLETES AND C. VILLANI, *On the trend to global equilibrium for spatially inhomogeneous kinetic systems: The Boltzmann equation*, Invent. Math., 159 (2005), pp. 245–316.
- [17] R. S. ELLIS AND M. A. PINSKY, *The first and second fluid approximations to the linearized Boltzmann equation*, J. Math. Pures Appl., 54 (1975), pp. 125–156.
- [18] F. FILBET, E. SONNENDRÜCKER, AND P. BERTRAND, *Conservative numerical schemes for the Vlasov equation*, J. Comput. Phys., 172 (2001), pp. 166–187.
- [19] F. FILBET AND L. PARESCHI, *A numerical method for the accurate solution of the Fokker-Planck-Landau equation in the nonhomogeneous case*, J. Comput. Phys., 179 (2002), pp. 1–26.
- [20] F. FILBET AND G. RUSSO, *High order numerical methods for the space non-homogeneous Boltzmann equation*, J. Comput. Phys., 186 (2003), pp. 457–480.
- [21] F. FILBET, L. PARESCHI, AND G. TOSCANI, *Accurate numerical methods for the collisional motion of (heated) granular flows*, J. Comput. Phys., 202 (2005), pp. 216–235.
- [22] F. FILBET AND G. RUSSO, *Accurate numerical methods for the Boltzmann equation*, in Modeling and Computational Methods for Kinetic Equations, Model Sumil. Sci Eng. Technol., Birkhäuser Boston, Boston, MA, 2004, pp. 117–145.
- [23] F. FILBET AND G. RUSSO, *A rescaling velocity method for kinetic equations: The homogeneous case*, in Proceedings of the Workshop on Modelling and Numerics of Kinetic Dissipative Systems (Lipari, Italy), L. Pareschi, G. Russo, and G. Toscani, eds., Nova Science Publishers, Inc., Hauppauge, NY, 2006.
- [24] E. GABETTA, L. PARESCHI, AND G. TOSCANI, *Relaxation schemes for nonlinear kinetic equations*, SIAM J. Numer. Anal., 34 (1997), pp. 2168–2194.
- [25] Y. GUO, *The Vlasov-Maxwell-Boltzmann system near Maxwellians*, Invent. Math., 153 (2003), pp. 593–630.
- [26] I. IBRAGIMOV AND S. RJASANOW, *Numerical solution of the Boltzmann equation on the uniform grid*, Computing, 69 (2002), pp. 163–186.
- [27] T.-P. LIU AND S.-H. YU, *The Green's function and large-time behavior of solutions for the one-dimensional Boltzmann equation*, Comm. Pure Appl. Math., 57 (2004), pp. 1543–1608.
- [28] P. A. MARKOWICH AND L. PARESCHI, *Fast conservative and entropic methods for the Boson Boltzmann equation*, Numer. Math., 99 (2005), pp. 509–532.
- [29] Y.-L. MARTIN, F. ROGIER, AND J. SCHNEIDER, *Une méthode déterministe pour la résolution de l'équation de Boltzmann inhomogène*, C. R. Acad. Sci. Paris Sér. I Math., 314 (1992), pp. 483–487.
- [30] S. MISCHLER AND C. MOUHOT, *Cooling process for inelastic Boltzmann equations for hard spheres, Part II: Self-similar solutions and tail behavior*, J. Statist. Phys., to appear.
- [31] C. MOUHOT, *Quantitative lower bounds for the full Boltzmann equation, Part I: Periodic boundary conditions*, Comm. Partial Differential Equations, 30 (2005), pp. 881–917.
- [32] C. MOUHOT AND L. PARESCHI, *Fast methods for the Boltzmann collision operator*, C. R. Math. Acad. Sci. Paris, 339 (2004), pp. 71–76.
- [33] C. MOUHOT AND L. PARESCHI, *Fast algorithms for computing the Boltzmann collision operator*, Math. Comp., to appear.
- [34] C. MOUHOT AND L. PARESCHI, *An $O(N \log_2 N)$ Algorithm for Computing Discrete Velocity Models*, in preparation.
- [35] G. NALDI, L. PARESCHI, AND G. TOSCANI, *Spectral methods for one-dimensional kinetic models of granular flows and numerical quasi elastic limit*, M2AN Math. Model. Numer. Anal., 37 (2003), pp. 73–90.
- [36] K. NANBU, *Direct simulation scheme derived from the Boltzmann equation. I. Monocomponent Gases*, J. Phys. Soc. Japan, 52 (1983), pp. 2042–2049.
- [37] A. PALCZEWSKI, J. SCHNEIDER, AND A. V. BOBYLEV, *A consistency result for a discrete-velocity model of the Boltzmann equation*, SIAM J. Numer. Anal., 34 (1997), pp. 1865–1883.
- [38] A. PALCZEWSKI AND J. SCHNEIDER, *Existence, stability, and convergence of solutions of discrete velocity models to the Boltzmann equation*, J. Statist. Phys., 91 (1998), pp. 307–326.
- [39] V. A. PANFEROV AND A. G. HEINTZ, *A new consistent discrete-velocity model for the Boltzmann equation*, Math. Methods Appl. Sci., 25 (2002), pp. 571–593.
- [40] L. PARESCHI AND B. PERTHAME, *A Fourier spectral method for homogeneous Boltzmann equations*, Transp. Theory Statist. Phys., 25 (1996), pp. 369–382.
- [41] L. PARESCHI AND G. RUSSO, *Numerical solution of the Boltzmann equation I. Spectrally accurate approximation of the collision operator*, SIAM J. Numer. Anal., 37 (2000), pp. 1217–1245.
- [42] L. PARESCHI, G. RUSSO, AND G. TOSCANI, *Fast spectral methods for the Fokker-Planck-Landau collision operator*, J. Comput. Phys., 165 (2000), pp. 216–236.

- [43] L. PARESCHI AND G. RUSSO, *On the stability of spectral methods for the homogeneous Boltzmann equation*, *Transp. Theory Statist. Phys.*, 29 (2000), pp. 431–447.
- [44] A. PULVIRENTI AND B. WENNERBERG, *A Maxwellian lower bound for solutions to the Boltzmann equation*, *Comm. Math. Phys.*, 183 (1997), pp. 145–160.
- [45] F. ROGIER AND J. SCHNEIDER, *A direct method for solving the Boltzmann equation*, *Transp. Theory Statist. Phys.*, 23 (1994), pp. 313–338.
- [46] G. A. SOD, *A numerical solution of Boltzmann's equation*, *Comm. Pure Appl. Math.*, 30 (1977), pp. 391–419.
- [47] R. STRAIN AND Y. GUO, *Almost exponential decay near Maxwellian*, *Comm. Partial Differential Equations*, 31 (2006), pp. 417–429.
- [48] S. UKAI, *On the existence of global solutions of mixed problem for non-linear Boltzmann equation*, *Proc. Japan Acad.*, 50 (1974), pp. 179–184.
- [49] C. VILLANI, *A review of mathematical topics in collisional kinetic theory*, in *Handbook of Mathematical Fluid Mechanics*, Vol. 1, S. Friedlander and D. Serre, eds., North-Holland, Amsterdam, 2002, pp. 91–309.