

Torus actions in the normalization problem

Jasmin Raissy

Dipartimento di Matematica "L. Tonelli"
Università di Pisa

KAWA 1, Workshop
Albi, January 29–31, 2010

Normalization Problem

Setting

Given $f: (\mathbb{C}^n, p) \rightarrow (\mathbb{C}^n, p)$ a germ of biholomorphism fixing p , we are interested in the dynamics of f in a neighbourhood of p .

Normalization Problem

Setting

Given $f: (\mathbb{C}^n, p) \rightarrow (\mathbb{C}^n, p)$ a germ of biholomorphism fixing p , we are interested in the dynamics of f in a neighbourhood of p .

i.e., for any q “sufficiently close” to p , we want to study the asymptotical behavior of $\{f^k(q)\}_{k \geq 1}$, where $f^k = f \circ \dots \circ f$.

Normalization Problem

Setting

Let $f: (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$ be germ of biholomorphism fixing O .

Normalization Problem

Setting

Let $f: (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$ be germ of biholomorphism fixing O .

Locally, using multi-index notation,

$$f(z) = \Lambda z + \sum_{\substack{Q \in \mathbb{N}^n \\ |Q| \geq 2}} f_Q z^Q,$$

where

$$z^Q := z_1^{q_1} \cdots z_n^{q_n}, \quad f_Q \in \mathbb{C}^n, \quad |Q| := \sum_{j=1}^n q_j,$$

Normalization Problem

Setting

Let $f: (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$ be germ of biholomorphism fixing O .

Locally, using multi-index notation,

$$f(z) = \Lambda z + \sum_{\substack{Q \in \mathbb{N}^n \\ |Q| \geq 2}} f_Q z^Q,$$

where

$$z^Q := z_1^{q_1} \cdots z_n^{q_n}, \quad f_Q \in \mathbb{C}^n, \quad |Q| := \sum_{j=1}^n q_j,$$

with Λ in Jordan normal form, i.e.,

$$\Lambda = \text{Diag}(\lambda_1, \dots, \lambda_n) + N$$

N = nilpotent matrix and $\lambda_1, \dots, \lambda_n \in \mathbb{C}^*$ not necessarily distinct.

Normalization Problem

Setting

Let $f: (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$ be germ of biholomorphism fixing O .

Locally, using multi-index notation,

$$f(z) = \Lambda z + \sum_{\substack{Q \in \mathbb{N}^n \\ |Q| \geq 2}} f_Q z^Q,$$

where

$$z^Q := z_1^{q_1} \cdots z_n^{q_n}, \quad f_Q \in \mathbb{C}^n, \quad |Q| := \sum_{j=1}^n q_j,$$

with Λ in Jordan normal form, i.e.,

$$\Lambda = \text{Diag}(\lambda_1, \dots, \lambda_n) + N$$

N = nilpotent matrix and $\lambda_1, \dots, \lambda_n \in \mathbb{C}^*$ not necessarily distinct.

Want to know whether $\exists \varphi: (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$, local holomorphic change of coordinates, $d\varphi_O = \text{Id}$, s.t. $\varphi^{-1} \circ f \circ \varphi$ has a **simple** form.

Normalization Problem

Linearization

Linearization problem

simple = linear

Normalization Problem

Linearization

Linearization problem

simple = **linear**

Idea: first to search for a formal solution of

$$f \circ \varphi = \varphi \circ \Lambda \tag{1}$$

and then to study the convergence of φ .

Normalization Problem

Linearization

Linearization problem

simple = linear


Idea: first to search for a formal solution of

$$f \circ \varphi = \varphi \circ \Lambda \quad (1)$$

and then to study the convergence of φ . We have to recursively solve, for each coordinate j ,

- $\varphi_j(z) = z_j + \sum_{|Q| \geq 2} \varphi_{Q,j} z^Q.$

- $\lambda^Q := \lambda_1^{q_1} \cdots \lambda_n^{q_n}$


$$(\lambda^Q - \lambda_j) \varphi_{Q,j} = \text{Polynomial}(f_{P,j}, \varphi_{R,k}, \text{ with } P \leq Q, R < Q) \quad (2)$$

Normalization Problem

Linearization

Linearization problem


simple = **linear**

Idea: first to search for a formal solution of

$$f \circ \varphi = \varphi \circ \Lambda \quad (1)$$

and then to study the convergence of φ . We have to recursively solve, for each coordinate j ,

- $\varphi_j(\mathbf{z}) = z_j + \sum_{|Q| \geq 2} \varphi_{Q,j} z^Q.$
- $\lambda^Q := \lambda_1^{q_1} \cdots \lambda_n^{q_n}$


$$(\lambda^Q - \lambda_j) \varphi_{Q,j} = \text{Polynomial}(f_{P,j}, \varphi_{R,k}, \text{ with } P \leq Q, R < Q) \quad (2)$$

- lexicographic order on \mathbb{N}^n
- 

Normalization Problem

Resonances

Definition

A **resonant multi-index** for $\lambda \in (\mathbb{C}^*)^n$, rel. to $j \in \{1, \dots, n\}$ is $Q \in \mathbb{N}^n$, with $|Q| \geq 2$, s.t.

$$\lambda^Q - \lambda_j = 0. \quad (3)$$

$$\text{Res}_j(\lambda) := \{Q \in \mathbb{N}^n \mid |Q| \geq 2, \lambda^Q - \lambda_j = 0\}.$$

Normalization Problem

Resonances

Definition

A **resonant multi-index** for $\lambda \in (\mathbb{C}^*)^n$, rel. to $j \in \{1, \dots, n\}$ is $Q \in \mathbb{N}^n$, with $|Q| \geq 2$, s.t.

$$\lambda^Q - \lambda_j = 0. \quad (3)$$

$\text{Res}_j(\lambda) := \{Q \in \mathbb{N}^n \mid |Q| \geq 2, \lambda^Q - \lambda_j = 0\}.$

Resonances = **obstruction** to *formal* linearization.

Normalization Problem

Poincaré-Dulac normal forms

Theorem (Poincaré-Dulac, 1904)

$\forall f$ as above $\exists \hat{\varphi}$ **formal** change of coord., $d\hat{\varphi}_0 = \text{Id}$, s.t.
 $\hat{\varphi}^{-1} \circ f \circ \hat{\varphi} = g \in \mathbb{C}[[z_1, \dots, z_n]]^n$ where $g(O) = O$, $dg_O = df_O$ and g has only resonant monomials,

$$g_j(z) = \lambda_j z_j + \varepsilon_j z_{j+1} + \sum_{\substack{|Q| \geq 2 \\ \lambda^Q = \lambda_j}} g_{Q,j} z^Q.$$

Moreover, the resonant terms of $\hat{\varphi}$ can be arbitrarily chosen, and that choice determines uniquely g^{res} and the remaining terms of $\hat{\varphi}$.

A germ of the form $\Lambda + g^{\text{res}}$, with g^{res} containing only resonant monomials is said in **Poincaré-Dulac normal form**.

Normalization Problem

Normalization Problem

Given $f, \exists \varphi: (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$, **holomorphic** change of coordinates, $d\varphi_O = \text{Id}$, s.t.

$\varphi^{-1} \circ f \circ \varphi$ is in Poincaré-Dulac normal form?

Normalization Problem

Normalization Problem

Given $f, \exists \varphi: (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$, **holomorphic** change of coordinates, $d\varphi_O = \text{Id}$, s.t.

$\varphi^{-1} \circ f \circ \varphi$ is in Poincaré-Dulac normal form?

Problem

Not uniqueness of the formal change of coordinates $\hat{\varphi}$ given by Poincaré-Dulac theorem, and not having explicit expression for g^{res} , make very difficult to give estimates for the convergence of $\hat{\varphi}$.

Torus Actions

Torus Actions

The same problem can be stated for germs of holomorphic vector field near a singular point.

Torus Actions

The same problem can be stated for germs of holomorphic vector field near a singular point.

In 2002, N.T. Zung found that to find a Poincaré-Dulac holomorphic normalization for a germ of holomorphic vector field is the same as to find (and linearize) a suitable torus action which preserves the vector field.

Torus Actions

The same problem can be stated for germs of holomorphic vector field near a singular point.

In 2002, N.T. Zung found that to find a Poincaré-Dulac holomorphic normalization for a germ of holomorphic vector field is the same as to find (and linearize) a suitable torus action which preserves the vector field.

Idea

We look for symmetries in the normalization problem and how to exploit them

Torus Actions

Germes commuting with a torus action

Theorem (–, 2009)

f commutes with a holom. effective \mathbb{T}^r -action on $(\mathbb{C}^n, 0)$, $1 \leq r \leq n$, with weight matrix $\Theta \in M_{n \times r}(\mathbb{Z})$



$\exists \varphi$ local holom. change of coord. s.t. $\varphi^{-1} \circ f \circ \varphi$ contains only Θ -resonant monomials.

Torus Actions

Germes commuting with a torus action

Theorem (–, 2009)

f commutes with a holom. effective \mathbb{T}^r -action on (\mathbb{C}^n, O) , $1 \leq r \leq n$, with weight matrix $\Theta \in M_{n \times r}(\mathbb{Z})$



$\exists \varphi$ local holom. change of coord. s.t. $\varphi^{-1} \circ f \circ \varphi$ contains only Θ -resonant monomials.

f **commutes with** $A: \mathbb{T}^r \times (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$, $A(x, O) = O$, means
$$f(A(x, z)) = A(x, f(z)).$$

Torus Actions

Germes commuting with a torus action

Theorem (–, 2009)

f commutes with a holom. effective \mathbb{T}^r -action on (\mathbb{C}^n, O) , $1 \leq r \leq n$, with weight matrix $\Theta \in M_{n \times r}(\mathbb{Z})$



$\exists \varphi$ local holom. change of coord. s.t. $\varphi^{-1} \circ f \circ \varphi$ contains only Θ -resonant monomials.

f **commutes with** $A: \mathbb{T}^r \times (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$, $A(x, O) = O$, means

$$f(A(x, z)) = A(x, f(z)).$$

A^{lin} is semi-simple and $\text{Sp}(A^{\text{lin}}(x, \cdot)) = \{\exp(2\pi i \sum_{k=1}^r x_k \theta_j^k)\}_{j=1, \dots, n}$

where $\Theta = (\theta_j^k) \in M_{n \times r}(\mathbb{Z})$ is **the weight matrix of A** .

Torus Actions

Germes commuting with a torus action

Theorem (–, 2009)

f commutes with a holom. effective \mathbb{T}^r -action on (\mathbb{C}^n, O) , $1 \leq r \leq n$, with weight matrix $\Theta \in M_{n \times r}(\mathbb{Z})$



$\exists \varphi$ local holom. change of coord. s.t. $\varphi^{-1} \circ f \circ \varphi$ contains only Θ -resonant monomials.

f **commutes with** $A: \mathbb{T}^r \times (\mathbb{C}^n, O) \rightarrow (\mathbb{C}^n, O)$, $A(x, O) = O$, means

$$f(A(x, z)) = A(x, f(z)).$$

A^{lin} is semi-simple and $\text{Sp}(A^{\text{lin}}(x, \cdot)) = \{\exp(2\pi i \sum_{k=1}^r x_k \theta_j^k)\}_{j=1, \dots, n}$

where $\Theta = (\theta_j^k) \in M_{n \times r}(\mathbb{Z})$ is **the weight matrix of A** .

$z^Q e_j$, with $|Q| \geq 1$, is **Θ -resonant** if

$$\langle Q, \theta^k \rangle := \sum_{h=1}^n q_h \theta_h^k = \theta_j^k \quad \forall k = 1, \dots, r.$$

Torus Actions

Definition

An **additive resonant multi-index** for $\theta \in \mathbb{C}^n$, rel. to $j \in \{1, \dots, n\}$ is $Q \in \mathbb{N}^n$, with $|Q| \geq 2$, s.t.

$$\langle Q, \theta \rangle = \theta_j. \quad (4)$$

$$\text{Res}_j^+(\theta) := \{Q \in \mathbb{N}^n \mid |Q| \geq 2, \langle Q, \theta \rangle = \theta_j\}.$$

Torus Actions

Definition

An **additive resonant multi-index** for $\theta \in \mathbb{C}^n$, rel. to $j \in \{1, \dots, n\}$ is $Q \in \mathbb{N}^n$, with $|Q| \geq 2$, s.t.

$$\langle Q, \theta \rangle = \theta_j. \quad (4)$$

$$\text{Res}_j^+(\theta) := \{Q \in \mathbb{N}^n \mid |Q| \geq 2, \langle Q, \theta \rangle = \theta_j\}.$$

$$\{Q \in \mathbb{N}^n \mid |Q| \geq 2, Q \text{ } \Theta\text{-resonant rel. to } j\} = \bigcap_{k=1}^r \text{Res}_j^+(\theta^k)$$

Torus Actions

Definition

An **additive resonant multi-index** for $\theta \in \mathbb{C}^n$, rel. to $j \in \{1, \dots, n\}$ is $Q \in \mathbb{N}^n$, with $|Q| \geq 2$, s.t.

$$\langle Q, \theta \rangle = \theta_j. \quad (4)$$

$$\text{Res}_j^+(\theta) := \{Q \in \mathbb{N}^n \mid |Q| \geq 2, \langle Q, \theta \rangle = \theta_j\}.$$

$$\{Q \in \mathbb{N}^n \mid |Q| \geq 2, Q \text{ } \Theta\text{-resonant rel. to } j\} = \bigcap_{k=1}^r \text{Res}_j^+(\theta^k)$$

Corollary (–, 2009)

f is holomorphically linearizable



it commutes with a \mathbb{T}^r -action, $1 \leq r \leq n$, with Θ having no resonances of degree $|Q| \geq 2$.

Torus Actions

Holomorphic Normalization

Torus Actions

$$\Theta$$

Holomorphic Normalization

$$\lambda_1, \dots, \lambda_n$$

Strategy

Torus Actions

$$\Theta$$

$$\bigcap_{k=1}^r \operatorname{Res}_j^+(\theta^k)$$

Holomorphic Normalization

$$\lambda_1, \dots, \lambda_n$$

$$\operatorname{Res}_j(\lambda)$$

Toric Degree

Definition

The **toric degree** of $\lambda \in (\mathbb{C}^*)^n$ is the min $r \in \mathbb{N}$ s.t. $\exists \alpha_1, \dots, \alpha_r \in \mathbb{C}^*$ and $\exists \theta^{(1)}, \dots, \theta^{(r)} \in \mathbb{Z}^n$ s.t.

$$[\varphi] = \left[\sum_{k=1}^r \alpha_k \theta^{(k)} \right] \in (\mathbb{C}/\mathbb{Z})^n,$$

where $[\varphi]$ is the unique in $(\mathbb{C}/\mathbb{Z})^n$ s.t. $\lambda = e^{2\pi i[\varphi]}$.

$\theta^{(1)}, \dots, \theta^{(r)}$ are a **r -tuple of toric vectors associated to λ** , with **toric coefficients** $\alpha_1, \dots, \alpha_r$.

Toric Degree

Definition

The **toric degree** of $\lambda \in (\mathbb{C}^*)^n$ is the min $r \in \mathbb{N}$ s.t. $\exists \alpha_1, \dots, \alpha_r \in \mathbb{C}^*$ and $\exists \theta^{(1)}, \dots, \theta^{(r)} \in \mathbb{Z}^n$ s.t.

$$[\varphi] = \left[\sum_{k=1}^r \alpha_k \theta^{(k)} \right] \in (\mathbb{C}/\mathbb{Z})^n,$$

where $[\varphi]$ is the unique in $(\mathbb{C}/\mathbb{Z})^n$ s.t. $\lambda = e^{2\pi i[\varphi]}$.

$\theta^{(1)}, \dots, \theta^{(r)}$ are a **r -tuple of toric vectors associated to λ** , with **toric coefficients** $\alpha_1, \dots, \alpha_r$.

Since $[\varphi] = [\sum_{j=1}^n \varphi_j e_j]$, the toric degree is well-defined and $1 \leq r \leq n$.

Toric Degree

Definition

The **toric degree** of $\lambda \in (\mathbb{C}^*)^n$ is the min $r \in \mathbb{N}$ s.t. $\exists \alpha_1, \dots, \alpha_r \in \mathbb{C}^*$ and $\exists \theta^{(1)}, \dots, \theta^{(r)} \in \mathbb{Z}^n$ s.t.

$$[\varphi] = \left[\sum_{k=1}^r \alpha_k \theta^{(k)} \right] \in (\mathbb{C}/\mathbb{Z})^n,$$

where $[\varphi]$ is the unique in $(\mathbb{C}/\mathbb{Z})^n$ s.t. $\lambda = e^{2\pi i[\varphi]}$.

$\theta^{(1)}, \dots, \theta^{(r)}$ are a **r -tuple of toric vectors associated to λ** , with **toric coefficients** $\alpha_1, \dots, \alpha_r$.

Since $[\varphi] = [\sum_{j=1}^n \varphi_j e_j]$, the toric degree is well-defined and $1 \leq r \leq n$.

- $\alpha_1, \dots, \alpha_r$ are \mathbb{Z} -independent,

Toric Degree

Definition

The **toric degree** of $\lambda \in (\mathbb{C}^*)^n$ is the min $r \in \mathbb{N}$ s.t. $\exists \alpha_1, \dots, \alpha_r \in \mathbb{C}^*$ and $\exists \theta^{(1)}, \dots, \theta^{(r)} \in \mathbb{Z}^n$ s.t.

$$[\varphi] = \left[\sum_{k=1}^r \alpha_k \theta^{(k)} \right] \in (\mathbb{C}/\mathbb{Z})^n,$$

where $[\varphi]$ is the unique in $(\mathbb{C}/\mathbb{Z})^n$ s.t. $\lambda = e^{2\pi i[\varphi]}$.

$\theta^{(1)}, \dots, \theta^{(r)}$ are a **r -tuple of toric vectors associated to λ** , with **toric coefficients** $\alpha_1, \dots, \alpha_r$.

Since $[\varphi] = [\sum_{j=1}^n \varphi_j e_j]$, the toric degree is well-defined and $1 \leq r \leq n$.

- $\alpha_1, \dots, \alpha_r$ are \mathbb{Z} -independent,
- $\theta^{(1)}, \dots, \theta^{(r)}$ are \mathbb{Q} -linearly independent.

Main Result

Theorem (–, 2009)

Take f as above. Then, *in all but one case*, f is holomorphically normalizable



f commutes with a holom. effective torus action on (\mathbb{C}^n, O) of dim *depending on* $\text{tordeg}(\lambda)$ with columns of Θ *related to toric vectors associated to* λ .

First distinction

Proposition (–, 2009)

\exists toric r -tuple assoc. to λ with coeff. \mathbb{Z} -independent with 1



\forall toric r -tuple assoc. to λ the coeff. are \mathbb{Z} -independent with 1.

First distinction

Proposition (–, 2009)

\exists toric r -tuple assoc. to λ with coeff. \mathbb{Z} -independent with 1



\forall toric r -tuple assoc. to λ the coeff. are \mathbb{Z} -independent with 1.

- **Torsion-free case:** $1, \alpha_1, \dots, \alpha_r$ \mathbb{Z} -independent

First distinction

Proposition (–, 2009)

\exists toric r -tuple assoc. to λ with coeff. \mathbb{Z} -independent with 1



\forall toric r -tuple assoc. to λ the coeff. are \mathbb{Z} -independent with 1.

- **Torsion-free case:** $1, \alpha_1, \dots, \alpha_r$ \mathbb{Z} -independent
- **Torsion case:** $1, \alpha_1, \dots, \alpha_r$ \mathbb{Z} -dependent

First distinction

Proposition (–, 2009)

\exists toric r -tuple assoc. to λ with coeff. \mathbb{Z} -independent with 1



\forall toric r -tuple assoc. to λ the coeff. are \mathbb{Z} -independent with 1.

- **Torsion-free case:** $1, \alpha_1, \dots, \alpha_r$ \mathbb{Z} -independent
- **Torsion case:** $1, \alpha_1, \dots, \alpha_r$ \mathbb{Z} -dependent

In the torsion case we can always consider **reduced** toric r -tuples, i.e., $\eta^{(1)}, \dots, \eta^{(r)}$ with coeff. β_1, \dots, β_r s.t. $\beta_1 = 1/m$ with $m \in \mathbb{N} \setminus \{0, 1\}$ and $m, \eta_1^{(1)}, \dots, \eta_n^{(1)}$ coprime; $\eta^{(2)}, \dots, \eta^{(r)}$ are called **reduced torsion-free toric vectors** assoc. to λ .

Torsion-free Case

Main Theorem in the torsion-free case

\forall r -tuple of toric vectors $\theta^{(1)}, \dots, \theta^{(r)}$ associated to λ

$$\mathrm{Res}_j(\lambda) = \bigcap_{k=1}^r \mathrm{Res}_j^+(\theta^{(k)}) \quad \forall j = 1, \dots, n$$

Torsion-free Case

Main Theorem in the torsion-free case

\forall r -tuple of toric vectors $\theta^{(1)}, \dots, \theta^{(r)}$ associated to λ

$$\mathrm{Res}_j(\lambda) = \bigcap_{k=1}^r \mathrm{Res}_j^+(\theta^{(k)}) \quad \forall j = 1, \dots, n$$

and

$$\lambda_j = \lambda_h \implies \theta_j^{(k)} = \theta_h^{(k)} \quad \forall k = 1, \dots, r.$$

Torsion-free Case

Main Theorem in the torsion-free case

\forall r -tuple of toric vectors $\theta^{(1)}, \dots, \theta^{(r)}$ associated to λ

$$\text{Res}_j(\lambda) = \bigcap_{k=1}^r \text{Res}_j^+(\theta^{(k)}) \quad \forall j = 1, \dots, n$$

and

$$\lambda_j = \lambda_h \implies \theta_j^{(k)} = \theta_h^{(k)} \quad \forall k = 1, \dots, r.$$

Main Theorem

f *in the torsion-free case* is holomorphically normalizable



f commutes with a holom. effective \mathbb{T}^r -action on $(\mathbb{C}^n, 0)$, $r = \text{tordeg}(\lambda)$,
with columns of Θ that are a *r -tuple of toric vectors associated to λ* .

Torsion Case

\forall reduced toric r -tuple $\eta^{(1)}, \dots, \eta^{(r)}$ associated to λ

$$\bigcap_{k=2}^r \operatorname{Res}_j^+(\eta^{(k)}) \supseteq \operatorname{Res}_j(\lambda) \supseteq \bigcap_{k=1}^r \operatorname{Res}_j^+(\eta^{(k)}). \quad (5)$$

Torsion Case

\forall reduced toric r -tuple $\eta^{(1)}, \dots, \eta^{(r)}$ associated to λ

$$\bigcap_{k=2}^r \text{Res}_j^+(\eta^{(k)}) \supseteq \text{Res}_j(\lambda) \supseteq \bigcap_{k=1}^r \text{Res}_j^+(\eta^{(k)}). \quad (5)$$

We have the following sub-cases:

- **Impure torsion case:** for a reduced toric r -tuple ($\Rightarrow \forall$)

$$\text{Res}_j(\lambda) = \bigcap_{k=2}^r \text{Res}_j^+(\eta^{(k)}) \quad \forall j = 1, \dots, n$$

Torsion Case

\forall reduced toric r -tuple $\eta^{(1)}, \dots, \eta^{(r)}$ associated to λ

$$\bigcap_{k=2}^r \text{Res}_j^+(\eta^{(k)}) \supseteq \text{Res}_j(\lambda) \supseteq \bigcap_{k=1}^r \text{Res}_j^+(\eta^{(k)}). \quad (5)$$

We have the following sub-cases:

- **Impure torsion case:** for a reduced toric r -tuple ($\Rightarrow \forall$)

$$\text{Res}_j(\lambda) = \bigcap_{k=2}^r \text{Res}_j^+(\eta^{(k)}) \quad \forall j = 1, \dots, n$$

- **Pure torsion case:** the first inclusion is always strict, and

Torsion Case

\forall reduced toric r -tuple $\eta^{(1)}, \dots, \eta^{(r)}$ associated to λ

$$\bigcap_{k=2}^r \text{Res}_j^+(\eta^{(k)}) \supseteq \text{Res}_j(\lambda) \supseteq \bigcap_{k=1}^r \text{Res}_j^+(\eta^{(k)}). \quad (5)$$

We have the following sub-cases:

- **Impure torsion case:** for a reduced toric r -tuple ($\Rightarrow \forall$)

$$\text{Res}_j(\lambda) = \bigcap_{k=2}^r \text{Res}_j^+(\eta^{(k)}) \quad \forall j = 1, \dots, n$$

- **Pure torsion case:** the first inclusion is always strict, and *either*
 - ▶ **λ can be simplified:** \exists a reduced toric r -tuple, said **simple**, s.t.
 $\text{Res}_j(\lambda) = \bigcap_{k=1}^r \text{Res}_j^+(\eta^{(k)}) \quad \forall j,$

Torsion Case

\forall reduced toric r -tuple $\eta^{(1)}, \dots, \eta^{(r)}$ associated to λ

$$\bigcap_{k=2}^r \text{Res}_j^+(\eta^{(k)}) \supseteq \text{Res}_j(\lambda) \supseteq \bigcap_{k=1}^r \text{Res}_j^+(\eta^{(k)}). \quad (5)$$

We have the following sub-cases:

- **Impure torsion case:** for a reduced toric r -tuple ($\Rightarrow \forall$)

$$\text{Res}_j(\lambda) = \bigcap_{k=2}^r \text{Res}_j^+(\eta^{(k)}) \quad \forall j = 1, \dots, n$$

- **Pure torsion case:** the first inclusion is always strict, and *either*
 - ▶ **λ can be simplified:** \exists a reduced toric r -tuple, said **simple**, s.t. $\text{Res}_j(\lambda) = \bigcap_{k=1}^r \text{Res}_j^+(\eta^{(k)}) \quad \forall j$, or
 - ▶ **λ cannot be simplified:** \forall reduced toric r -tuple, $\exists j$ s.t. the inclusions in (5) are strict.

Impure Torsion Case

Main Theorem in the impure torsion case

Main Theorem

f in the impure torsion case is holomorphically normalizable



*f commutes with a holom. effective \mathbb{T}^{r-1} -action on (\mathbb{C}^n, O) ,
 $r = \text{tordeg}(\lambda)$, with columns of Θ that are reduced torsion-free toric
vectors associated to λ .*

Pure Torsion Case

λ can be simplified

Main Theorem

f in the pure torsion case s.t. λ can be simplified is holomorphically normalizable



Pure Torsion Case

λ can be simplified

Main Theorem

f in the pure torsion case s.t. λ can be simplified is holomorphically normalizable



- *df_0 diagonalizable: f commutes with a holom. effective \mathbb{T}^r -action on $(\mathbb{C}^n, 0)$, $r = \text{tordeg}(\lambda)$, with columns of Θ that are a **reduced simple r -tuple of toric vectors associated to λ** ;*

Pure Torsion Case

λ can be simplified

Main Theorem

f in the pure torsion case s.t. λ can be simplified is holomorphically normalizable



- *df_O diagonalizable: f commutes with a holom. effective \mathbb{T}^r -action on (\mathbb{C}^n, O) , $r = \text{tordeg}(\lambda)$, with columns of Θ that are a **reduced simple r -tuple of toric vectors associated to λ** ;*
- *df_O not diagonalizable: if $\lambda_j = \lambda_h \Rightarrow \eta_j^{(k)} = \eta_h^{(k)} \quad \forall k = 1, \dots, r$, then same statement as above.*

Pure Torsion Case

λ cannot be simplified

Proposition (–, 2009)

f in the pure torsion case s.t. λ cannot be simplified. If f commutes with a holom. effective \mathbb{T}^r -action on (\mathbb{C}^n, O) , $r = \text{tordeg}(\lambda)$, with columns of Θ that are reduced toric vectors associated to λ , then f is holomorphically normalizable.

Pure Torsion Case

λ cannot be simplified

Proposition (–, 2009)

f in the pure torsion case s.t. λ cannot be simplified. If f commutes with a holom. effective \mathbb{T}^r -action on (\mathbb{C}^n, O) , $r = \text{tordeg}(\lambda)$, with columns of Θ that are reduced toric vectors associated to λ , then f is holomorphically normalizable.

Proposition (–, 2009)

f in the pure torsion case s.t. λ cannot be simplified. If f is holomorphically normalizable, then f commutes with a holom. effective \mathbb{T}^{r-1} -action on (\mathbb{C}^n, O) , $r = \text{tordeg}(\lambda)$, with columns of Θ that are reduced torsion-free toric vectors associated to λ .

Remark: Torsion

Écalle, in 1992, introduced the following notion of torsion.

Definition

The *torsion* of $\lambda \in (\mathbb{C}^*)^n$ is the natural integer τ such that

$$\frac{1}{\tau}\mathbb{Z} = \mathbb{Q} \cap \left(\mathbb{Z} \bigoplus_{1 \leq j \leq n} \left(\frac{\log(\lambda_j)}{2\pi i} \mathbb{Z} \right) \right).$$

Remark: Torsion

Écalle, in 1992, introduced the following notion of torsion.

Definition

The *torsion* of $\lambda \in (\mathbb{C}^*)^n$ is the natural integer τ such that

$$\frac{1}{\tau}\mathbb{Z} = \mathbb{Q} \cap \left(\mathbb{Z} \bigoplus_{1 \leq j \leq n} \left(\frac{\log(\lambda_j)}{2\pi i} \mathbb{Z} \right) \right).$$

Lemma (–, 2009)

$\lambda \in (\mathbb{C}^*)^n$ is *torsion-free* \iff its torsion is 1

Remarks

- algorithmic way to compute resonances

Remarks

- algorithmic way to compute resonances
- examples for each case

Remarks

- algorithmic way to compute resonances
- examples for each case \implies consistent classification

Remarks

- algorithmic way to compute resonances
- examples for each case \implies consistent classification
- torsion is not enough to measure the difference between germs of holomorphic vector fields and germs of biholomorphisms

Remarks

- algorithmic way to compute resonances
- examples for each case \implies consistent classification
- torsion is not enough to measure the difference between germs of holomorphic vector fields and germs of biholomorphisms
- example of techniques to construct torus actions

Construction of Torus Action

Theorem (–, 2009)

Let f be as above and **commute with a set of integrable holomorphic vector fields** X_1, \dots, X_m , $1 \leq m \leq n$. Then f commutes with a holom. effective \mathbb{T}^r -action on (\mathbb{C}^n, O) , $r = \text{tordeg}(X_1)$, with columns of the weight matrix that are a r -tuple of toric vectors associated to X_1 .

Where, if $1 \leq m \leq n$, f **commute with a set of integrable holomorphic vector fields** if $\exists X_1, \dots, X_m$ s.t.

$$df(X_j) = X_j \circ f$$

$\forall j = 1, \dots, m$ that are **integrable**, i.e.,

- X_1, \dots, X_m germs of holom. v.f. of (\mathbb{C}^n, O) , $X_j(O) = 0$, $\text{order}(X_j) = 1$, $[X_j, X_k] = 0 \forall j, k$, and $X_1 \wedge \dots \wedge X_m \neq 0$;
- $\exists g_1, \dots, g_{n-m}$ germs of holom. functions of (\mathbb{C}^n, O) s.t. $X_j(g_k) = 0 \forall j, k$, and $dg_1 \wedge \dots \wedge dg_{n-m} \neq 0$.

Construction of Torus Action

Toric degree of a vector field

Writing $X = \sum_{j=1}^n \varphi_j \mathbf{z}_j \frac{\partial}{\partial \mathbf{z}_j} + \dots$, the **toric degree of X** is the min $r \in \mathbb{N}$ s.t. $\exists \alpha_1, \dots, \alpha_r \in \mathbb{C}^*$ and $\exists \theta^{(1)}, \dots, \theta^{(r)} \in \mathbb{Z}^n$ s.t.

$$\varphi = \sum_{k=1}^r \alpha_k \theta^{(k)}.$$

Thanks

Examples

Example (Torsion-free case)

$$\left[\sqrt{2} \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix} + 2i \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} \right] \in (\mathbb{C}/\mathbb{Z})^3$$

Example (Torsion case)

$$\left[\frac{1}{7} \begin{pmatrix} 3 \\ 2 \\ -1 \end{pmatrix} + 2i \begin{pmatrix} 2 \\ 3 \\ 1 \end{pmatrix} \right] \in (\mathbb{C}/\mathbb{Z})^3$$

Example (Impure torsion case)

$$\left[\frac{1}{3} \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix} + \sqrt{2} \begin{pmatrix} -12 \\ 0 \\ 0 \\ 1 \end{pmatrix} + \sqrt{3} \begin{pmatrix} 0 \\ 5 \\ 2 \\ 0 \end{pmatrix} \right] \in (\mathbb{C}/\mathbb{Z})^4$$

has toric degree 3.

Example (Impure torsion case)

$$\text{Res}_1^+(\eta^{(2)}) = \{(q_1, q_2, q_3, 12(q_1 - 1)) \in \mathbb{N}^4 \mid 13q_1 + q_2 + q_3 \geq 14\}$$

$$\text{Res}_2^+(\eta^{(2)}) = \{(q_1, q_2, q_3, 12q_1) \in \mathbb{N}^4 \mid 13q_1 + q_2 + q_3 \geq 2\}$$

$$\text{Res}_3^+(\eta^{(2)}) = \text{Res}_2^+(\eta^{(2)})$$

$$\text{Res}_4^+(\eta^{(2)}) = \{(q_1, q_2, q_3, 12q_1 + 1) \in \mathbb{N}^4 \mid 13q_1 + q_2 + q_3 \geq 1\},$$

and

$$\text{Res}_1^+(\eta^{(3)}) = \{(q_1, 0, 0, q_4) \in \mathbb{N}^4 \mid q_1 + q_4 \geq 2\}$$

$$\text{Res}_2^+(\eta^{(3)}) = \{(q_1, 1, 0, q_4) \in \mathbb{N}^4 \mid q_1 + q_4 \geq 1\}$$

$$\text{Res}_3^+(\eta^{(3)}) = \{(q_1, 0, 1, q_4) \in \mathbb{N}^4 \mid q_1 + q_4 \geq 1\}$$

$$\text{Res}_4^+(\eta^{(3)}) = \text{Res}_1^+(\eta^{(3)}).$$

$$\forall P = (p, 0, 0, 12p) \text{ with } p \geq 1 \quad \langle P, \eta^{(1)} \rangle = 12p \in 3\mathbb{Z}.$$

Then it is easy to verify that for $j = 1, \dots, 4$

$$\text{Res}_j([\varphi]) = \text{Res}_j^+(\eta^{(2)}) \cap \text{Res}_j^+(\eta^{(3)}).$$

Examples

Example (Pure torsion case that can be simplified)

$$[\varphi] = \left[\frac{1}{3} \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} + \sqrt{2} \begin{pmatrix} 1 \\ 6 \\ 0 \\ 0 \end{pmatrix} + \sqrt{3} \begin{pmatrix} 0 \\ 0 \\ -1 \\ 5 \end{pmatrix} \right] \in (\mathbb{C}/\mathbb{Z})^4,$$

has toric degree 3. We have $\text{Res}_j^+(\eta^{(1)}) = \emptyset$, for $j = 1, \dots, 4$, and it is not difficult to verify that

$$\text{Res}_2(\lambda) = \{(0, 1, 5q, q) \mid q \in \mathbb{N}^*\} \neq \emptyset$$

$$\text{Res}_j(\lambda) = \text{Res}_j^+(\eta^{(2)}) \cap \text{Res}_j^+(\eta^{(3)}) \quad j = 1, 3, 4.$$

Examples

Example (Pure torsion case that can be simplified)

However, we can write

$$[\varphi] = \left[\frac{1}{3} \begin{pmatrix} 1 \\ -2 \\ 1 \\ -5 \end{pmatrix} + \sqrt{2} \begin{pmatrix} 1 \\ 6 \\ 0 \\ 0 \end{pmatrix} + \sqrt{3} \begin{pmatrix} 0 \\ 0 \\ -1 \\ 5 \end{pmatrix} \right],$$

and it is not difficult to verify that, in this representation, we have, for $j = 1, \dots, 4$

$$\mathrm{Res}_j(\lambda) = \bigcap_{k=1}^3 \mathrm{Res}_j^+(\xi^{(k)}).$$

Examples

Example (Pure torsion case that cannot be simplified)

$$\left[\frac{1}{7} \begin{pmatrix} 1 \\ 3 \end{pmatrix} + \sqrt{2} \begin{pmatrix} 1 \\ -6 \end{pmatrix} \right] \in (\mathbb{C}/\mathbb{Z})^2,$$

has toric degree 2 and torsion 7. We have

$$\text{Res}_1^+(\eta^{(2)}) = \{(6h+1, h) \mid h \geq 1\}, \quad \text{Res}_2^+(\eta^{(2)}) = \{(6h, h+1) \mid h \geq 1\},$$

$$\text{Res}_1^+(\eta^{(1)}) \cap \text{Res}_1^+(\eta^{(2)}) = \text{Res}_2^+(\eta^{(1)}) \cap \text{Res}_2^+(\eta^{(2)}) = \emptyset,$$

then

$$\text{Res}_1^+(\eta^{(2)}) \supset \text{Res}_1(\lambda) = \{(42h+1, 7h) \mid h \geq 1\} \supset \text{Res}_1^+(\eta^{(1)}) \cap \text{Res}_1^+(\eta^{(2)})$$

$$\text{Res}_2^+(\eta^{(2)}) \supset \text{Res}_2(\lambda) = \{(42h, 7h+1) \mid h \geq 1\} \supset \text{Res}_2^+(\eta^{(1)}) \cap \text{Res}_2^+(\eta^{(2)}).$$

Furthermore, it is easy to check that λ cannot be simplified.