

# Chapter 3

## Selfadjoint operators

[Draft version, November 16, 2022]


Let  $\mathcal{H}$  be a Hilbert space.

### 3.1 Selfadjoint operators

#### 3.1.1 Symmetric operators

**Definition 3.1.** Let  $A$  be an operator on  $\mathcal{H}$ . We say that  $A$  is symmetric if

$$\forall \varphi, \psi \in \text{Dom}(A), \quad \langle A\varphi, \psi \rangle_{\mathcal{H}} = \langle \varphi, A\psi \rangle_{\mathcal{H}}.$$

 Ex. 3.1

*Remark 3.2.* If  $A$  is symmetric then  $\langle A\varphi, \varphi \rangle_{\mathcal{H}} \in \mathbb{R}$  for all  $\varphi \in \text{Dom}(A)$ . The converse is also true, as can be seen from the polarization formula

$$\begin{aligned} \forall \varphi, \psi \in \text{Dom}(A), \quad 4\langle A\varphi, \psi \rangle &= \langle A(\varphi + \psi), \varphi + \psi \rangle - \langle A(\varphi - \psi), \varphi - \psi \rangle \\ &\quad + i\langle A(\varphi + i\psi), \varphi + i\psi \rangle - i\langle A(\varphi - i\psi), \varphi - i\psi \rangle. \end{aligned}$$

**Definition 3.3.** Let  $A$  be a symmetric operator on  $\mathcal{H}$ .

- (i) We say that  $A$  is non-negative (and we write  $A \geq 0$ ) if  $\langle A\varphi, \varphi \rangle_{\mathcal{H}} \geq 0$  for all  $\varphi \in \text{Dom}(A)$ .
- (ii) We say that  $A$  is semi-bounded from below if there exists  $\gamma \in \mathbb{R}$  such that  $A - \gamma \geq 0$  (we can write  $A \geq \gamma$ ). Equivalently,  $\langle A\varphi, \varphi \rangle_{\mathcal{H}} \geq \gamma \|\varphi\|_{\mathcal{H}}^2$  for all  $\varphi \in \text{Dom}(A)$ . In this case we say that  $\gamma$  is a lower bound for  $A$ .

**Proposition 3.4.** Let  $A$  be a symmetric and densely defined operator on  $\mathcal{H}$ . Then  $A^*$  is a closed extension of  $A$ .

*Proof.* Let  $\psi \in \text{Dom}(A)$ . For all  $\varphi \in \text{Dom}(A)$  we have  $\langle A\varphi, \psi \rangle = \langle \varphi, A\psi \rangle$  so  $\psi \in \text{Dom}(A^*)$  and  $A^*\psi = A\psi$ . This proves that  $A^*$  is an extension of  $A$ . Moreover  $A^*$  is closed by Proposition 2.48.  $\square$

**Proposition 3.5.** Let  $A$  be a symmetric operator on  $\mathcal{H}$ . The eigenvalues of  $A$  (if any) are real, and two eigenvectors of  $A$  associated to different eigenvalues are orthogonal.

*Proof.* • Let  $\lambda \in \mathbb{C}$  and assume that for some  $\varphi \in \text{Dom}(A) \setminus \{0\}$  we have  $A\varphi = \lambda\varphi$ . Then  $\lambda \|\varphi\|_{\mathcal{H}}^2 = \langle A\varphi, \varphi \rangle_{\mathcal{H}} \in \mathbb{R}$ . This implies that  $\lambda \in \mathbb{R}$ .

• Now let  $\lambda, \mu$  be two distinct eigenvalues of  $A$ . Let  $\varphi \in \ker(A - \lambda)$  and  $\psi \in \ker(A - \mu)$ . Then

$$(\mu - \lambda)\langle \psi, \varphi \rangle_{\mathcal{H}} = \langle \mu\psi, \varphi \rangle_{\mathcal{H}} - \langle \psi, \lambda\varphi \rangle_{\mathcal{H}} = \langle A\psi, \varphi \rangle_{\mathcal{H}} - \langle \psi, A\varphi \rangle_{\mathcal{H}} = 0.$$

Since  $\mu - \lambda \neq 0$ , this implies that  $\langle \psi, \varphi \rangle_{\mathcal{H}} = 0$ .  $\square$

**Proposition 3.6.** Let  $A$  be a non-negative and densely defined operator. Let  $\varphi \in \text{Dom}(A)$  be such that  $\langle A\varphi, \varphi \rangle_{\mathcal{H}} = 0$ . Then  $A\varphi = 0$ .

*Proof.* Since  $A$  is non-negative, we can apply the Cauchy-Schwarz inequality to the sesquilinear form  $(\zeta, \psi) \mapsto \langle A\zeta, \psi \rangle_{\mathcal{H}}$  on  $\text{Dom}(A)$ . Then for all  $\psi \in \text{Dom}(A)$  we have

$$|\langle A\varphi, \psi \rangle_{\mathcal{H}}| \leq |\langle A\varphi, \varphi \rangle_{\mathcal{H}}| |\langle A\psi, \psi \rangle_{\mathcal{H}}| = 0.$$

Since  $\text{Dom}(A)$  is dense in  $\mathcal{H}$ , this proves that  $A\varphi = 0$ .  $\square$

**Proposition 3.7.** *Let  $A$  be a symmetric operator on  $\mathcal{H}$ .*

(i) *For  $z \in \mathbb{C} \setminus \mathbb{R}$  and  $\varphi \in \text{Dom}(A)$  we have*

$$\|(A - z)\varphi\|_{\mathcal{H}} \geq |\text{Im}(z)| \|\varphi\|_{\mathcal{H}}.$$

(ii) *Assume moreover that  $A$  is bounded from below and let  $\gamma \in \mathbb{R}$  be such that  $A \geq \gamma$ . Then for  $\lambda < \gamma$  and  $\varphi \in \text{Dom}(A)$  we have*

$$\|(A - z)\varphi\|_{\mathcal{H}} \geq (\gamma - \lambda) \|\varphi\|_{\mathcal{H}}.$$

*Proof.* Let  $\varphi \in \text{Dom}(A)$ .

• Let  $z \in \mathbb{C} \setminus \mathbb{R}$ ,  $\lambda = \text{Re}(z)$  and  $\varepsilon = \text{Im}(z)$ . We have

$$\|(A - z)\varphi\|^2 = \|(A - \lambda)\varphi\|^2 + \varepsilon^2 \|\varphi\|^2 + 2 \text{Re} \langle (A - \lambda)\varphi, -i\varepsilon\varphi \rangle.$$

Since

$$\langle (A - \lambda)\varphi, -i\varepsilon\varphi \rangle = i\varepsilon \langle A\varphi, \varphi \rangle - i\varepsilon\lambda \|\varphi\|^2 \in i\mathbb{R},$$

this gives

$$\|(A - z)\varphi\|^2 = \|(A - \lambda)\varphi\|^2 + \varepsilon^2 \|\varphi\|^2 \geq \varepsilon^2 \|\varphi\|^2. \quad \square$$

• Similarly, if  $A - \gamma \geq 0$  then for  $\lambda \in ]-\infty, \gamma[$  we have

$$\begin{aligned} \|(A - \lambda)\varphi\|_{\mathcal{H}}^2 &= \|(A - \gamma)\varphi\|_{\mathcal{H}}^2 + (\gamma - \lambda)^2 \|\varphi\|_{\mathcal{H}}^2 + 2(\gamma - \lambda) \text{Re} \langle (A - \gamma)\varphi, \varphi \rangle_{\mathcal{H}} \\ &\geq (\gamma - \lambda)^2 \|\varphi\|_{\mathcal{H}}^2, \end{aligned}$$

and the second statement follows.

### 3.1.2 Selfadjoint operators

**Definition 3.8.** *An operator  $A$  on  $\mathcal{H}$  is said to be selfadjoint if it is densely defined and  $A^* = A$ .*

$\varnothing$  Ex. 3.2

**Definition 3.9.** *An operator  $A$  on  $\mathcal{H}$  is said to be skew-adjoint if it is densely defined and  $A^* = -A$ .*

*Remark 3.10.* An operator  $A$  is skew-adjoint if and only if  $iA$  is selfadjoint. Then one usually only discusses the properties of selfadjoint operators, and we can deduce similar properties for skew-adjoint operators.

*Example 3.11.* • The Laplacian  $H = -\Delta$  on  $L^2(\mathbb{R}^d)$  (with domain  $\text{Dom}(H) = H^2(\mathbb{R}^d)$ ) is selfadjoint. The Laplacian  $H_0 = -\Delta$  with domain  $C_0^\infty(\mathbb{R}^d)$  is symmetric but not self-adjoint (in particular  $C_0^\infty(\mathbb{R}^d) \subsetneq H^2(\mathbb{R}^d) \subset \text{Dom}(H_0^*)$ ). However,  $H_0$  has a selfadjoint extension ( $H$ ).

*Example 3.12.* The Dirichlet and Neumann Laplacians on  $]0, 1[$  (introduced in Section 2.5.2) are selfadjoint.

*Example 3.13.* The harmonic oscillator introduced in Section 2.4 is selfadjoint.

*Remark 3.14.* A selfadjoint operator is closed by Proposition 2.47.

**Proposition 3.15.** *Let  $A$  be a selfadjoint operator on  $\mathcal{H}$ . Then*

$$\overline{\text{Ran}(A)} = \ker(A)^\perp.$$

**Definition 3.16.** Let  $A$  be a selfadjoint operator on  $\mathcal{H}$ . Let  $F$  be a subspace of  $\mathcal{H}$ . We say that  $F$  is reducing for the operator  $A$  (or that it reduces  $A$ ) if it is closed and the orthogonal projection  $\Pi$  on  $F$  satisfies  $\Pi A \subset A \Pi$ .

*Remark 3.17.* If  $F$  reduces  $A$ , then so does  $F^\perp$ .

**Proposition 3.18.** Let  $A$  be a selfadjoint operator on  $\mathcal{H}$ . Let  $F$  be a reducing subspace for  $A$ . Then the restriction  $A_F$  of  $A$  on  $F$  is a selfadjoint operator on  $F$ .

*Proof.* The restriction  $A_F$  of the symmetric operator  $A$  is symmetric. Let  $\psi \in \text{Dom}(A_F^*)$ . For all  $\varphi \in \text{Dom}(A) \cap F$  we have

$$\langle A_F \varphi, \psi \rangle = \langle \varphi, A_F^* \psi \rangle$$

Then for all  $\varphi = \varphi_F + \varphi^\perp$  with  $\varphi_F \in F$  and  $\varphi^\perp \in F^\perp$  we have

$$\langle A \varphi, \psi \rangle = \langle A_F \varphi_F, \psi \rangle = \langle \varphi_F, A_F^* \psi \rangle = \langle \varphi, A_F^* \psi \rangle.$$

This proves that  $\psi \in \text{Dom}(A^*) = \text{Dom}(A)$ , so  $\psi \in \text{Dom}(A) \cap F = \text{Dom}(A_F)$  and  $A_F^* \psi = A^* \psi = A \psi = A_F \psi$ . This proves that  $A_F^* \subset A_F$ , and finally  $A_F$  is selfadjoint by Proposition 3.4.  $\square$

**Proposition 3.19.** Let  $A$  be a selfadjoint operator on  $\mathcal{H}$ . Then  $\ker(A)$  is reducing for  $A$ .

*Proof.* Since  $A$  is closed,  $\ker(A)$  is closed in  $\mathcal{H}$ . Let  $\Pi$  be the orthogonal projection on  $\ker(A)$ . For all  $\varphi \in \text{Dom}(A)$  we have  $A \varphi \in \text{Ran}(A) \subset \ker(A)^\perp$ , so  $\Pi A \varphi = 0$ . On the other hand we have  $\Pi \varphi \in \ker(A) \subset \text{Dom}(A)$  and  $A \Pi \varphi = 0$ . This proves that  $\Pi A \subset A \Pi$ .  $\square$

**Proposition 3.20.** If  $A$  and  $B$  are two selfadjoint operators on  $\mathcal{H}$  such that  $A \subset B$  then  $A = B$ .

*Proof.* We have  $A \subset B = B^* \subset A^* = A$ , so  $A = B$ .  $\square$

### 3.1.3 A criterion for self-adjointness

**Proposition 3.21.** Let  $A$  be a symmetric and densely defined operator on  $\mathcal{H}$  and  $z \in \mathbb{C} \setminus \mathbb{R}$ . Then the following assertions are equivalent.

- (i)  $A$  is self-adjoint.
- (ii)  $A$  is closed and  $z, \bar{z} \in \rho(A)$ .
- (iii)  $A$  is closed and  $\ker(A^* - z) = \ker(A^* - \bar{z}) = \{0\}$ .
- (iv)  $\text{Ran}(A - z) = \text{Ran}(A - \bar{z}) = \mathcal{H}$ .

*Proof.* • (i)  $\Rightarrow$  (iii). Assume that  $A$  is self-adjoint. In particular,  $A$  is closed. Moreover,  $\ker(A^* - z) = \ker(A - z) = \{0\}$  by Proposition 3.7. Similarly,  $\ker(A^* - \bar{z}) = \{0\}$ .

• (iii)  $\Rightarrow$  (iv). By Proposition 2.46 we have  $\text{Ran}(A - z) = \ker(A^* - \bar{z})^\perp = \{0\}$ , so  $\text{Ran}(A - z)$  is dense in  $\mathcal{H}$ . On the other hand,  $(A - z)$  has closed range by Propositions 3.7 and 2.34, so  $\text{Ran}(A - z) = \mathcal{H}$ . Similarly,  $\text{Ran}(A - \bar{z}) = \mathcal{H}$ .

• (iv)  $\Rightarrow$  (i). We already know by Proposition 3.4 that  $A^*$  is an extension of  $A$ . Let  $\varphi \in \text{Dom}(A^*)$ . Since  $(A - z)$  is surjective, there exists  $\psi \in \text{Dom}(A)$  such that  $(A^* - z)\varphi = (A - z)\psi$ . On the other hand, we have  $\psi \in \text{Dom}(A^*)$  and  $A^* \psi = A \psi$  so  $(A^* - z)\varphi = (A^* - z)\psi$ . By Proposition 2.46 we have  $\ker(A^* - z) = \text{Ran}(A - \bar{z})^\perp = \{0\}$ , so  $\varphi = \psi \in \text{Dom}(A)$ . This proves that  $\text{Dom}(A) = \text{Dom}(A^*)$ , and hence  $A = A^*$ .

• (ii)  $\Rightarrow$  (iv) is clear.

• (iii)  $\Rightarrow$  (ii).  $A$  is closed by (iii). By Proposition 3.7 we already know that  $A - z$  is injective. It is surjective by (iv) so it is bijective and  $z \in \rho(A)$ . Similarly,  $\bar{z} \in \rho(A)$ .  $\square$   Ex. 3.3, 3.4

The proof of the implication (iv)  $\implies$  (i) also holds if  $z \in \mathbb{R}$ . This gives the following sufficient condition.

**Corollary 3.22.** Let  $A$  be a symmetric operator on  $\mathcal{H}$ . Assume that there exists  $\lambda \in \mathbb{R}$  such that  $\text{Ran}(A - \lambda) = \mathcal{H}$ . Then  $A$  is selfadjoint.

Combined with Proposition 3.7 this also gives the following result.

**Corollary 3.23.** Let  $A$  be a symmetric operator on  $\mathcal{H}$ . Assume that  $A \geq \gamma$  for some  $\gamma \in \mathbb{R}$ . If there exists  $\lambda \in ]-\infty, \gamma[$  such that  $\text{Ran}(A - \lambda)$  is dense in  $\mathcal{H}$ , then  $A$  is selfadjoint.

### 3.1.4 Essentially selfadjoint operators

We have seen that if  $A$  is symmetric then  $A \subset A^*$ . It may happen that  $A$  is not selfadjoint because we have chosen the domain too small. Given a symmetric operator, the question is then whether it has a selfadjoint extension.

We know from Proposition 3.4 that a densely defined and symmetric operator is always closable, so the first try is to look at its closure.

**Definition 3.24.** *Let  $A$  be a densely defined symmetric operator on  $\mathcal{H}$ . We say that  $A$  is essentially selfadjoint if its closure  $\overline{A}$  is selfadjoint.*

**Proposition 3.25.** *Let  $A$  be a densely defined symmetric operator on  $\mathcal{H}$ . Then  $A$  is essentially selfadjoint if and only if  $\overline{A} = A^*$ .*

*Proof.* • By Proposition 2.48 we have  $\overline{A^*} = A^*$ . If  $A$  is essentially selfadjoint, we also have  $\overline{A^*} = \overline{A}$ , and hence  $\overline{A} = A^*$ .

• Conversely, assume that  $\overline{A} = A^*$ . By Proposition 2.48 again we have  $A^{**} = \overline{A}$ , so  $\overline{A^*} = A^{**} = \overline{A}$ .  $\square$

We will see below that a symmetric operator may have many selfadjoint extensions. However, when it is essentially selfadjoint, the extension is unique.

**Proposition 3.26.** *Let  $A$  be a densely defined symmetric operator on  $\mathcal{H}$ . If  $A$  is essentially selfadjoint then  $\overline{A}$  is the unique selfadjoint extension of  $A$ .*

*Proof.* Let  $B$  be a selfadjoint extension of  $A$ . Since it is a closed extension of  $A$ , it is an extension of  $\overline{A}$ . Since  $B$  and  $\overline{A}$  are selfadjoint, we have  $B = \overline{A}$  by Proposition 3.20.  $\square$

**Proposition 3.27.** *Let  $A$  be a densely defined symmetric operator on  $\mathcal{H}$ . Let  $z \in \mathbb{C} \setminus \mathbb{R}$ . The following assertions are equivalent.*

- (i)  $A$  is essentially selfadjoint ;
- (ii)  $\ker(A^* - z) = \ker(A^* - \overline{z}) = \{0\}$  ;
- (iii)  $\overline{\text{Ran}(A - z)} = \overline{\text{Ran}(A - \overline{z})} = \mathcal{H}$ .

*Proof.* • Assume that  $A$  is essentially selfadjoint. In particular,  $A$  is closable and  $\overline{A^*} = A^*$  by Proposition 2.48. By Proposition 3.21 applied to the selfadjoint operator  $\overline{A}$ , we have  $\ker(A^* - z) = \ker(A^* - \overline{z}) = \{0\}$ .

- Conversely, assume that (ii) holds. Since  $\overline{A^*} \subset A^*$  we have  $\ker(\overline{A^*} - z) = \ker(\overline{A^*} - \overline{z}) = \{0\}$ . By Proposition 3.21,  $\overline{A}$  is selfadjoint.
- Finally (ii) and (iii) are equivalent by Proposition 2.46.  $\square$

### 3.1.5 Examples of closed symmetric operators which are not essentially selfadjoint

We consider on  $L^2(0, 1)$  the operator  $A$  which acts as

$$A = i \frac{d}{dx}$$

on the domain

$$\text{Dom}(A) = H_0^1(0, 1).$$

Then  $A$  is closed (by Example 2.29 and symmetric: for  $u, v \in H_0^1(0, 1)$  we have by the Green formula

$$\begin{aligned} \langle Au, v \rangle_{L^2(0,1)} &= i \int_0^1 u'(x) \overline{v(x)} \, dx = i(u(1)\overline{v(1)} - u(0)\overline{v(0)}) - i \int_0^1 u(x) \overline{v'(x)} \, dx \\ &= \langle u, Av \rangle_{L^2(0,1)}. \end{aligned}$$

Notice that for the boundary terms it was not necessary that both  $u$  and  $v$  vanish.

Now we compute  $A^*$ . Let  $v \in \text{Dom}(A^*)$ . We have  $v \in L^2(0, 1)$  and for all  $\phi \in C_0^\infty(]0, 1[)$  we have

$$\int_{\mathbb{R}} i\phi'(x)\overline{v(x)} dx = \langle A\phi, v \rangle_{L^2(0,1)} = \langle \phi, A^*v \rangle_{L^2(0,1)} = \int_{\mathbb{R}} \phi(x)\overline{(A^*v)(x)} dx.$$

This prove that in the sense of distributions we have  $v' \in L^2(0, 1)$  and

$$A^*v = iv'.$$

Conversely, if  $v \in H^1(0, 1)$  then the same computation as above shows that

$$\forall u \in \text{Dom}(A), \quad \langle iu', v \rangle_{L^2(0,1)} = \langle u, iv' \rangle_{L^2(0,1)},$$

so  $v \in \text{Dom}(A^*)$  (and we recover  $A^*v = iv'$ ). This proves that  $\text{Dom}(A^*) = H^1(0, 1) \neq \text{Dom}(A)$ . Thus  $A$  is not selfadjoint.

Notice that for  $z \in \mathbb{C}$  the function  $x \mapsto e^{-izx}$  belongs to  $\ker(A^* - z)$ . In particular,  $\ker(A^* - z) \neq \{0\}$ . By Proposition 3.21, this confirms that  $A$  cannot be selfadjoint. It is not even essentially selfadjoint. Moreover, for  $z \in \mathbb{C}$  we have by Proposition 2.46

$$\overline{\text{Ran}(A - z)} = \ker(A^* - \bar{z})^\perp \neq \mathcal{H}.$$

This proves that  $\sigma(A) = \mathbb{C}$ .

Now the question is: does  $A$  have a selfadjoint extension? The answer is: yes, many! Assume that  $\tilde{A}$  is a selfadjoint extension of  $A$ . Then  $\tilde{A} = \tilde{A}^* \subset A^*$ . Let  $v \in \text{Dom}(\tilde{A}) \setminus \text{Dom}(A)$ . For all  $u \in \text{Dom}(\tilde{A})$  we have

$$0 = \langle \tilde{A}u, v \rangle - \langle u, \tilde{A}v \rangle = i(u(1)\bar{v}(1) - u(0)\bar{v}(0)).$$

Assume that  $v(1) = 0$ . Since  $v$  is not in  $\text{Dom}(A)$  we have  $v(0) \neq 0$ . Then for all  $u \in \text{Dom}(\tilde{A})$  we have  $u(0) = 0$ . This gives a contradiction since  $v(0) \neq 0$ . This proves that  $v(1) \neq 0$ . We set  $\alpha = \bar{v}(0)/\bar{v}(1)$ . Then for all  $u \in \text{Dom}(\tilde{A})$  we have

$$u(1) = \alpha u(0).$$

In particular we have  $v(1) = \alpha v(0)$ . Since by definition we have  $v(0) = \bar{\alpha}v(1)$ , this proves that  $|\alpha| = 1$ . This proves that there exists  $\alpha \in \mathbb{U}$  such that  $\text{Dom}(\tilde{A}) \subset D_\alpha$ , where we have set

$$D_\alpha = \{u \in H^1(0, 1) : u(1) = \alpha u(0)\}.$$

For  $\alpha \in \mathbb{U}$  we denote by  $A_\alpha$  the operator defined by  $A_\alpha u = iu'$  for  $u$  in  $\text{Dom}(A_\alpha) = D_\alpha$ . In particular,  $A_\alpha$  is an extension of  $A$  and  $A^*$  is an extension of  $A_\alpha$  for all  $\alpha$ .

We check that  $A_\alpha$  is selfadjoint. For  $u, v \in \text{Dom}(A_\alpha)$  we have

$$\begin{aligned} \langle A_\alpha u, v \rangle &= i \int_{\mathbb{R}} u'(x)\bar{v}(x) dx \\ &= i\bar{u}(1)v(1) - iu(0)\bar{v}(0) - i \int_{\mathbb{R}} u(x)\bar{v}'(x) dx \\ &= i(|\alpha|^2 - 1)u(0)\bar{v}(0) - i \int_{\mathbb{R}} u(x)\bar{v}'(x) dx \\ &= \langle u, A_\alpha v \rangle. \end{aligned}$$

Then  $A_\alpha$  is symmetric, and hence  $A_\alpha^*$  is an extension of  $A_\alpha$ . Now let  $v \in \text{Dom}(A_\alpha^*)$ . The same computation with  $u \in C_0^\infty(]0, 1[)$  shows that  $v \in H^1(0, 1)$  and  $A_\alpha^*v = iv'$ . Then for all  $u \in \text{Dom}(A_\alpha)$  we have

$$0 = \langle A_\alpha u, v \rangle - \langle u, A_\alpha^*v \rangle = -iu(0)(\bar{\alpha}v(1) - v(0)).$$

This proves that  $v(1) = \alpha v(0)$ , so  $v \in \text{Dom}(A_\alpha)$ , and finally  $A_\alpha^* = A_\alpha$ .

All this proves that the operators  $A_\alpha$  for  $\alpha \in \mathbb{U}$  are the selfadjoint extensions of  $A$ .

Moreover we have seen that if  $\tilde{A}$  is a selfadjoint extension of  $A$  then we have  $\tilde{A} \subset A_\alpha$  for some  $\alpha \in \mathbb{U}$ , and hence  $\tilde{A} = A_\alpha$ . So finally, the operators  $A_\alpha$  for  $\alpha \in \mathbb{U}$  are exactly the selfadjoint extensions of  $A$ .

*Example 3.28.* We consider the previous example but on  $L^2(0, +\infty)$ :

$$A = i \frac{d}{dx}, \quad \text{Dom}(A) = H_0^1(0, +\infty).$$

With the same proof we see that  $A$  is symmetric with  $\text{Dom}(A^*) = H^1(0, +\infty)$ . The difference is that in this case  $A$  has no selfadjoint extension. Indeed, assume by contradiction that  $\tilde{A}$  is a selfadjoint extension of  $A$ . We can check by direct computation that  $\text{Ran}(A + i) = L^2(0, +\infty)$  (or equivalently, that  $\text{Ker}(A^* - i) = \{0\}$ ), so  $A$  has no selfadjoint extension by Exercise 3.3.

### 3.1.6 Friedrichs extension

We have seen in the previous paragraph that a symmetric operator which is not selfadjoint can have many selfadjoint extensions, and it is also possible that it does not have any.

In this paragraph we consider the case of lower semibounded symmetric operators and choose in an abstract setting a selfadjoint extension. This ensures in particular that such an operator has at least one selfadjoint extension.

**Definition 3.29.** Let  $A$  be a densely defined lower bounded symmetric operator on  $\mathcal{H}$ . Let  $\alpha_0 \in \mathbb{R}$  be such that  $A + \alpha_0 \geq 0$ . For  $\alpha > \alpha_0$  we consider the quadratic form associated to  $A + \alpha$  on  $\text{Dom}(A + \alpha) = \text{Dom}(A)$

$$\mathfrak{q}_{A+\alpha} : \varphi \in \text{Dom}(A) \mapsto \langle (A + \alpha)\varphi, \varphi \rangle_{\mathcal{H}} = \langle A\varphi, \varphi \rangle_{\mathcal{H}} + \alpha \|\varphi\|_{\mathcal{H}}^2.$$

The closure of  $\text{Dom}(A)$  for the norm  $\|\varphi\|_A = \sqrt{\mathfrak{q}_{A+\alpha}(\varphi)}$  is called the form domain of  $A$ .

The definition of the form domain does not depend on  $\alpha > \alpha_0$ .

*Example 3.30.* We consider on  $L^2(0, 1)$  the Laplacian  $H_0 = -\partial^2$  with domain  $\text{Dom}(H_0) = C_0^\infty(]0, 1[)$ . For all  $u \in C_0^\infty(]0, 1[)$  we have

$$\mathfrak{q}_{H_0+1}(u, u) = \langle -\Delta u, u \rangle_{L^2(0,1)} + \|u\|_{L^2(0,1)}^2 = \|u\|_{H^1(0,1)}^2.$$

The closure of  $C_0^\infty(]0, 1[)$  for the  $H^1$  norm is  $H_0^1(0, 1)$ . Then the form domain of  $H_0$  is  $H_0^1(0, 1)$ .

**Proposition 3.31.** Let  $\mathcal{V}$  be a Hilbert space densely and continuously embedded in  $\mathcal{H}$ . Let  $\mathfrak{q}$  be a continuous sesquilinear form on  $\mathcal{V}$  such that, for some  $\alpha_0 > 0$ ,

$$\forall \varphi \in \mathcal{V}, \quad \mathfrak{q}(\varphi) \geq \alpha_0 \|\varphi\|_{\mathcal{H}}^2. \quad (3.1)$$

Let  $A$  be the operator given by the representation theorem (Theorem 2.52). Then  $A$  is selfadjoint on  $\mathcal{H}$  and  $A \geq \alpha_0$ . Then  $\mathcal{V}$  is the form domain of  $A$ .

*Proof.* For all  $\varphi \in \text{Dom}(A)$  we have  $\langle A\varphi, \varphi \rangle = \mathfrak{q}(\varphi, \varphi)$  by definition of  $A$ . By continuity of  $\mathfrak{q}$  and (3.1), the norm  $\varphi \mapsto \sqrt{\mathfrak{q}(\varphi, \varphi)}$  is equivalent to the norm  $\|\cdot\|_{\mathcal{V}}$ . Since  $\text{Dom}(A)$  is dense in  $\mathcal{V}$  by Theorem 2.52, the closure of  $\text{Dom}(A)$  for  $\|\cdot\|_{\mathcal{V}}$  is  $\mathcal{V}$ . Finally, as for Remark 3.2, since the quadratic form takes real values it is symmetric. Then we deduce that  $A$  is selfadjoint by Theorem 2.52.  $\square$

*Example 3.32.* The form domain of the Dirichlet Laplacian on  $]0, 1[$  (see Example 2.57) is  $H_0^1(0, 1)$  and the form domain of the Neumann Laplacian (see Example 2.56) is  $H^1(0, 1)$ .

**Definition 3.33.** Let  $A$  be a lower bounded symmetric operator on  $\mathcal{H}$  and let  $\mathcal{V}$  be the form domain of  $A$ . Let  $\alpha_0 \in \mathbb{R}$  be such that  $A + \alpha_0 \geq 0$ . Let  $\alpha > 0$ . We denote by  $A_\alpha$  the operator associated to the coercive quadratic form  $\mathfrak{q}_{A+\alpha}$  by the representation theorem (Theorem 2.52). Then we define the Friedrichs extension  $A_F$  of  $A$  by  $A_F = A_\alpha - \alpha$ .

This definition does not depend on the choice of  $\alpha$ .

*Example 3.34.* Let  $H_0$  be the operator of Example 3.30. Its Friedrichs extension is the Dirichlet Laplacian on  $]0, 1[$ .

*Remark 3.35.* If  $A$  is a non-negative selfadjoint operator, we have  $A_F = A$ . Indeed, we consider the quadratic form  $q$  associated with  $A + 1$ . For all  $\varphi, \psi \in \text{Dom}(A)$ ,

$$q(\varphi, \psi) = \langle A\varphi, \psi \rangle_{\mathcal{H}} + \langle \varphi, \psi \rangle_{\mathcal{H}}.$$

Then  $\varphi \mapsto \sqrt{q(\varphi, \varphi)}$  defines a norm on  $\text{Dom}(A)$  and we denote by  $\mathcal{V}$  the closure. Then  $q$  extends to a continuous and coercive form on  $\mathcal{V}$ . We denote by  $T$  the corresponding operator given by the representation theorem. Let  $\varphi \in \text{Dom}(A)$ . For all  $\psi \in \text{Dom}(A)$  we have

$$|q(\varphi, \psi)| \leq (\|A\varphi\|_{\mathcal{H}} + \|\varphi\|_{\mathcal{H}}) \|\psi\|_{\mathcal{H}}.$$

Then, by density of  $\text{Dom}(A)$  in  $\mathcal{V}$  we obtain the same inequality for all  $\psi \in \mathcal{V}$ . Then  $\varphi \in \text{Dom}(T)$  and

$$T\varphi = A\varphi + \varphi.$$

Now let  $\varphi \in \text{Dom}(T)$ . For all  $\psi \in \text{Dom}(A)$  we have

$$|\langle A\psi, \varphi \rangle_{\mathcal{H}}| = |q(\psi, \varphi)| = |q(\varphi, \psi)| \leq C_{\varphi} \|\psi\|_{\mathcal{H}}.$$

This proves that  $\varphi \in \text{Dom}(A^*) = \text{Dom}(A)$ . Finally we have  $T = A + 1$  and  $A_F = T - 1 = A$ .

### 3.1.7 Relatively bounded perturbations of self-adjoint operators

**Definition 3.36.** Let  $A$  and  $B$  be operators on  $\mathbf{E}$ . We say that  $B$  is  $A$ -bounded if  $\text{Dom}(A) \subset \text{Dom}(B)$  and there exist  $a, b \geq 0$  such that

$$\forall \varphi \in \text{Dom}(A), \quad \|B\varphi\|_{\mathbf{E}} \leq a \|A\varphi\|_{\mathbf{E}} + b \|\varphi\|_{\mathbf{E}}. \quad (3.2)$$

The  $A$ -bound of  $B$  is the infimum of all  $a \geq 0$  for which there exists  $b$  such that (3.2) holds.

*Remark 3.37.*  $B$  is  $A$ -bounded if and only if  $\text{Dom}(A) \subset \text{Dom}(B)$  and  $B$  is a continuous map from  $(\text{Dom}(A), \|\cdot\|_{\text{Dom}(A)})$  to  $\mathbf{E}$ .

*Remark 3.38.* If  $B$  is bounded then it is  $A$  bounded with  $A$ -bound 0 (we can take  $a = 0$  and  $b = \|B\|_{\mathcal{L}(\mathbf{E})}$  in (3.2)).

*Remark 3.39.* The  $A$ -bound of  $B$  is defined as the infimum of all possible  $a$  in (3.2). This infimum is not necessarily attained. In particular,  $B$  can be unbounded but  $A$ -bounded with  $A$ -bound 0. For example, if  $B$  is a symmetric operator on  $\mathcal{H}$  then  $B$  is  $B^2$ -bounded with bound 0. Indeed,

$$\text{Dom}(B^2) = \{\varphi \in \text{Dom}(B) : B\varphi \in \text{Dom}(B)\} \subset \text{Dom}(B)$$

and for  $\varepsilon > 0$  and  $\varphi \in \text{Dom}(B^2)$  we have

$$0 \leq \|(\varepsilon^2 B^2 - 1)\varphi\|^2 = \varepsilon^4 \|B^2\varphi\|^2 + \|\varphi\|^2 - 2\varepsilon^2 \|B\varphi\|^2,$$

so

$$\|B\varphi\|^2 \leq \frac{\varepsilon^2}{2} \|B^2\varphi\|^2 + \frac{\varepsilon^{-2}}{2} \|\varphi\|^2 \leq \frac{1}{4} (\varepsilon \|B^2\varphi\| + \varepsilon^{-1} \|\varphi\|)^2.$$

Thus (3.2) holds with  $a = \varepsilon/4$  and  $b = 1/(4\varepsilon)$  for all  $\varepsilon > 0$  and  $B$  is  $B^2$ -bounded with  $B^2$ -bound 0 (but (3.2) cannot hold with  $a = 0$  if  $B$  is not bounded).

We give examples of operators which are relatively bounded with respect to the usual Laplacian on  $\mathbb{R}^d$ . We denote by  $H_0$  the Laplacian  $-\Delta$  on  $L^2(\mathbb{R}^d)$ , with domain  $H^2(\mathbb{R}^d)$ .

*Example 3.40.* Let  $\beta, V \in L^\infty(\mathbb{R}^d)$  and  $j \in \llbracket 1, d \rrbracket$ . Then  $\beta(x)\partial_j$  and  $V$  are  $H_0$ -bounded with  $H_0$ -bound equal to 0. Indeed for  $u \in H^2(\mathbb{R}^d)$ ,

$$\|\partial_j u\|^2 = \langle \partial_j u, \partial_j u \rangle = \langle -\partial_j^2 u, u \rangle \leq \langle -\Delta u, u \rangle \leq \|H_0 u\| \|u\| \leq \varepsilon \|H_0 u\|^2 + \frac{\|u\|^2}{4\varepsilon}.$$

**Theorem 3.41** (Kato-Rellich). Let  $A$  be a selfadjoint operator on the Hilbert space  $\mathcal{H}$ . Let  $B$  be a symmetric operator on  $\mathcal{H}$ . Assume that  $B$  is  $A$ -bounded with bound smaller than 1.

- (i) The operator  $A + B$ , defined on the domain  $\text{Dom}(A + B) = \text{Dom}(A)$ , is selfadjoint.
- (ii) Let  $\mathcal{D} \subset \text{Dom}(A)$ . If  $A$  is essentially selfadjoint on  $\mathcal{D}$ , then so is  $A + B$ .

*Proof.* The operator  $A + B$  is symmetric as the sum of two symmetric operators. There exist  $a \in [0, 1[$  and  $b \geq 0$  such that (3.2) holds. Let  $\beta > 0$ . We recall that for  $\varphi \in \text{Dom}(A)$  we have

$$\|(A - i\beta)\varphi\|^2 = \|A\varphi\|^2 + \beta^2 \|\varphi\|^2,$$

so

$$\|B\varphi\| \leq a \|A\varphi\| + b \|\varphi\| \leq (a + b\beta^{-1}) \|(A - i\beta)\varphi\|.$$

Let  $\psi \in \mathcal{H}$ . Applied with  $\varphi = (A - i\beta)^{-1}\psi \in \text{Dom}(A)$ , this inequality gives

$$\|B(A - i\beta)^{-1}\psi\| \leq (a + b\beta^{-1}) \|\psi\|.$$

Assume that  $|\beta| > \frac{b}{1-a}$ . Then  $T = B(A - i\beta)^{-1}$  is bounded with bound smaller than 1, so  $(1 + T)$  has a bounded inverse on  $\mathcal{H}$ . We deduce that

$$\text{Ran}(A + B - i\beta) = \text{Ran}((1 + T)(A - i\beta)) = \mathcal{H}.$$

We similarly prove that  $\text{Ran}(A + B + i\beta) = \mathcal{H}$ . By Proposition 3.21, this proves that  $A + B$  is selfadjoint.  $\square$

**Proposition 3.42.** *Assume that  $d \leq 3$ . Let  $V$  be a potential (Borel function) on  $\mathbb{R}^d$ . We assume that we can write  $V = V_2 + V_\infty$  where  $V_2 \in L^2(\mathbb{R}^d)$  and  $V_\infty \in L^\infty(\mathbb{R}^d)$ . Then the Schrödinger operator  $H = H_0 + V$  is selfadjoint on  $L^2(\mathbb{R}^d)$  with domain  $\text{Dom}(H) = \text{Dom}(H_0) = H^2(\mathbb{R}^d)$ .*

*Proof.* Let  $u \in H^2(\mathbb{R}^d)$ . For  $\varepsilon > 0$  we have

$$\begin{aligned} \|u\|_{L^\infty(\mathbb{R}^d)} &\leq \|\hat{u}\|_{L^1(\mathbb{R}^d)} \leq \left\| (1 + \varepsilon^2 |\xi|^2)^{-1} \right\|_{L^2(\mathbb{R}^d)} \left\| (1 + \varepsilon^2 |\xi|^2) \hat{u} \right\|_{L^2(\mathbb{R}^d)} \\ &\leq C_\varepsilon (\varepsilon^2 \|\Delta u\|_{L^2(\mathbb{R}^d)} + \|u\|_{L^2(\mathbb{R}^d)}), \end{aligned}$$

where

$$C_\varepsilon = \sqrt{\int_{\mathbb{R}^d} (1 + \varepsilon^2 |\xi|^2)^{-2} d\xi}.$$

We have  $C_\varepsilon = \varepsilon^{-\frac{d}{2}} C_1$ , so

$$\begin{aligned} \|Vu\|_{L^2} &\leq \|V_2\|_{L^2} \|u\|_{L^\infty} + \|V_\infty\| \|u\|_{L^2} \\ &\leq \varepsilon^{2-\frac{d}{2}} C_1 \|V_2\| \|\Delta u\|_{L^2} + (\varepsilon^{-\frac{d}{2}} C_1 \|V_2\|_{L^2} + \|V_\infty\|_{L^\infty}) \|u\|_{L^2}. \end{aligned}$$

Applied with  $\varepsilon > 0$  small enough this proves that  $V$  is  $H_0$ -bounded with  $H_0$ -bound smaller than 1. We conclude with Theorem 3.41.  $\square$

*Remark 3.43.* We can prove that the same conclusion holds for  $V \in L^p(\mathbb{R}^d) + L^\infty(\mathbb{R}^d)$  for  $p \geq 2$  if  $d = 4$  and  $p \in [2, \frac{2d}{d-4}[$  if  $d \geq 5$ .

*Example 3.44.* Let  $d \leq 3$  and  $\alpha \in [0, \frac{d}{2}[$ . Then for any  $c \in \mathbb{R}$  the operator

$$H = H_0 + \frac{c}{|x|^\alpha}$$

is well-defined and selfadjoint on the domain  $\text{Dom}(H) = H^2(\mathbb{R}^d)$ .



## 3.2 Spectrum of selfadjoint operators

### 3.2.1 Basic properties

**Proposition 3.45.** *Let  $A$  be a selfadjoint operator on  $\mathcal{H}$ . Then  $\sigma(A) \subset \mathbb{R}$  and for  $z \in \rho(A)$  we have*

$$\|(A - z)^{-1}\|_{\mathcal{L}(\mathcal{H})} = \frac{1}{\text{dist}(z, \sigma(A))}. \quad (3.3)$$

In particular,  $\sigma(A) \neq \emptyset$ .

*Proof.* The first statement follows from Proposition 3.21. Let  $z \in \rho(A)$ . By Proposition 2.49 we have

$$((A - z)^{-1})^* = (A^* - \bar{z})^{-1} = (A - \bar{z})^{-1}.$$

Since  $(A - z)^{-1}$  and  $(A - \bar{z})^{-1}$  commute,  $(A - z)^{-1}$  is a bounded normal operator on  $\mathcal{H}$ . Then, by Propositions 1.46 and 1.15,

$$\|(A - z)^{-1}\|_{\mathcal{L}(\mathcal{H})} = \sup_{\mu \in \sigma((A - z)^{-1})} |\mu| = \sup_{\lambda \in \sigma(A)} |\lambda - z|^{-1} = \frac{1}{\inf_{\lambda \in \sigma(A)} |\lambda - z|}.$$

The proposition follows.  $\square$

**Proposition 3.46.** *Let  $A$  be a selfadjoint operator on  $\mathcal{H}$  and  $\lambda \in \mathbb{R}$ .*

(i) *Let  $\varepsilon > 0$ . If there exists  $\varphi \in \text{Dom}(A) \setminus \{0\}$  such that  $\|(A - \lambda)\varphi\|_{\mathcal{H}} \leq \varepsilon \|\varphi\|_{\mathcal{H}}$  then  $\sigma(A) \cap [\lambda - \varepsilon, \lambda + \varepsilon] \neq \emptyset$ .*

(ii)  *$\lambda \in \sigma(A)$  if and only if there exists a sequence  $(\varphi_n)_{n \in \mathbb{N}}$  in  $\text{Dom}(A)$  such that  $\|\varphi_n\|_{\mathcal{H}} = 1$  for all  $n \in \mathbb{N}$  and*

$$\|(A - \lambda)\varphi_n\|_{\mathcal{H}} \xrightarrow{n \rightarrow +\infty} 0.$$

*Such a sequence is called a Weyl sequence.*

*Proof.* • Assume that  $[\rho - \varepsilon, \rho + \varepsilon] \subset \rho(A)$ . Since  $\rho(A)$  is open there exists  $\varepsilon_1 > \varepsilon$  such that  $[\rho - \varepsilon_1, \rho + \varepsilon_1] \subset \rho(A)$ . By Proposition 3.45 we have  $\|(A - \lambda)^{-1}\|_{\mathcal{L}(\mathcal{H})} \leq \varepsilon_1^{-1}$ . Then for  $\varphi \in \text{Dom}(A) \setminus \{0\}$  we have

$$\|\varphi\| \leq \|(A - \lambda)^{-1}\| \|(A - \lambda)\varphi\| \leq \frac{\|(A - \lambda)\varphi\|}{\varepsilon_1},$$


so  $\|(A - \lambda)\varphi\| \geq \varepsilon_1 \|\varphi\| > \varepsilon \|\varphi\|$ . This prove the first statement by contradiction.

• If a Weyl sequence exists then  $\lambda \in \sigma(A)$  by Proposition 2.21 (we can also use the first statement). Now assume that there exists  $c > 0$  such that

$$\forall \varphi \in \text{Dom}(A), \quad \|(A - \lambda)\varphi\|_{\mathcal{H}} \geq c \|\varphi\|_{\mathcal{H}}.$$

Then  $A - \lambda$  is injective with closed range by Proposition 2.34. On the other hand, by Proposition 2.46,

$$\overline{\text{Ran}(A - \lambda)} = \ker((A - \lambda)^*)^\perp = \ker(A - \lambda)^\perp = \mathcal{H}.$$

This proves that  $\lambda \in \rho(A)$ .  $\square$   Ex. 3.6

### 3.2.2 Discrete and essential spectra

**Proposition 3.47.** *Let  $A$  be a selfadjoint operator on  $\mathcal{H}$ . Assume that  $\lambda$  is an isolated element of  $\sigma(A)$ . Let  $\Pi_\lambda$  be the corresponding Riesz projection. Then  $\Pi_\lambda$  is the orthogonal projection on  $\ker(A - \lambda)$ . In particular,  $\lambda$  is an eigenvalue of  $A$  and if  $\dim(\ker(A - \lambda)) < \infty$ , then its geometric and algebraic multiplicities coincide.*

*Proof.* Let  $r > 0$  be so small that  $\sigma(A) \cap D(\lambda, 2r) = \{\lambda\}$ . We have

$$\Pi_\lambda = -\frac{1}{2i\pi} \int_{\mathcal{C}(\lambda, r)} (A - \zeta)^{-1} d\zeta.$$

Then

$$\Pi_\lambda^* = \frac{1}{2i\pi} \int_{\mathcal{C}(\lambda, r)} (A - \bar{\zeta})^{-1} d\zeta = \Pi_\lambda,$$

so  $\Pi_\lambda$  is an orthogonal projection. By Proposition 2.66 we have  $\ker(A - \lambda) \subset \text{Ran}(\Pi_\lambda)$ .

For  $\varphi \in \mathcal{H}$  we have

$$\begin{aligned} (A - \lambda)\Pi\varphi &= -\frac{1}{2i\pi} \int_{\mathcal{C}(\lambda, r)} (A - \lambda)(A - \zeta)^{-1}\varphi d\zeta \\ &= -\frac{1}{2i\pi} \int_{\mathcal{C}(\lambda, r)} (\varphi + (\zeta - \lambda)(A - \zeta)^{-1}\varphi) d\zeta. \end{aligned} \quad (3.4)$$

The map  $\zeta \mapsto (\zeta - \lambda)(A - \zeta)^{-1}$  is analytic in  $D(\lambda, r) \setminus \{\lambda\}$ . By (3.3) it is also bounded. Thus it extends to an analytic function on  $D(\lambda, r)$  and (3.4) vanishes. This proves that  $\text{Ran}(\Pi) \subset \ker(A - \lambda)$ , so  $\text{Ran}(\Pi) = \ker(A - \lambda)$ . Finally,  $\text{Ran}(\Pi)$  cannot be  $\{0\}$  (since  $\lambda$  belongs to the spectrum of the restriction of  $A$  to  $\text{Ran}(\Pi)$ ), so  $\lambda$  is an eigenvalue of  $A$ .  $\square$

**Corollary 3.48.** *Let  $A$  be a selfadjoint operator on  $\mathcal{H}$  and let  $\lambda$  be an isolated element of  $\sigma(A)$ . Let  $\mathbf{G}$  be a reducing subspace for  $A$  and let  $A_{\mathbf{G}}$  be the restriction of  $A$  to  $\mathbf{G}$ . If  $\mathbf{G} \subset \ker(A - \lambda)^\perp$  then  $\sigma(A_{\mathbf{G}}) \subset \sigma(A) \setminus \{\lambda\}$ .*

*Proof.* By Proposition 2.59 we have  $\sigma(A_{\mathbf{G}}) \subset \sigma(A)$ . Moreover,  $A_{\mathbf{G}}$  is a selfadjoint operator by Proposition 3.18 and  $\lambda$  is not an eigenvalue of  $A_{\mathbf{G}}$  since  $\ker(A_{\mathbf{G}} - \lambda) = \ker(A - \lambda) \cap \mathbf{G} = \{0\}$ . By Proposition 3.47,  $\lambda \in \rho(A_{\mathbf{G}})$ .  $\square$

**Lemma 3.49.** *Let  $A$  be a selfadjoint operator on  $\mathcal{H}$ . Let  $\lambda \in \sigma(A)$ . Assume that  $\ker(A - \lambda)$  has finite dimension and that there exists  $c > 0$  such that*

$$\forall \varphi \in \ker(A - \lambda)^\perp, \quad \|(A - \lambda)\varphi\| \geq c \|\varphi\|. \quad (3.5)$$

*Then  $\lambda$  is isolated in  $\sigma(A)$ .*

*Proof.* Let  $\mathbf{F} = \ker(A - \lambda)$  and  $\mathbf{G} = \mathbf{F}^\perp$ . Then  $\mathbf{F}$  and  $\mathbf{G}$  are closed. Let  $\Pi$  be the orthogonal projection on  $\mathbf{F}$ . Let  $A_{\mathbf{F}}$  and  $A_{\mathbf{G}}$  be the restrictions of  $A$  to  $\mathbf{F}$  and  $\mathbf{G}$ . We have  $\sigma(A_{\mathbf{F}}) = \{\lambda\}$ . On the other hand,  $A_{\mathbf{G}}$  is a selfadjoint operator on  $\mathbf{G}$  such that  $\ker(A_{\mathbf{G}} - \lambda) = \{0\}$ . Then  $\overline{\text{Ran}(A_{\mathbf{G}} - \lambda)} = \ker(A_{\mathbf{G}} - \lambda)^\perp = \mathbf{G}$ . By (3.5) and Proposition 2.34,  $\text{Ran}(A_{\mathbf{G}} - \lambda)$  is closed so  $\lambda \in \rho(A_{\mathbf{G}})$ . Since  $\rho(A_{\mathbf{G}})$  is open, there exists  $\varepsilon > 0$  such that  $] \lambda - \varepsilon, \lambda + \varepsilon [ \subset \rho(A_{\mathbf{G}})$ . Then, by Proposition 2.59,  $] \lambda - \varepsilon, \lambda + \varepsilon [ \setminus \{\lambda\} \subset \rho(A_{\mathbf{F}}) \cap \rho(A_{\mathbf{G}}) = \rho(A)$ .  $\square$

**Proposition 3.50 (Weyl Criterion).** *Let  $A$  be a selfadjoint operator on  $\mathcal{H}$  and  $\lambda \in \mathbb{R}$ . The following assertions are equivalent.*

- (i)  $\lambda \in \sigma_{\text{ess}}(A)$ .
- (ii) *There exists a sequence  $(\varphi_n)_{n \in \mathbb{N}}$  in  $\text{Dom}(A)$  such that  $\|\varphi_n\|_{\mathcal{H}} = 1$  for all  $n \in \mathbb{N}$ ,  $\varphi_n$  goes weakly to 0 and  $\|(A - \lambda)\varphi_n\|_{\mathcal{H}} \rightarrow 0$  as  $n \rightarrow \infty$ .*
- (iii) *There exists a sequence  $(\varphi_n)_{n \in \mathbb{N}}$  in  $\text{Dom}(A)$  such that  $\|\varphi_n\|_{\mathcal{H}} = 1$  for all  $n \in \mathbb{N}$ ,  $(\varphi_n)_{n \in \mathbb{N}}$  has no convergent subsequence in  $\mathcal{H}$  and  $\|(A - \lambda)\varphi_n\|_{\mathcal{H}} \rightarrow 0$  as  $n \rightarrow \infty$ .*

$\circlearrowleft$  Ex. 3.7

*Proof.* We set  $\mathbf{F} = \ker(A - \lambda)$  and  $\mathbf{G} = \ker(A - \lambda)^\perp$ . We denote by  $A_{\mathbf{G}}$  the restriction of  $A$  to  $\mathbf{G}$ .

- Assume that  $\lambda \in \sigma_{\text{ess}}(A)$ . If  $\dim(\mathbf{F}) = \infty$  then we can construct an orthonormal sequence  $(\varphi_n)_{n \in \mathbb{N}}$  in  $\mathbf{F}$ . Now assume that  $\dim(\mathbf{F}) < \infty$ . By Lemma 3.49, (3.5) cannot hold, so there exists a normalized sequence  $(\varphi_n)_{n \in \mathbb{N}}$  in  $\mathbf{G}$  such that  $\|(A - \lambda)\varphi_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . For  $\psi \in \mathbf{F}$

we have  $\langle \psi, \varphi_n \rangle = 0$  for all  $n \in \mathbb{N}$ . It remains to prove that  $\langle \psi, \varphi_n \rangle \rightarrow 0$  for all  $\psi \in \mathbf{G}$ . It is enough to prove this for  $\psi$  in a dense subset of  $\mathbf{G}$ . We have

$$\overline{\text{Ran}(A_{\mathbf{G}} - \lambda)}^\perp = \ker(A_{\mathbf{G}} - \lambda) = \{0\},$$

so it is enough to consider  $\psi \in \text{Ran}(A_{\mathbf{G}} - \lambda)$ . In this case we consider  $\zeta \in \text{Dom}(A_{\mathbf{G}})$  such that  $\psi = (A_{\mathbf{G}} - \lambda)\zeta$  and write

$$\langle \psi, \varphi_n \rangle = \langle (A_{\mathbf{G}} - \lambda)\zeta, \varphi_n \rangle = \langle \zeta, (A_{\mathbf{G}} - \lambda)\varphi_n \rangle \xrightarrow{n \rightarrow +\infty} 0.$$

This proves that  $\varphi_n \rightarrow 0$  as  $n \rightarrow \infty$ . This proves (i)  $\implies$  (iii).

- A normalized sequence which goes weakly to 0 cannot have a convergent subsequence, so (ii)  $\implies$  (iii).
- Assume that there exists a sequence  $(\varphi_n)_{n \in \mathbb{N}}$  as in (iii). By Proposition 3.46 we have  $\lambda \in \sigma(A)$ . Assume by contradiction that  $\lambda \in \sigma_{\text{disc}}(A)$ . For  $n \in \mathbb{N}$  we write  $\varphi_n = \psi_n + \psi_n^\perp$  where  $\psi_n \in F$  and  $\psi_n^\perp \in \mathbf{G} \cap \text{Dom}(A)$ . We have

$$(A_{\mathbf{G}} - \lambda)\psi_n^\perp = (A - \lambda)\psi_n^\perp = (A - \lambda)\varphi_n \xrightarrow{n \rightarrow \infty} 0.$$

Since  $\lambda \in \rho(A_{\mathbf{G}})$  by Corollary 3.48, we deduce that  $\psi_n^\perp \rightarrow 0$  as  $n \rightarrow \infty$ . In particular,  $\|\varphi_n - \psi_n\|_{\mathcal{H}} \rightarrow 0$ . But the sequence  $(\psi_n)_{n \in \mathbb{N}}$  is in  $F$  which has finite dimension, so it has a convergent subsequence. This gives a contradiction and proves that  $\lambda \in \sigma_{\text{ess}}(A)$ .  $\square$

**Proposition 3.51.** *Let  $A$  be a selfadjoint operator on  $\mathcal{H}$  and  $\lambda \in \sigma_{\text{ess}}(A)$ . Let  $N \in \mathbb{N}^*$  and  $\varepsilon > 0$ . There exists an orthonormal family  $(\varphi_n)_{1 \leq n \leq N}$  such that*

$$\forall n \in \llbracket 1, N \rrbracket, \quad \|(A - \lambda)\varphi_n\|_{\mathcal{H}} \leq \varepsilon.$$

*Proof.* • If  $\lambda$  is isolated, it is an eigenvalue of infinite multiplicity, so we can consider an orthonormal family  $(\varphi_n)_{1 \leq n \leq N}$  in  $\ker(A - \lambda)$ .

- Now assume that  $\lambda$  is not isolated. We fix distinct elements  $\lambda_1, \dots, \lambda_N$  of  $\sigma(A)$  such that, for all  $n \in \llbracket 1, N \rrbracket$ ,

$$|\lambda_n - \lambda| \leq \frac{\varepsilon}{2}. \quad (3.6)$$

Let  $\eta \in ]0, 1]$ . Let  $n \in \llbracket 1, N \rrbracket$ . By Proposition 3.50 we can consider  $\psi_n \in \text{Dom}(A)$  such that  $\|\psi_n\|_{\mathcal{H}} = 1$  and

$$\|(A - \lambda_n)\psi_n\|_{\mathcal{H}} \leq \eta.$$

We set  $\tilde{\varphi}_1 = \psi_1$  and for  $n \in \llbracket 2, N \rrbracket$  we define by induction

$$\tilde{\varphi}_n = \psi_n - \sum_{k=1}^{n-1} \langle \tilde{\varphi}_k, \psi_n \rangle_{\mathcal{H}} \tilde{\varphi}_k.$$

- We prove by induction on  $n \in \llbracket 1, N \rrbracket$  that there exists a constant  $C_n > 0$  independent of  $\eta \in ]0, 1]$  such that

$$\|(A - \lambda_n)\tilde{\varphi}_n\| \leq C_n \eta \quad \text{and} \quad \left| \|\tilde{\varphi}_n\| - 1 \right| \leq C_n \eta. \quad (3.7)$$

This is clear for  $n = 1$ . Now assume that this holds up to order  $n - 1$  for some  $n \in \llbracket 2, N \rrbracket$ . For  $k \in \llbracket 1, n - 1 \rrbracket$  we have

$$(\lambda_n - \lambda_k) \langle \tilde{\varphi}_k, \psi_n \rangle = \langle (A - \lambda_k)\tilde{\varphi}_k, \psi_n \rangle - \langle \tilde{\varphi}_k, (A - \lambda_n)\psi_n \rangle,$$

so, for some  $\tilde{C}_{k,n} > 0$ ,

$$|\langle \tilde{\varphi}_k, \psi_n \rangle| \leq \frac{C_k \eta + (1 + C_k \eta) \eta}{|\lambda_k - \lambda_n|} \leq \tilde{C}_{k,n} \eta.$$

Then

$$\left| \|\tilde{\varphi}_n\| - 1 \right| \leq \|\tilde{\varphi}_n - \psi_n\| \leq \sum_{k=1}^{n-1} |\langle \tilde{\varphi}_k, \psi_n \rangle| \|\tilde{\varphi}_k\|$$

and

$$\|(A - \lambda_n)\tilde{\varphi}_n\| \leq \|(A - \lambda_n)\psi_n\| + \sum_{k=1}^{n-1} |\langle \tilde{\varphi}_k, \psi_n \rangle| (\|(A - \lambda_k)\tilde{\varphi}_k\| + |\lambda_k - \lambda_n| \|\tilde{\varphi}_k\|).$$

We deduce (3.7). If  $\eta$  is chosen small enough then for  $n \in \llbracket 1, N \rrbracket$  we can set

$$\varphi_n = \frac{\tilde{\varphi}_n}{\|\tilde{\varphi}_n\|}.$$

Then there exists  $C > 0$  such that for  $n \in \llbracket 1, N \rrbracket$  and  $\eta \in ]0, 1]$  we have

$$\|(A - \lambda_n)\varphi_n\| \leq C\eta.$$

It remains to chose  $\eta$  smaller that  $\varepsilon/(2C)$  and conclude with (3.6).  $\square$

### 3.2.3 Min-max principle

We consider on  $\mathcal{H}$  a self-adjoint operator  $A$  bounded from below.

**Proposition 3.52.** *We have*

$$\min(\sigma(A)) = \inf_{\varphi \in \text{Dom}(A) \setminus \{0\}} \frac{\langle A\varphi, \varphi \rangle_{\mathcal{H}}}{\|\varphi\|_{\mathcal{H}}^2}. \quad (3.8)$$

*Proof.* We denote by  $\mu_1$  the right-hand side of (3.8).

• Let  $\lambda \in \sigma(A)$ . By the Weyl criterion (Proposition 3.50) there exists a sequence  $(\varphi_n)$  such that  $\|\varphi_n\| = 1$  for all  $n$  and  $\|(A - \lambda)\varphi_n\| \rightarrow 0$ . This implies in particular

$$\mu_1 \leq \langle A\varphi_n, \varphi_n \rangle \xrightarrow{n \rightarrow \infty} \lambda,$$

so  $\mu_1 \leq \min(\sigma(A))$ .

• Now assume by contradiction that  $\mu_1 \in \rho(A)$ . We set  $R = (A - \mu_1)^{-1}$ . For  $\eta, \psi \in \mathcal{H}$  we set

$$\mathfrak{q}(\eta, \psi) = \langle R\eta, \psi \rangle_{\mathcal{H}}.$$

For  $\eta \in \mathcal{H}$  and  $\psi = R\eta \in \text{Dom}(A)$  we have

$$\mathfrak{q}(\eta, \eta) = \langle \psi, (A - \mu_1)\psi \rangle \geq 0,$$

so  $\mathfrak{q}$  is a non-negative sesquilinear form on  $\mathcal{H}$ . Let  $(\psi_n)$  be a sequence in  $\text{Dom}(A)$  such that  $\|\psi_n\|_{\mathcal{H}} = 1$  for all  $n \in \mathbb{N}$  and

$$\langle A\psi_n, \psi_n \rangle \xrightarrow{n \rightarrow +\infty} \mu_1.$$

For  $n \in \mathbb{N}$  we set  $\eta_n = (A - \mu_1)\psi_n$ . Then by the Cauchy-Schwarz inequality we have

$$\begin{aligned} 1 &= \|\psi_n\|_{\mathcal{H}}^2 = \mathfrak{q}(\eta_n, \psi_n) \\ &\leq \mathfrak{q}(\eta_n, \eta_n)^{\frac{1}{2}} \mathfrak{q}(\psi_n, \psi_n)^{\frac{1}{2}} \\ &= \langle \psi_n, (A - \mu_1)\psi_n \rangle^{\frac{1}{2}} \langle R\psi_n, \psi_n \rangle^{\frac{1}{2}} \\ &\xrightarrow{n \rightarrow +\infty} 0. \end{aligned}$$

This gives a contradiction and proves that  $\mu_1(A) \in \sigma(A)$ , and in particular  $\mu_1(A) \geq \min(\sigma(A))$ . The conclusion follows.  $\square$

**Theorem 3.53** (Min-max Theorem). *Let  $A$  be a lower-bounded self-adjoint operator on  $\mathcal{H}$ . We denote by  $(\lambda_k)_{k \in \mathbb{N}^*, k \leq N}$  with  $N \in \mathbb{N} \cup \{\infty\}$  the non-decreasing sequence of eigenvalues (counted with multiplicities) smaller than  $\inf \sigma_{\text{ess}}(A)$ . For  $n \in \mathbb{N}^*$  (with  $n \leq \dim(\mathcal{H})$  if  $\mathcal{H}$  is of finite dimension) we have*

$$\inf_{\substack{F \subset \text{Dom}(A) \\ \dim(F) = n}} \sup_{\varphi \in F \setminus \{0\}} \frac{\langle A\varphi, \varphi \rangle_{\mathcal{H}}}{\|\varphi\|_{\mathcal{H}}^2} = \begin{cases} \lambda_n & \text{if } n \leq N, \\ \inf \sigma_{\text{ess}}(A) & \text{if } n > N. \end{cases}$$

*Proof.* For  $n \in \mathbb{N}^*$  we set

$$\mu_n = \inf_{\substack{F \subset \text{Dom}(A) \\ \dim(F)=n}} \sup_{\varphi \in F \setminus \{0\}} \frac{\langle A\varphi, \varphi \rangle_{\mathcal{H}}}{\|\varphi\|_{\mathcal{H}}^2}.$$

- We set  $\mathcal{N} = \llbracket 1, N \rrbracket$  if  $N \in \mathbb{N}$  and  $\mathcal{N} = \mathbb{N}$  if  $N = +\infty$ . We consider an orthonormal family  $(\varphi_k)_{k \in \mathcal{N}}$  such that  $\varphi_k \in \text{Dom}(A)$  and  $A\varphi_k = \lambda_k \varphi_k$  for all  $k \in \mathcal{N}$ . For  $n \in \mathcal{N}$  we set  $F_n = \text{span}(\varphi_1, \dots, \varphi_n)$ . We also set  $\eta = \inf \sigma_{\text{ess}}(A)$ .
- Let  $n \in \mathcal{N}$ . Let  $\varphi \in F_n$  such that  $\|\varphi\| = 1$ . We can write  $\varphi = \sum_{k=1}^n \alpha_k \varphi_k$  with  $\sum_{k=1}^n |\alpha_k|^2 = 1$ . Then we have

$$\langle A\varphi, \varphi \rangle = \sum_{k=1}^n |\alpha_k|^2 \lambda_k \leq \lambda_n,$$

so  $\mu_n \leq \lambda_n$ .

- Let  $F$  be a subspace of  $\text{Dom}(A)$  of dimension  $n$ . By Corollary 3.48, the restriction of  $A$  to  $F_{n-1}^\perp$  is selfadjoint and its spectrum is included in  $[\lambda_n, +\infty[$ . There exists  $\varphi \in F \cap F_{n-1}^\perp$  with  $\|\varphi\| = 1$ . For such a  $\varphi$  we have  $\langle A\varphi, \varphi \rangle \geq \lambda_n$  by Proposition 3.52. This proves that  $\mu_n \geq \lambda_n$ . Then  $\mu_n = \lambda_n$  and the infimum is a minimum.
- Now assume that  $N$  is finite and consider  $n > N$ . As in the previous step, we see that  $\mu_n \geq \eta$ . Then let  $\varepsilon > 0$ . Since  $\eta \in \sigma_{\text{ess}}(A)$  there exists by Proposition 3.51 an orthonormal family  $(\psi_k)_{1 \leq k \leq n}$  of vectors in  $\text{Dom}(A)$  such that

$$\forall k \in \llbracket 1, n \rrbracket, \quad \|\psi_k\|_{\mathcal{H}} = 1 \quad \text{and} \quad \|(A - \eta)\psi_k\|_{\mathcal{H}} \leq \frac{\varepsilon}{\sqrt{n}}.$$

Let  $\psi \in F = \text{span}(\psi_1, \dots, \psi_n)$  such that  $\|\psi\| = 1$ . We write  $\psi = \sum_{k=1}^n \alpha_k \psi_k$  with  $\sum_{k=1}^n |\alpha_k|^2 = 1$ . Then we have

$$\begin{aligned} \langle A\psi, \psi \rangle &\leq \eta + \|(A - \eta)\psi\| \\ &\leq \eta + \sum_{k=1}^n |\alpha_k| \|(A - \eta)\psi_k\|_{\mathcal{H}} \\ &\leq \eta + \left( \sum_{k=1}^n \|(A - \eta)\psi_k\|^2 \right)^{\frac{1}{2}} \\ &\leq \eta + \varepsilon. \end{aligned}$$

This proves that

$$\mu_n \leq \max_{\substack{\psi \in F \\ \|\psi\|_{\mathcal{H}}=1}} \langle A\psi, \psi \rangle_{\mathcal{H}} \leq \eta + \varepsilon.$$

Finally  $\mu_n = \eta$ . □

*Remark 3.54.* • Let  $F$  be a finite dimensional subspace of  $\text{Dom}(A)$ . Since the unit sphere  $\mathbb{S}_F$  of  $F$  is compact and the map  $\varphi \mapsto \langle A\varphi, \varphi \rangle$  is continuous on  $\mathbb{S}_F$ , we have

$$\sup_{\varphi \in F \setminus \{0\}} \frac{\langle A\varphi, \varphi \rangle_{\mathcal{H}}}{\|\varphi\|_{\mathcal{H}}^2} = \sup_{\varphi \in \mathbb{S}_F} \langle A\varphi, \varphi \rangle_{\mathcal{H}} = \max_{\varphi \in \mathbb{S}_F} \langle A\varphi, \varphi \rangle_{\mathcal{H}} = \max_{\varphi \in F \setminus \{0\}} \frac{\langle A\varphi, \varphi \rangle_{\mathcal{H}}}{\|\varphi\|_{\mathcal{H}}^2}.$$

- Let  $n \in \mathcal{N}$ . We have seen that

$$\inf_{\substack{F \subset \text{Dom}(A) \\ \dim(F)=n}} \sup_{\varphi \in F \setminus \{0\}} \frac{\langle A\varphi, \varphi \rangle_{\mathcal{H}}}{\|\varphi\|_{\mathcal{H}}^2} = \lambda_n = \sup_{\varphi \in F_n \setminus \{0\}} \frac{\langle A\varphi, \varphi \rangle_{\mathcal{H}}}{\|\varphi\|_{\mathcal{H}}^2},$$

so the infimum is a minimum.

- When  $n > N$ , the infimum is not necessarily reached. Consider for instance the usual Laplacian  $H_0$  on  $\mathbb{R}^d$ . We have  $\min \sigma(H_0) = \sigma_{\text{ess}}(H_0) = 0$  and there is no  $\varphi \in H^2(\mathbb{R}^2)$  such that  $\langle H_0\varphi, \varphi \rangle$  is equal to 0.

This Min-max Theorem has an equivalent Max-min version. See Exercise 3.8.

☞ Ex. 3.8

**Corollary 3.55.** *Let  $a < \inf \sigma_{\text{ess}}(A)$ . Assume that there exists a subspace  $V$  of  $\text{Dom}(A)$  of dimension  $n \in \mathbb{N}^*$  such that*

$$\forall \varphi \in V, \quad \langle A\varphi, \varphi \rangle_{\mathcal{H}} \leq a \|\varphi\|_{\mathcal{H}}^2.$$

*Then  $A$  has at least  $n$  eigenvalues (counted with multiplicities) not greater than  $a$ .*

**Proposition 3.56.** *Let  $A$  be a lower bounded selfadjoint operator  $A$  on  $\mathcal{H}$ . Let  $\mathfrak{q}_A$  be the corresponding quadratic form and let  $\mathcal{V}_A$  be the form domain of  $A$  (see Definition 3.29).*

(i) *We have*

$$\min \sigma(A) = \inf_{\varphi \in \mathcal{V}_A \setminus \{0\}} \frac{\mathfrak{q}_A(\varphi)}{\|\varphi\|_{\mathcal{H}}^2}. \quad (3.9)$$

(ii) *The right-hand side of (3.9) is a minimum if and only if  $\min \sigma(A)$  is an eigenvalue, and in this case the minimizers are the eigenvectors corresponding to the eigenvalue  $\min \sigma(A)$ .*

*Proof.* • We set

$$\mu_1 = \min \sigma(A) = \inf_{\varphi \in \text{Dom}(A) \setminus \{0\}} \frac{\langle A\varphi, \varphi \rangle}{\|\varphi\|^2} \quad \text{and} \quad \tilde{\mu}_1 = \inf_{\varphi \in \mathcal{V}_A \setminus \{0\}} \frac{\mathfrak{q}_A(\varphi)}{\|\varphi\|_{\mathcal{H}}^2}.$$

Since  $\text{Dom}(A) \subset \mathcal{V}_A$  and  $\mathfrak{q}_A(\varphi) = \langle A\varphi, \varphi \rangle$  for  $\varphi \in \text{Dom}(A)$ , we have  $\tilde{\mu}_1 \leq \mu_1$ . After translation we can assume that  $\mu_1 > 0$ . Then by definition of the form domain,  $\text{Dom}(A)$  is dense in  $\mathcal{V}_A$  for the norm defined by  $\mathfrak{q}_A$ , so we also have  $\mu_1 \leq \tilde{\mu}_1$ . This gives the first statement.

• Now assume that  $\mu_1$  is an eigenvalue of  $A$ . Then for a corresponding eigenvector  $\varphi$  we have

$$\frac{\mathfrak{q}_A(\varphi)}{\|\varphi\|^2} = \frac{\langle A\varphi, \varphi \rangle}{\|\varphi\|^2} = \mu_1,$$

so  $\tilde{\mu}_1$  is a minimum and  $\varphi$  is a minimizer. Conversely, assume that  $\varphi$  is a minimizer for  $\tilde{\mu}_1$  with  $\|\varphi\| = 1$ . Let  $\psi \in \text{Dom}(A)$ . The map

$$\Phi : t \mapsto \frac{\mathfrak{q}_A(\varphi + t\psi)}{\|\varphi + t\psi\|_{\mathcal{H}}^2}$$

is well defined for  $|t|$  small enough, it is smooth and it reaches its minimum at  $t = 0$ . Thus  $\Phi'(0) = 0$ , which implies that

$$\text{Re } \mathfrak{q}_A(\varphi, \psi) = \tilde{\mu}_1 \text{Re } \langle \varphi, \psi \rangle.$$

Since we can replace  $\psi$  by  $i\psi$ , this gives

$$\forall \psi \in \text{Dom}(A), \quad \mathfrak{q}_A(\varphi, \psi) = \langle \tilde{\mu}_1 \varphi, \psi \rangle.$$

This proves that  $\varphi \in \text{Dom}(A)$  and  $A\varphi = \tilde{\mu}_1 \varphi$ . Then  $\tilde{\mu}_1$  is an eigenvalue of  $A$  and  $\varphi$  is a corresponding eigenvector.  $\square$

*Example 3.57.* Let  $\Omega$  be a bounded open set of  $\mathbb{R}^d$ . We denote by  $H_0$  the Dirichlet Laplacian on  $\Omega$  ( $H_0 = -\Delta$ ,  $\text{Dom}(H_0) = H^2(\Omega) \cap H_0^1(\Omega)$ ). The form domain of  $H_0$  is  $H_0^1(\Omega)$  and the corresponding quadratic form is  $\mathfrak{q}_{H_0} : u \mapsto \|\nabla u\|_{L^2(\Omega)}^2$ . We will see in Chapter 4 that  $H_0$  has no essential spectrum. Then by Proposition 3.56 the first eigenvalue of  $H_0$  is given by

$$\lambda_1(H_0) = \inf_{u \in H_0^1(\Omega) \setminus \{0\}} \frac{\|\nabla u\|_{L^2(\Omega)}^2}{\|u\|_{L^2(\Omega)}^2}.$$

By the Poincaré inequality we have  $\lambda_1(H_0) > 0$ .

### 3.3 Exercises

**Exercise 3.1.** Let  $\Omega$  be an open subset of  $\mathbb{R}^d$ . We consider on  $L^2(\Omega)$  the operators  $H_0$  and  $H$  which act as  $-\Delta$  on the domains  $\text{Dom}(H_0) = C_0^\infty(\Omega)$  and  $\text{Dom}(H) = H^2(\Omega)$ . Are  $H_0$  and  $H$  symmetric operators ?

**Exercise 3.2.** Let  $\mathcal{H}_1$  and  $\mathcal{H}_2$  be two Hilbert spaces. Let  $U : \mathcal{H}_1 \rightarrow \mathcal{H}_2$  be a unitary operator. Let  $A_1$  be an operator on  $\mathcal{H}_1$  and  $A_2$  an operator on  $\mathcal{H}_2$ . Assume that  $\text{Dom}(A_2) = U\text{Dom}(A_1)$  and  $A_2 = UA_1U^*$ . Prove that  $A_1$  is selfadjoint on  $\mathcal{H}_1$  if and only if  $A_2$  is selfadjoint on  $\mathcal{H}_2$ .

**Exercise 3.3.** Let  $A$  be a symmetric operator on the Hilbert space  $\mathcal{H}$ . Assume that  $A$  is not selfadjoint but  $\text{Ran}(A - i) = \mathcal{H}$  or  $\text{Ran}(A + i) = \mathcal{H}$ . Prove that  $A$  has no selfadjoint extension.

**Exercise 3.4.** Let  $m > 0$ . We consider the Hilbert space  $\mathcal{H} = H^1(\mathbb{R}^d) \times L^2(\mathbb{R}^d)$  the operator

$$\mathcal{W} = \begin{pmatrix} 0 & 1 \\ \Delta - m & 0 \end{pmatrix}$$

defined on the domain  $\text{Dom}(\mathcal{W}) = H^2(\mathbb{R}^d) \times H^1(\mathbb{R}^d)$ . Prove that  $\mathcal{W}$  is skew-adjoint if  $\mathcal{H}$  is endowed with the Hilbert structure corresponding to the norm defined by

$$\|(u, v)\|_{\mathcal{H}}^2 = \|\nabla u\|_{L^2(\mathbb{R}^d)}^2 + m \|u\|_{L^2(\mathbb{R}^d)}^2 + \|v\|_{L^2(\mathbb{R}^d)}^2.$$

**Exercise 3.5.** Let  $A_0$  be the operator of Example 3.30.

1. What is the adjoint of  $A_0$  ?
2. Compute  $\ker(A_0^* - z)$  for  $z \in \mathbb{C} \setminus \mathbb{R}$ .
3. For  $u \in H^2(0, 1)$  we set

$$Bu = \begin{pmatrix} u(0) \\ u'(0) \\ u(1) \\ u'(1) \end{pmatrix}.$$

Prove that there exists a matrix  $M \in M_4(\mathbb{C})$  (to be explicated) such that an operator  $A$  is a selfadjoint extension of  $A_0$  if and only if there exists a subspace  $F$  of  $\mathbb{C}^4$  such that  $MF = F^\perp$  and

$$A = -\frac{d^2}{dx^2}, \quad \text{Dom}(A) = \{u \in H^2(0, 1) : Bu \in F\}.$$

4. Give some examples of selfadjoint extensions of  $A_0$  ?

**Exercise 3.6.** Give an example of an operator  $A$  and  $\lambda \in \mathbb{C}$  such that  $\lambda \in \sigma(A)$  but there is no corresponding Weyl sequence.

**Exercise 3.7.** We consider the Laplacian  $H = -\Delta$  on  $L^2(\mathbb{R})$ , with domain  $H^2(\mathbb{R})$ . Let  $\lambda > 0$ . Construct a sequence  $(\varphi_n)$  in  $H^2(\mathbb{R})$  such that  $\|\varphi_n\| = 1$ ,  $\|(H - \lambda)\varphi_n\| \rightarrow 0$  and  $\varphi_n$  goes weakly to 0 in  $L^2(\mathbb{R})$ .

**Exercise 3.8.** Prove the following version of the Min-Max Theorem. Let  $A$  be a self-adjoint operator on  $\mathcal{H}$ . Assume that  $A$  is semi-bounded from below. For  $n \in \mathbb{N}^*$  (with  $n \leq \dim(\mathcal{H})$  if  $\mathcal{H}$  is of finite dimension) we set

$$\mu_n(A) = \sup_{\varphi_1, \dots, \varphi_{n-1} \in \mathcal{H}} \inf_{\substack{\varphi \in \text{span}(\varphi_1, \dots, \varphi_{n-1})^\perp \\ \varphi \in \text{Dom}(A) \setminus \{0\}}} \frac{\langle A\varphi, \varphi \rangle_{\mathcal{H}}}{\|\varphi\|_{\mathcal{H}}^2}.$$

The sequence  $(\mu_n)_{n \in \mathbb{N}^*}$  is non-decreasing and for  $n \in \mathbb{N}^*$  one of the following statements hold.

- (i)  $\mu_n(A) < \inf \sigma_{\text{ess}}(A)$  and  $\mu_n$  is the  $n$ -th eigenvalue of  $A$  counted with multiplicities,
- (ii)  $\mu_n(A) = \inf \sigma_{\text{ess}}(A)$ .

