

Symmetry and monotonicity of least energy solutions

Jaeyoung BYEON*, Louis JEANJEAN† and Mihai MARIȘ‡

Abstract

We give a simple proof of the fact that for a large class of quasilinear elliptic equations and systems the solutions that minimize the corresponding energy in the set of all solutions are radially symmetric. We require just continuous nonlinearities and no cooperative conditions for systems. Thus, in particular, our results cannot be obtained by using the moving planes method. In the case of scalar equations, we also prove that any least energy solution has a constant sign and is monotone with respect to the radial variable. Our proofs rely on results in [15, 6] and answer questions from [3, 12].

1 Introduction

We consider the system of partial differential equations

$$-\operatorname{div}(|\nabla u_i|^{p-2} \nabla u_i) = g_i(u), \quad i = 1, \dots, m, \quad (1)$$

where $u = (u_1, \dots, u_m) : \mathbf{R}^N \rightarrow \mathbf{R}^m$, $1 < p \leq N$, $|(y_1, \dots, y_N)|^p = \left(\sum_{j=1}^N y_j^2\right)^{\frac{p}{2}}$, $g_i(0) = 0$ and there exists $G \in C^1(\mathbf{R}^m \setminus \{0\}, \mathbf{R}) \cap C(\mathbf{R}^m, \mathbf{R})$ such that $g_i(u) = \frac{\partial G}{\partial u_i}(u)$ for $u \neq 0$.

Formally, solutions of (1) are critical points of the following energy functional

$$S(u) = \frac{1}{p} \int_{\mathbf{R}^N} \sum_{i=1}^m |\nabla u_i|^p dx - \int_{\mathbf{R}^N} G(u) dx.$$

The aim of this note is to prove, under general assumptions, that those solutions of (1) which minimize the energy S in the set of all solutions are radially symmetric (up to a translation in \mathbf{R}^N). In the scalar case we also study the sign and monotonicity of these solutions. We do not consider here the problem of existence of solutions (respectively of least energy solutions) for (1). We believe that our results cover all situations where the existence of a least energy solution is already known in the literature.

We begin with some definitions. Let Π be an affine hyperplane in \mathbf{R}^N , let Π^+ and Π^- be the two closed half-spaces determined by Π and s_Π the symmetry with respect

*Department of Mathematics and PMI, Pohang University of Science and Technology, Pohang, Kyungbuk 790-784, Republic of Korea, e-mail: jbyeon@postech.ac.kr

†Département de Mathématiques UMR 6623, Université de Franche-Comté, 16, Route de Gray, 25030 Besançon, France, e-mail: louis.jeanjean@univ-fcomte.fr

‡Département de Mathématiques UMR 6623, Université de Franche-Comté, 16, Route de Gray, 25030 Besançon, France, e-mail: mihai.maris@univ-fcomte.fr

to Π (i.e. $s_\Pi(x) = 2p_\Pi(x) - x$, where p_Π is the orthogonal projection onto Π). Given a function f defined on \mathbf{R}^N , we define

$$f_{\Pi^+}(x) = \begin{cases} f(x) & \text{if } x \in \Pi^+ \\ f(s_\Pi(x)) & \text{if } x \in \Pi^- \end{cases}, \quad f_{\Pi^-}(x) = \begin{cases} f(x) & \text{if } x \in \Pi^- \\ f(s_\Pi(x)) & \text{if } x \in \Pi^+. \end{cases} \quad (2)$$

For $\sigma > 0$, we denote $f_\sigma(x) = f(\frac{x}{\sigma})$. We say that a space \mathcal{X} of functions defined on \mathbf{R}^N is admissible if \mathcal{X} is nonempty and

- (i) $\mathcal{X} \subset L^1_{loc}(\mathbf{R}^N, \mathbf{R}^m)$ and $\text{measure}(\{x \mid |u(x)| > \alpha\}) < \infty$ for any $u \in \mathcal{X}$ and $\alpha > 0$;
- (ii) $g_i(u) \in L^1_{loc}(\mathbf{R}^N)$ for any $u \in \mathcal{X}$ and $i = 1, \dots, m$;
- (iii) $\sum_{i=1}^m |\nabla u_i|^p$ and $G(u)$ belong to $L^1(\mathbf{R}^N)$ if $u \in \mathcal{X}$;
- (iv) $u_\sigma \in \mathcal{X}$ for any $u \in \mathcal{X}$ and $\sigma > 0$;
- (v) $u_{\Pi^+}, u_{\Pi^-} \in \mathcal{X}$ whenever $u \in \mathcal{X}$ and Π is an affine hyperplane in \mathbf{R}^N .

Let \mathcal{X} be an admissible function space. We note that from (i) and (iii), $G(0) = 0$. A function $u \in \mathcal{X}$ is a solution of (1) if it satisfies (1) in $\mathcal{D}'(\mathbf{R}^N)$. If (1) admits solutions in \mathcal{X} , we say that \underline{u} is a *least energy solution* if \underline{u} is a nontrivial solution of (1) and

$$S(\underline{u}) = \inf\{S(u) \mid u \in \mathcal{X} \setminus \{0\}, u \text{ is a solution of (1)}\}.$$

We introduce the functionals

$$J(u) = \frac{1}{p} \int_{\mathbf{R}^N} \sum_{i=1}^m |\nabla u_i|^p dx \quad \text{and} \quad V(u) = \int_{\mathbf{R}^N} G(u) dx.$$

Clearly, these functionals are well-defined on any admissible function space. As we will see, the least energy solutions of (1) come from the following minimization problem:

$$\text{minimize } J(u) \text{ in the set } \{u \in \mathcal{X} \mid V(u) = \lambda\}. \quad (\mathcal{P}_\lambda)$$

We shall prove that under some general conditions (see **(C1)**-**(C3)** or **(D1)**-**(D3)** below), all least energy solutions of (1) in the set \mathcal{X} are radially symmetric, up to a translation in \mathbf{R}^N .

It is easy to see that $J(u_\sigma) = \sigma^{N-p} J(u)$ and $V(u_\sigma) = \sigma^N V(u)$. If $V(u) > 0$ for some $u \in \mathcal{X}$, we have $V(u_\sigma) = 1$ for $\sigma = V(u)^{-\frac{1}{N}}$. Then, denoting

$$T = \inf \{J(u) \mid u \in \mathcal{X} \text{ and } V(u) = 1\},$$

we see that

$$J(v) \geq T (V(v))^{\frac{N-p}{N}} \quad \text{for any } v \in \mathcal{X} \text{ satisfying } V(v) > 0. \quad (3)$$

It is clear that u is a minimizer for problem (\mathcal{P}_λ) above ($\lambda > 0$) if and only if u_{σ_1} is a minimizer for (\mathcal{P}_1) , where $\sigma_1 = \lambda^{-\frac{1}{N}}$.

We assume first that $1 < p < N$ and the following conditions are satisfied.

(C1) $T > 0$ and problem (\mathcal{P}_1) has a minimizer $u_* \in \mathcal{X}$;

(C2) Any minimizer $u \in \mathcal{X}$ of (\mathcal{P}_1) is a C^1 function and satisfies the Euler-Lagrange system of equations

$$-\operatorname{div}(|\nabla u_i|^{p-2} \nabla u_i) = \alpha g_i(u) \quad \text{in } \mathcal{D}'(\mathbf{R}^N) \quad (4)$$

for $i = 1, \dots, m$ and some $\alpha \in \mathbf{R}$;

(C3) Any solution $u \in \mathcal{X}$ of (4) (and not only any minimizer!) satisfies the Pohozaev identity

$$(N - p)J(u) = \alpha NV(u). \quad (5)$$

A few comments are in order. Clearly, the most important of the conditions above is **(C1)**. To our knowledge, the existence of a minimizer for (\mathcal{P}_1) , under sufficiently general assumptions on the functions g_i and for arbitrary $m \in \mathbf{N}^*$ and $p \in (1, \infty)$, is still an open problem. However, several particular cases have been extensively studied in the literature. A series of papers has been devoted to the case $p = 2$ and fairly optimal conditions on g_i that guarantee **(C1)** have been found by Berestycki-Lions [1] for $m = 1$ and by Brezis-Lieb [3] for $m \geq 1$. In the case $m = 1$ and $1 < p < N$ the existence of a minimizer for (\mathcal{P}_1) has also been proved in [9] under general assumptions on $g = g_1$ (similar to the assumptions in [1]). Under the conditions considered in [1] and [2], the functionals J and V are well defined on $H^1(\mathbf{R}^N)$ and this is clearly an admissible function space. The settings in [3] and [9] also correspond to our assumptions.

If $T > 0$ and (\mathcal{P}_1) admits minimizers, in most applications it is quite standard to prove that **(C2)** and **(C3)** hold. This is indeed the case under the assumptions in [1, 3, 9].

Next we consider the case $p = N$. Note that in this case the Pohozaev identity (5) becomes $\alpha NV(u) = 0$; hence any "reasonable" solution u of (1) should satisfy $V(u) = 0$. Since we are interested in nontrivial solutions, we consider the minimization problem

$$\text{minimize } J(u) \text{ in the set } \{ u \in \mathcal{X} \setminus \{0\} \mid V(u) = 0 \}. \quad (\mathcal{P}'_0)$$

We assume that the following conditions are satisfied.

- (D1)** $T_0 := \inf\{J(u) \mid u \in \mathcal{X}, u \neq 0, V(u) = 0\} > 0$ and (\mathcal{P}'_0) admits a minimizer u_0 ;
- (D2)** Any minimizer $u \in \mathcal{X}$ of (\mathcal{P}'_0) is C^1 and satisfies the Euler-Lagrange equations (4) for some $\alpha > 0$;
- (D3)** Any solution $u \in \mathcal{X}$ of (4) (with $\alpha > 0$) satisfies the Pohozaev identity $V(u) = 0$.

For $p = N = 2$, fairly optimal conditions on g_i that guarantee **(D1)**-**(D3)** have been found by Berestycki-Gallouët-Kavian [2] for $m = 1$ and by Brezis-Lieb [3] for $m \geq 1$.

In the next section we show that least energy solutions are minimizers of (\mathcal{P}_λ) for some particular choice of λ if $1 < p < N$, respectively minimizers of (\mathcal{P}'_0) if $p = N$. Then we obtain the radial symmetry of such solutions as a direct consequence of the general results in [15] (in the case $N = p$, we need some extra-argument in addition to the results in [15]).

In the third section we consider the scalar case $m = 1$ and we prove that least energy solutions have constant sign and, if they tend to zero at infinity, then they are monotone with respect to the radial variable.

In the final section we make some connections with related results of symmetry and monotonicity in the literature. Let us just mention that, especially in the scalar case, the symmetry and monotonicity of solutions of (1) have been studied by many authors, see

e.g. [11, 16, 8, 7] and references therein. In most of these works it is assumed that the solutions are nonnegative and some further assumptions on the nonlinearity g are made. They require, at least, g to be Lipschitz continuous and to satisfy a cooperative condition in the case of systems.

In the present work, we do not make any additional assumptions on g , except those that guarantee the existence of least energy solutions (basically, we need g to be merely continuous and to satisfy some growth conditions near zero and infinity, but we do not need any sign or monotonicity assumption; see [3] and [9]). We prove that our solutions have constant sign and our results are valid as well for compactly supported solutions and for solutions that do not vanish. Of course, there is a price we have to pay: our method works only for least energy solutions, not for any nonnegative solution of (1).

2 Variational characterization and symmetry

We begin with the case $1 < p < N$.

Lemma 1 *Assume that $1 < p < N$ and the conditions (C1)-(C3) hold.*

- (i) *Let u be a minimizer for (\mathcal{P}_1) . Then u_{σ_0} is a least action solution of (1), where $\sigma_0 = \left(\frac{N-p}{N}T\right)^{\frac{1}{p}}$, and $S(u_{\sigma_0}) = p(N-p)^{\frac{N}{p}-1}N^{-\frac{N}{p}}T^{\frac{N}{p}}$.*
- (ii) *Let v be a least energy solution for (1). Then v is a minimizer for (\mathcal{P}_λ) , where $\lambda = \left(\frac{N-p}{N}T\right)^{\frac{N}{p}}$.*

Proof. (i) By (C2) we know that $u \in C^1$ and u satisfies (4) for some $\alpha \in \mathbf{R}$. Then (5) implies $(N-p)J(u) = \alpha NV(u)$, which gives $\alpha = \frac{N-p}{N}T > 0$. It is easy to see that u_{σ_0} satisfies (1) for $\sigma_0 = \alpha^{\frac{1}{p}}$ and

$$S(u_{\sigma_0}) = \sigma_0^{N-p}J(u) - \sigma_0^N V(u) = \sigma_0^{N-p}T - \sigma_0^N = p(N-p)^{\frac{N}{p}-1}N^{-\frac{N}{p}}T^{\frac{N}{p}}.$$

Let $w \in \mathcal{X}$, $w \neq 0$, be a solution of (1). By (C3) we have $(N-p)J(w) = NV(w)$. If $J(w) = 0$, we have $\nabla w = 0$ a.e. on \mathbf{R}^N , hence w must be constant. Since $\text{measure}\{x \in \mathbf{R}^N \mid |w(x)| > \alpha\} < \infty$ for any $\alpha > 0$, we infer that $w = 0$, a contradiction. Thus $J(w) > 0$ and $V(w) = \frac{N-p}{N}J(w) > 0$. On the other hand, by (3) we get $J(w) \geq T(V(w))^{\frac{N-p}{N}}$, i.e. $J(w) \geq T\left(\frac{N-p}{N}J(w)\right)^{\frac{N-p}{N}}$, which gives

$$J(w) \geq \left(\frac{N-p}{N}\right)^{\frac{N-p}{p}} T^{\frac{N}{p}}. \quad (6)$$

Combined with Pohozaev identity, this implies

$$S(w) = J(w) - V(w) = \frac{p}{N}J(w) \geq p(N-p)^{\frac{N}{p}-1}N^{-\frac{N}{p}}T^{\frac{N}{p}} = S(u_{\sigma_0}) \quad (7)$$

and we infer that u_{σ_0} is a least energy solution for (1).

(ii) Conversely, let v be a least energy solution for (1). Then $(N-p)J(v) = NV(v)$ by (C3), hence $S(v) = \frac{p}{N}J(v)$. It is obvious that the inequalities (6) and (7) above are satisfied with $w = v$. On the other hand, $S(v) = S(u_{\sigma_0})$ and we infer that v must satisfy (7) with equality sign, that is,

$$J(v) = \left(\frac{N-p}{N}\right)^{\frac{N-p}{p}} T^{\frac{N}{p}} \quad \text{and} \quad V(v) = \frac{N-p}{N}J(v) = \left(\frac{N-p}{N}\right)^{\frac{N}{p}} T^{\frac{N}{p}}.$$

A simple scaling argument shows that v is a minimizer for (\mathcal{P}_λ) , where $\lambda = \left(\frac{N-p}{N}\right)^{\frac{N}{p}} T^{\frac{N}{p}}$; equivalently, v_{σ_1} is a minimizer for (\mathcal{P}_1) , where $\sigma_1 = \left(\frac{N-p}{N}T\right)^{-\frac{1}{p}} = \sigma_0^{-1}$. This completes the proof of Lemma 1. \square

The symmetry of least energy solutions will follow from Lemma 1 and a general symmetry result in [15]. For the convenience of the reader, we recall here that result.

Theorem 2 ([15]) *Let $N \geq 2$. Assume that $u : \mathbf{R}^N \rightarrow \mathbf{R}^m$ belongs to some function space \mathcal{Y} and solves the minimization problem*

$$\begin{aligned} & \text{minimize } \int_{\mathbf{R}^N} F(u(x), |\nabla u(x)|) dx \\ & \text{in the set } \left\{ u \in \mathcal{Y} \mid \int_{\mathbf{R}^N} H(u(x), |\nabla u(x)|) dx = \lambda \neq 0 \right\}. \end{aligned} \quad (\mathcal{P})$$

Suppose that the following conditions are satisfied:

- (A1) *For any $v \in \mathcal{Y}$ and any affine hyperplane Π in \mathbf{R}^N we have $v_{\Pi^+}, v_{\Pi^-} \in \mathcal{Y}$.*
- (A2) *Problem (\mathcal{P}) admits minimizers in \mathcal{Y} and any minimizer is a C^1 function on \mathbf{R}^N .*

Then, after a translation, u is radially symmetric.

Lemma 1 implies that least energy solutions solve the minimization problem (\mathcal{P}_λ) for some $\lambda > 0$. Conditions **(C1)**, **(C2)** and property **(v)** in the definition of admissible spaces imply that (\mathcal{P}_λ) satisfies the assumptions of Theorem 2. Thus we get:

Proposition 3 *Assume that $1 < p < N$ and **(C1)**-**(C3)** hold. Then (1) admits a least energy solution and each least energy solution is radially symmetric (up to a translation in \mathbf{R}^N).*

Now we turn our attention to the case $p = N$.

Proposition 4 *Assume that $p = N$ and **(D1)**-**(D3)** hold. Then (1) admits a least energy solution and any least energy solution solves (\mathcal{P}'_0) .*

Moreover, if we assume that G is either negative or positive in some ball $B_{\mathbf{R}^m}(0, \varepsilon) \setminus \{0\}$ and $u \in \mathcal{X}$ is a least energy solution such that $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$, then u is radially symmetric (up to a translation in \mathbf{R}^N).

Proof. Let u_0 be a minimizer for (\mathcal{P}'_0) . By **(D2)** and **(D3)** we have $V(u_0) = 0$ and u_0 satisfies (4) for some $\alpha > 0$. Let $u_1 = (u_0)_\sigma$, where $\sigma = \alpha^{\frac{1}{p}}$. It is easy to see that u_1 solves (1) and $S(u_1) = J(u_1) - V(u_1) = J(u_0) - \sigma^N V(u_0) = J(u_0) = T_0$. For any solution $u \in \mathcal{X}$, $u \neq 0$ of (1) we have $V(u) = 0$ by **(D3)** and $S(u) = J(u) \geq T_0 = J(u_1)$. Hence u_1 is a least energy solution.

If v is a least energy solution, then $V(v) = 0$ by **(D3)** and $J(v) = S(v) = S(u_1) = T_0$, thus v solves (\mathcal{P}'_0) .

Although Theorem 2 does not apply directly to minimizers of problem (\mathcal{P}'_0) (because the value of the constraint in (\mathcal{P}'_0) is zero), its proof can still be adapted to those minimizers. Indeed, the only place where the assumption $\lambda \neq 0$ is needed in Theorem 2 is to show that for any $e \in S^{N-1}$ there exists an affine hyperplane Π orthogonal to e such that

$$\int_{\Pi^-} H(u(x), |\nabla u(x)|) dx = \int_{\Pi^+} H(u(x), |\nabla u(x)|) dx = \frac{\lambda}{2}. \quad (8)$$

From (8) it follows then easily that u_{Π^+} and u_{Π^-} are also minimizers. (In fact, if $N = 2$ the assumption $\lambda \neq 0$ was also used in the proof of Theorem 2 to show that a minimizer u of (\mathcal{P}) could not be of the form $u(x) = \tilde{u}(|x|)$ on \mathbf{R}^2 , with $\tilde{u} : [0, \infty) \rightarrow \mathbf{R}^m$ periodic and nonconstant. In our setting it is clear that no minimizer u of (\mathcal{P}'_0) could be of this form because $J(u)$ is finite.)

In the present case we will use the fact that $G(u)$ has a constant sign in a neighborhood of ∞ to find hyperplanes that “split the constraint in two equal parts.” A similar idea has already been used in [14]. Henceforth we assume that u is a least action solution, $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$ and, say, $G(\xi) < 0$ for $0 < |\xi| < \varepsilon$. For $e \in S^{N-1}$ and $t \in \mathbf{R}$, we denote $\Pi_{e,t} = \{x \in \mathbf{R}^N \mid x \cdot e = t\}$, $\Pi_{e,t}^- = \{x \in \mathbf{R}^N \mid x \cdot e < t\}$ and $\Pi_{e,t}^+ = \{x \in \mathbf{R}^N \mid x \cdot e > t\}$. We claim that for any $e \in S^{N-1}$, there exists $t_e \in \mathbf{R}$ such that

$$\int_{\Pi_{e,t_e}^-} G(u(x)) dx = \int_{\Pi_{e,t_e}^+} G(u(x)) dx = 0 \quad \text{and} \quad u_{\Pi_{e,t_e}^-} \neq 0, u_{\Pi_{e,t_e}^+} \neq 0. \quad (9)$$

To see this, fix $e \in S^{N-1}$ and define $\varphi_e^\pm(t) = \int_{\Pi_{e,t}^\pm} G(u(x)) dx$, respectively. It follows that φ_e^+ and φ_e^- are continuous because $G(u) \in L^1(\mathbf{R}^N)$. Since u is continuous, $u \neq 0$, $\lim_{|x| \rightarrow \infty} u(x) = 0$ and $G < 0$ on $B_{\mathbf{R}^m}(0, \varepsilon) \setminus \{0\}$, it is not hard to see that there exist $t^-, t^+ \in \mathbf{R}$, $t^- < t^+$ such that

$$\varphi_e^-(t^-) < 0, \quad \varphi_e^+(t^+) < 0 \quad \text{and} \quad u_{\Pi_{e,t^-}^-} \neq 0, \quad u_{\Pi_{e,t^+}^+} \neq 0.$$

Since $\varphi_e^+(t^-) = V(u) - \varphi_e^-(t^-) = -\varphi_e^-(t^-)$, it follows that $\varphi_e^+(t^+) < 0 < \varphi_e^+(t^-)$. From the mean value property, we see that there exists $t_e \in (t^-, t^+)$ satisfying (9). It is clear that $u_{\Pi_{e,t_e}^-}, u_{\Pi_{e,t_e}^+} \in \mathcal{X} \setminus \{0\}$ because \mathcal{X} is admissible and (9) implies that $V(u_{\Pi_{e,t_e}^-}) = V(u_{\Pi_{e,t_e}^+}) = 0$, hence $J(u_{\Pi_{e,t_e}^-}) \geq T_0$, $J(u_{\Pi_{e,t_e}^+}) \geq T_0$. On the other hand, it is easy to see that $J(u_{\Pi_{e,t_e}^-}) + J(u_{\Pi_{e,t_e}^+}) = 2J(u) = 2T_0$. Thus $J(u_{\Pi_{e,t_e}^-}) = J(u_{\Pi_{e,t_e}^+}) = T_0$ and $u_{\Pi_{e,t_e}^-}, u_{\Pi_{e,t_e}^+}$ are also minimizers for (\mathcal{P}'_0) . Then arguing exactly as in the proof of Theorem 2 in [15], it follows that after a translation, u is radially symmetric. \square

Remark 5 The situation is different for $p > N$. The system (1) may still have solutions in some cases, and least energy solutions may also exist. For instance, if $N = 1$ and $p = 2$ it can be proved, under suitable assumptions on g , that (1) admits a finite energy solution which is unique up to translations; hence it is a least energy solution (and it is symmetric with respect to a point).

The existence and the symmetry of least energy solutions for (1) in the case $p > N \geq 2$ would be interesting problems to consider.

Note that whenever (1) admits finite energy solutions in the case $p > N$, they cannot admit a variational characterization as in Lemma 1 or Proposition 4 above. Indeed, any reasonable solution u of (1) should satisfy the Pohozaev identity $(N - p)J(u) = NV(u)$; if u is nontrivial, then necessarily $V(u) < 0$. It turns out that in any admissible function space \mathcal{X} , a condition like **(C1)** cannot hold for $p > N$, no matter what the nonlinearity g is. More precisely, denote

$$T_\lambda := \inf\{J(u) \mid u \in \mathcal{X} \text{ and } V(u) = \lambda\}.$$

Let $\lambda \neq 0$. We claim that either the set $\{u \in \mathcal{X} \mid V(u) = \lambda\}$ is empty (thus $T_\lambda = -\infty$), or we have $T_\lambda = 0$. To see this we argue by contradiction and we assume that there is some

$\lambda \neq 0$ such that $T_\lambda > 0$. Let $u \in \mathcal{X}$ be such that $V(u) = \lambda$ and $J(u) < 2T_\lambda$. Choose $\varepsilon > 0$ sufficiently small, so that $\varepsilon < \frac{1}{2}$ and $2\varepsilon^{\frac{p}{N}-1} < 1$. Let $\varphi(t) = \int_{\{x_1 < t\}} G(u) dx$. The function φ is continuous, $\lim_{t \rightarrow -\infty} \varphi(t) = 0$ and $\lim_{t \rightarrow \infty} \varphi(t) = \lambda$, hence there exist $t_1, t_2 \in \mathbf{R}$, $t_1 < t_2$ such that $\varphi(t_1) = \frac{\varepsilon}{2}\lambda$ and $\varphi(t_2) = (1 - \frac{\varepsilon}{2})\lambda$. Let $u_1(x) = \begin{cases} u(x) & \text{if } x_1 \leq t_1, \\ u(2t_1 - x_1, x') & \text{if } x_1 > t_1, \end{cases}$
 $u_2(x) = \begin{cases} u(x) & \text{if } x_1 > t_2, \\ u(2t_2 - x_1, x') & \text{if } x_1 \leq t_2, \end{cases}$ where $x' = (x_2, \dots, x_N)$. A simple change of variables shows that $V(u_1) = 2 \int_{\{x_1 < t_1\}} G(u) dx = \varepsilon\lambda$ and $V(u_2) = 2 \int_{\{x_1 > t_2\}} G(u) dx = \varepsilon\lambda$. Since $J(u_1) + J(u_2) = \frac{2}{p} \int_{\{x_1 < t_1\} \cup \{x_1 > t_2\}} \sum_{i=1}^m |\nabla u_i|^p dx \leq 2J(u)$, we see that $J(u_1) \leq J(u)$ or $J(u_2) \leq J(u)$. Assume that $J(u_1) \leq J(u) < 2T_\lambda$. For $\sigma = \varepsilon^{-\frac{1}{N}}$ we have $V((u_1)_\sigma) = \sigma^N V(u_1) = \lambda$ and $J((u_1)_\sigma) = \sigma^{N-p} J(u_1) \leq \sigma^{N-p} J(u) < \varepsilon^{\frac{p}{N}-1} 2T_\lambda < T_\lambda$, contradicting the definition of T_λ . Our claim is thus proved.

3 Monotonicity results

Throughout this section we assume that $m = 1$. Given a measurable function $f : \mathbf{R}^N \rightarrow [0, \infty)$ such that $\text{measure}(\{x \in \mathbf{R}^N \mid f(x) > \alpha\})$ is finite for any $\alpha > 0$, we denote by f^* the Schwarz rearrangement of f . We consider the following additional conditions for an admissible space \mathcal{X} .

- (vi) For any $u \in \mathcal{X}$ and $t \geq 0, s \leq 0$, we have $\min(u, t) \in \mathcal{X}$ and $\max(u, s) \in \mathcal{X}$.
- (vii) If $u \in \mathcal{X}$ and is a radial function and $u \geq 0$ (respectively $u \leq 0$), then $u^* \in \mathcal{X}$ (respectively $-(-u)^* \in \mathcal{X}$).

Note that assumption (vii) is needed only in the proof of Theorem 8 below.

Proposition 6 *Let \mathcal{X} be an admissible function space such that for any $v \in \mathcal{X}$ the functions $v_+ = \max(v, 0)$ and $v_- = \min(v, 0)$ belong to \mathcal{X} . Assume that $1 < p < N$ and (C1) holds. If $u \in \mathcal{X}$ is a solution of (\mathcal{P}_λ) for some $\lambda > 0$, then u does not change sign.*

Proof. This is a simple consequence of scaling. Indeed, let u be as above. It is clear that $V(u_+) + V(u_-) = V(u) = \lambda$ and $J(u_+) + J(u_-) = J(u)$. If $V(u_-) < 0$, then necessarily $V(u_+) > \lambda$. For $\sigma = \left(\frac{\lambda}{V(u_+)}\right)^{\frac{1}{N}} \in (0, 1)$ we have $V((u_+)_\sigma) = \sigma^N V(u_+) = \lambda$ and $J((u_+)_\sigma) = \sigma^{N-p} J(u_+) \leq \sigma^{N-p} J(u) < J(u)$, contradicting the fact that u is a minimizer. Thus necessarily $V(u_-) \geq 0$. In the same way $V(u_+) \geq 0$, therefore $V(u_-), V(u_+) \in [0, \lambda]$. Using inequality (3) (which trivially holds if $V(v) = 0$), we get

$$T\lambda^{\frac{N-p}{N}} = J(u) = J(u_+) + J(u_-) \geq TV(u_+)^{\frac{N-p}{N}} + TV(u_-)^{\frac{N-p}{N}},$$

which gives

$$1 \geq \left(\frac{V(u_+)}{\lambda}\right)^{\frac{N-p}{N}} + \left(\frac{V(u_-)}{\lambda}\right)^{\frac{N-p}{N}}. \quad (10)$$

Since $V(u_+) + V(u_-) = \lambda$, (10) implies that either $V(u_+) = 0$ or $V(u_-) = 0$. If $V(u_-) = 0$ and $V(u_+) = \lambda$ we see that u_+ satisfies the constraint and

$$J(u_+) = J(u) - J(u_-) \leq J(u). \quad (11)$$

Since u is a minimizer, we must have equality in (11) and this gives $J(u_-) = 0$, hence $u_- = 0$ and $u = u_+ \geq 0$. Similarly $V(u_+) = 0$ implies $u = u_- \leq 0$. \square

Proposition 7 *Let an admissible space \mathcal{X} satisfy the condition (vi). Assume that $p = N$ and (D1) holds. We have:*

- (a) *if $G < 0$ on $[-\varepsilon, 0) \cup (0, \varepsilon]$ for some $\varepsilon > 0$, then $u \in \mathcal{X}$ is a minimizer of (\mathcal{P}'_0) if and only if it solves the problem*

$$\text{minimize } J(v) \text{ in the set } \{v \in \mathcal{X} \mid v \neq 0, V(v) \geq 0\}; \quad (\mathcal{P}''_0)$$

- (b) *if $G > 0$ on $[-\varepsilon, 0) \cup (0, \varepsilon]$, then $u \in \mathcal{X}$ solves (\mathcal{P}'_0) if and only if it solves the problem*

$$\text{minimize } J(v) \text{ in the set } \{v \in \mathcal{X} \mid v \neq 0, V(v) \leq 0\}. \quad (\mathcal{P}'''_0)$$

Moreover, any minimizer of (\mathcal{P}''_0) or (\mathcal{P}'''_0) does not change sign.

Proof. It clearly suffices to prove (a).

Consider $v \in \mathcal{X}$ such that $v \geq 0$ a.e. and $V(v) > 0$. For $t \geq 0$ we define $v^t(x) = \min(v(x), t)$. By (vi) we have $v^t \in \mathcal{X}$. We claim that there exists $t_* > 0$ such that $V(v^{t_*}) = 0$.

The continuity of G , properties (i) and (iii) in the definition of admissible spaces and the dominated convergence theorem imply that the mapping $t \mapsto V(v^t) = \int_{\mathbf{R}^N} G(v^t(x)) dx$ is continuous on $(0, \infty)$. Since $G(v^\varepsilon(x)) < 0$ whenever $v(x) \neq 0$ and we cannot have $v(x) = 0$ a.e. because $V(v) > 0$, we infer that $V(v^\varepsilon) < 0$.

We claim that there exists $t_0 > \varepsilon$ such that $V(v^{t_0}) > 0$. Two situations may occur:

Case 1. There exists an increasing sequence $t_n \rightarrow \infty$ such that $\{G(t_n)\}_{n=1}^\infty$ is bounded from below. Let $m = \inf_{n \geq 1} G(t_n)$. By dominated convergence we get

$$V(v^{t_n}) - V(v) = \int_{\{v \geq t_n\}} G(t_n) - G(v(x)) dx \geq \int_{\{v \geq t_n\}} m - G(v(x)) dx \rightarrow 0 \quad \text{as } n \rightarrow \infty;$$

hence $V(v^{t_n}) \geq \frac{1}{2}V(v) > 0$ for n sufficiently large.

Case 2. $G(s) \rightarrow -\infty$ as $s \rightarrow \infty$. Then, since $v \geq 0$ a.e. and $V(v) > 0$, we see that the set $A = \{s > 0 \mid G(s) > 0\}$ is nonempty. Let $M = \sup A < \infty$. It follows that $G(s) \leq 0$ for $s \geq M$. It is clear that $M > \varepsilon$ and $V(v^M) \geq V(v) > 0$. The claim is thus proved.

Now the continuity of the mapping $t \mapsto V(v^t)$ implies that there exists $t_* \in (\varepsilon, t_0)$ such that $V(v^{t_*}) = 0$. Similarly, if $w \in \mathcal{X}$, $w \leq 0$ a.e. and $V(w) > 0$ there is some $\hat{t} > 0$ such that $V(-(-w)^{\hat{t}}) = 0$.

Next let $u_0 \in \mathcal{X}$ be a minimizer of (\mathcal{P}'_0) . Suppose $V(u) > 0$ for some $u \in \mathcal{X}$. Then at least one of the quantities $V(u_+)$ and $V(u_-)$ is positive. If $V(u_+) > 0$, take $t_* > 0$ such that $V(u_+^{t_*}) = 0$. We have $u_+^{t_*} \in \mathcal{X} \setminus \{0\}$ and

$$J(u) \geq J(u_+) \geq J(u_+^{t_*}) \geq J(u_0) = T_0. \quad (12)$$

Hence $\inf\{J(u) \mid u \in \mathcal{X}, u \neq 0, V(u) \geq 0\} = J(u_0) = T_0$ and u_0 is a solution of (\mathcal{P}''_0) .

Conversely, assume that u is a solution of (\mathcal{P}''_0) . We prove that

$$V(u_+) = V(u_-) = V(u) = 0. \quad (13)$$

We argue again by contradiction. If (13) does not hold, the inequality $V(u_+) + V(u_-) = V(u) \geq 0$ implies that at least one of the quantities $V(u_+)$ and $V(u_-)$ must be positive. Suppose that $V(u_+) > 0$. As above we find $t_* > 0$ such that $V(u_+^{t_*}) = 0$ and then (12) holds for u . Moreover, since u is a minimizer of (\mathcal{P}'_0) we have $J(u) \leq T_0$ and therefore all inequalities in (12) are in fact equalities. But $J(u_+) = J(u_+^{t_*})$ implies $\int_{\{u > t_*\}} |\nabla u|^p dx = 0$, hence $\nabla u = 0$ a.e. on $\{u > t_*\}$ which gives $\nabla((u - t_*)_+) = 0$ a.e. and we infer that $(u - t_*)_+ = 0$ a.e., that is $u \leq t_*$ a.e. Then we have $u_+ = u_+^{t_*}$ and consequently $V(u_+) = V(u_+^{t_*}) = 0$, contrary to our assumption. We argue similarly if $V(u_-) > 0$ and (13) is proved. Since $V(u) = 0$ and $J(u) = T_0 = J(u_0)$, we see that u solves (\mathcal{P}'_0) .

Lastly we show that if u is a minimizer of (\mathcal{P}''_0) , then either $u_+ = 0$ a.e. or $u_- = 0$ a.e. (but we cannot have $u_+ = u_- = 0$ a.e. because $J(u) = T_0 > 0$). Indeed, if $u^+ \neq 0$ and $u^- \neq 0$, (13) would imply $J(u_+) \geq T_0$ and $J(u_-) \geq T_0$ and this would give

$$T_0 = J(u) = J(u_+) + J(u_-) \geq 2T_0 > 0,$$

which is a contradiction. This completes the proof. \square

Next we prove the monotonicity of scalar minimizers.

Theorem 8 *Let \mathcal{X} be an admissible space satisfying the conditions (vi) and (vii). We assume that conditions (C1)-(C3) hold if $1 < p < N$, respectively conditions (D1)-(D3) hold if $p = N$. In the case $p = N$, we also assume that there exists $\varepsilon > 0$ such that either $G > 0$ or $G < 0$ on $[-\varepsilon, 0) \cup (0, \varepsilon]$. Then any least energy solution u of (1) such that $\lim_{|x| \rightarrow \infty} u(x) = 0$ is, up to a translation, radially symmetric and monotone with respect to $r = |x| \in [0, \infty)$.*

Proof. Symmetry follows directly from Propositions 3 and 4. Hence there is a function $\tilde{u} : [0, \infty) \rightarrow \mathbf{R}$ such that $u(x) = \tilde{u}(|x|) = \tilde{u}(r)$. From Lemma 1 and Proposition 4 we know that any least energy solution is a minimizer of (\mathcal{P}_λ) for some $\lambda > 0$, respectively of (\mathcal{P}'_0) . We will show that whenever $u(x) = \tilde{u}(r)$ solves one of these minimization problems and tends to zero at infinity, \tilde{u} is monotone on $[0, \infty)$. The proof relies on Lemma 9 below. The first part of this Lemma is well known and the second part is a simple consequence of Lemma 3.2 p. 163 in [6].

Lemma 9 ([6]) *Let w be a nonnegative measurable function defined on \mathbf{R}^N such that for any $t > 0$ the function $(w - t)_+$ belongs to $W^{1,p}(\mathbf{R}^N)$ and has compact support. Then we have*

$$\int_{\mathbf{R}^N} |\nabla w^*|^p dx \leq \int_{\mathbf{R}^N} |\nabla w|^p dx. \quad (14)$$

Moreover, if equality holds in (14) then for any $t \in (0, \text{sup ess}(w))$ the level set $\{x \in \mathbf{R}^N \mid w(x) > t\}$ is equivalent to a ball.

Now let u be as above. From Proposition 6 and Proposition 7, we know that u has constant sign; hence we may assume that $u \geq 0$. Since $u \in C^1$ and $\lim_{|x| \rightarrow \infty} u(x) = 0$, we see that u is bounded and $(u - t)_+$ belongs to $W^{1,p}(\mathbf{R}^N)$ and has compact support for any $t > 0$. By assumption (vii) we have $u^* \in \mathcal{X}$. It is clear that $V(u^*) = V(u)$, $u^* \neq 0$ if $u \neq 0$, and Lemma 9 implies that $J(u^*) \leq J(u)$. Since u is a minimizer of (\mathcal{P}_λ) (respectively of (\mathcal{P}'_0)), we have necessarily $J(u) \leq J(u^*)$, and hence $J(u) = J(u^*)$. Using Lemma 9 again we infer that for any $t \in (0, \text{sup}(u))$, the set $E_t = \{x \in \mathbf{R}^N \mid u(x) > t\}$ is equivalent to a ball.

If \tilde{u} is not nonincreasing, there exist $0 \leq r_1 < r_2$ such that $0 < \tilde{u}(r_1) < \tilde{u}(r_2)$. Since $\tilde{u}(x) \rightarrow 0$ as $|x| \rightarrow \infty$, there exists $r_3 > r_2$ such that $u(r_3) = u(r_1)$. Denoting $a = u(r_1)$ and $b = u(r_2)$, we see that for any $t \in (a, b)$, E_t is nonempty and is not equivalent to a ball, which is a contradiction. This completes the proof of Theorem 8. \square

4 Some remarks and examples

Remark 10 In the scalar case $m = 1$ it is well known (see for example the Introduction of [5]) that if g is odd then any least energy solution has a constant sign. In Remark II.6 of [12], Lions raised the question (for $p = 2$ and $N \geq 3$) whether this remains true without assuming g odd. Proposition 6 gives an affirmative answer for any $1 < p < N$ and Proposition 7, under some mild additional assumptions, for $p = N$. Previous partial results were obtained by Brock [5], using rearrangement arguments, assuming that $1 < p \leq 2$, the minimizer u satisfies $u(x) \rightarrow 0$ as $|x| \rightarrow \infty$ and $g \in C^{0,p-1}(\mathbf{R})$. Nothing was proved for $p > 2$.

Remark 11 If $N \geq 3$, $p = 2$, $m = 1$ and under the assumption that g is odd, the existence of least energy solutions for (1) has been proved in [1] by showing that problem (\mathcal{P}_1) admits a minimizer. The minimizer found in [1] was radial by construction, but it was not known whether all least energy solutions were radially symmetric. The existence of a minimizer for (\mathcal{P}_1) without the oddness assumption on g has also been proved in [12], but nothing was known about the symmetry or the sign of such minimizers. Our results imply that any least energy solution is radially symmetric, has constant sign and is monotone with respect to the radial variable, no matter whether g is odd or not.

In the case $N \geq 2$, $p = 2$, $m \in \mathbf{N}^*$, the existence of least energy solutions is also known (see [3] for general results, historical notes, comments and further references). If $N > 2$, the existence of a minimizer for (\mathcal{P}_λ) and the existence of least energy solutions have been proved in [3] under very general assumptions on the functions g_i . It has also been shown that the solutions are smooth (Theorem 2.3 p. 105 in [3]) and satisfy the Pohozaev identity (Lemma 2.4 p. 104 in [3]). However, as already mentioned in [3] p. 99, the existence of radially symmetric least energy solutions was not clear. Indeed, the Schwarz symmetrization that lead to a radial minimizer in [1] could not be used in [3] because of the general assumptions on the nonlinearity made there. In fact, it is known that the Schwarz rearrangements may be used for systems only if the nonlinearity satisfies a cooperative condition.

Proposition 3 above implies that all least energy solutions of the system considered in [3] are radially symmetric.

If $N = 2$ and $G(\xi) < 0$ for $0 < |\xi| \leq \varepsilon$, the existence of least energy solutions and the existence of minimizers for (\mathcal{P}'_0) have been proved in [2, 3]. It has also been shown that such solutions are smooth, satisfy the Pohozaev identity and tend to 0 as $|x| \rightarrow \infty$. Therefore Proposition 4 implies that any least energy solution is radially symmetric.

We have to mention that if $p = 2$ and if the minimizers of (\mathcal{P}_λ) satisfy a unique continuation principle, it has already been proved in [13] that any minimizer is radially symmetric (modulo translation). In [13] no cooperative condition is required when $m \geq 2$ but using a unique continuation principle require in particular g to be C^1 . Our results are still valid when a unique continuation principle fails (e.g., for minimizers with compact support). Note that compactly supported minimizers may occur in some applications (cf. Theorem 3.2 (ii) p. 111 in [3]; see also [15] for such an example). In the scalar case $m = 1$, [13] does not say anything about the sign of the minimizers.

However, in the case $p = 2$ and $N \geq 3$, the symmetry, positivity and monotonicity of minimizers for problem (\mathcal{P}_λ) have been proved in [10] in the "zero-mass case" (that is, when $g(0) = g'(0) = 0$). The proofs in [10] rely on some sharp estimates of the decay of solutions at infinity (which are a consequence of the "zero-mass" condition) and on a result in [6]. Note that in [10] it is not assumed that g is continuous on \mathbf{R} , but it is assumed that $g \geq 0$ on $(0, \infty)$ and $g \leq 0$ on $(-\infty, 0)$, respectively.

Remark 12 If $1 < p < N$ and $m = 1$, it has been proved in [9], under general conditions on g , that problem (\mathcal{P}_λ) admits minimizers (thus (1) has least energy solutions). The minimizers found in [9] were radially symmetric by construction. It follows from Proposition 3 that any least energy solution is radially symmetric.

If, in addition to the assumptions of Theorem 8, it is assumed that g is locally Lipschitz on $(0, \infty)$ and non-increasing on some interval $(0, s_0)$ and $1 < p < 2$, it has been proved in [8] that any nonnegative solution of (1) is radially symmetric and that $u(x) = \tilde{u}(|x|)$ satisfies $\tilde{u}'(r) < 0$ whenever $r > 0$ and $\tilde{u}(r) > 0$. The same result is true when $p > 2$ if it is assumed in addition that the critical set of the solution u is reduced to one point (see [16]). These assumptions are not necessary for us but, of course, we only deal with least energy solutions.

Remark 13 (i) The symmetry results in Section 2 hold without any change if we replace the functional J by a functional of the form $\int_{\mathbf{R}^N} \sum_{i=1}^m A_i(u, \nabla u_i) dx$ where $\xi \rightarrow A_i(u, \xi)$ is p -homogeneous for any $i = 1, \dots, m$.

(ii) Our method still works for more general functionals of the form

$$\tilde{J}(u) = \frac{1}{p} \int_{\mathbf{R}^N} |x|^\alpha \sum_{i=1}^m A_i(u) |\nabla u_i|^p dx \quad \text{and} \quad \tilde{V}(u) = \int_{\mathbf{R}^N} |x|^\beta G(u) dx.$$

In this case, using Theorem 1 in [15], we obtain that minimizers (and the corresponding minimum action solutions) are axially symmetric.

Functionals of this type appear, e.g., in the Caffarelli-Kohn-Nirenberg problem (which consists in minimizing $\int_{\mathbf{R}^N} |\nabla u|^q |x|^{-aq} dx$ under the constraint $\int_{\mathbf{R}^N} |u|^p |x|^{-bp} dx = \text{const.}$, where $q > 1$, $p > 1$, $a \leq b < \frac{N}{q}$ and $0 < \frac{1}{q} - \frac{1}{p} = \frac{1+a-b}{N}$). It has been proved that minimizers for this problem exist and, in general, are not radially symmetric (see [4] and references therein).

Acknowledgment The research of the first author was supported in part by KRF-2007-412-J02301 of Korea Research Foundation.

References

- [1] H. BERESTYCKI, P.-L. LIONS, *Nonlinear scalar field equations, I. Existence of a ground state*, Arch. Rational Mech. Anal. 82 (1983), 313-345.
- [2] H. BERESTYCKI, T. GALLOUËT, O. KAVIAN, *Equations de champs scalaires euclidiens non linéaires dans le plan*, C.R. Acad. Sc. Paris Série I - Math. 297 (1983), 307-310 and Publications du Laboratoire d'Analyse Numérique, Université de Paris VI, 1984.
- [3] H. BRÉZIS, E. H. LIEB, *Minimum Action Solutions of Some Vector Field Equations*, Comm. Math. Phys. 96 (1984), 97-113.

- [4] J. BYEON, Z.-Q. WANG, *Symmetry breaking of extremal functions for the Caffarelli-Kohn-Nirenberg inequalities*, Comm. Contemp. Math. 4 (2002), 457–465.
- [5] F. BROCK, *Positivity and radial symmetry of solutions to some variational problems in \mathbf{R}^N* , J. Math. Anal. Appl. 296 (2004), 226–243.
- [6] J. E. BROTHERS, W.P. ZIEMER, *Minimal rearrangements of Sobolev functions*, J. reine angew. Math. 384 (1988), 153-179.
- [7] J. BUSCA, B. SIRAKOV, *Symmetry results for semilinear elliptic systems in the whole space*, J. Diff. Eq. 163, No. 1 (2000), 41-56.
- [8] L. DAMASCELLI, F. PACELLA, M. RAMASWAMY, *Symmetry of ground states of p -Laplace equations via the moving plane method*, Arch. Rational Mech. Anal. 148 (1999), 291-308.
- [9] A. FERRERO, F. GAZZOLA, *On subcriticality assumptions for the existence of ground states of quasilinear elliptic equations*, Adv. Diff. Eq. 8, No. 9 (2003), 1081-1106.
- [10] M. FLUCHER, S. MÜLLER, *Radial symmetry and decay rate of variational ground states in the zero mass case*, SIAM. J. Math. Anal. 29, No. 3 (1998), 712-719.
- [11] B. GIDAS, W. N. NI, L. NIRENBERG, *Symmetry of positive solutions of nonlinear elliptic equations in \mathbf{R}^n* , Adv. Math. Supp. Stud. 7A (1981), 369-403.
- [12] P.L. LIONS, *The concentration-compactness principle in the Calculus of Variations, The locally compact case, Part 2*, Ann. Inst. H. Poincaré Anal. Non Linéaire 1 (1984), 223-283.
- [13] O. LOPES, *Radial symmetry of minimizers for some translation and rotation invariant functionals*, J. Diff. Eq. 124 (1996), 378-388.
- [14] O. LOPES, M. MONTENEGRO, *Symmetry of mountain pass solutions for some vector field equations*, J. Dyn. Diff. Eq. 18, No. 4 (2006), 991-999.
- [15] M. MARIŞ, *On the symmetry of minimizers*, Arch. Rational Mech. Anal., in press, DOI 10.1007/s00205-008-0136-2; arXiv:0712.3386 (www.arxiv.org)
- [16] J. SERRIN, H. ZOU, *Symmetry of ground states of quasilinear elliptic equations*, Arch. Rational Mech. Anal. 148 (1999), 265-290.