

KMAS9AA1 – Algebraic Topology


Exercise Sheet 4

1. Computation of Homologies

- 1) Let (M, m) and (N, n) be two pointed spaces with open neighbourhoods deformation retracting to the respective points. Show that $H_d(M \vee N, *) = H_d(M, n) \oplus H_d(N, n)$.
Remark: This condition is satisfied by manifolds, but also by CW complexes.
- 2) The *connected sum* $M \# N$ of two connected manifolds M and N of the same dimension n is obtained by removing a small neighborhood of a point¹ formed by an open disc from each, and gluing the resulting manifolds along the two spheres S^{n-1} that appear. For example, $\Sigma_g \# \Sigma_{g'} = \Sigma_{g+g'}$, where Σ_g is the oriented surface of genus g .
Show that for $i \neq n-1, n$, we have $\tilde{H}_i(M \# N) = \tilde{H}_i(M) \oplus \tilde{H}_i(N)$.
Show that if R is a field, the dimension of $\tilde{H}_i(M \# N)$ and $\tilde{H}_i(M) \oplus \tilde{H}_i(N)$ differ by at most 1.
- 3) Compute the homology of the torus, seeing it as square identified opposite edges. This can be done by taking the Mayer–Vietoris sequence on U being a small disc inside the square and V being a thickening of the complement of U , in such a way that $U \cap V \sim S^1$.
- 4) Compute the homology of Σ_2 . This can be done using the strategy of the last exercise, but also seeing $\Sigma_2 = \Sigma_1 \# \Sigma_1$.
- 5) Let $R = \mathbb{Z}$. Suppose that a topological space is written as a union of two open subspaces $X = U \cup V$ and consider the associated Mayer–Vietoris long exact sequence. Assuming that $U \cap V$ is path connected, use the explicit construction of the connecting morphism $\delta: H_1(X) \rightarrow H_0(U \cap V)$ to show that is the zero map.
- 6) Compute the homology of $S^n \times S^m$.
- 7) Compute the homology $H(\Sigma_2, A)$ and $H(\Sigma_2, B)$, where A and B are the following circles²:

¹Up to homeomorphism it doesn't matter which point we choose.

²Drawing from [H, Exercise 2.1.17]



2. Products, coproducts, pullbacks and pushouts

Let \mathcal{C} be a category.

- 1) Recall the notions in the title of this exercise and show that if they exist they are unique up to unique isomorphism.
- 2) Show that the category of fields does not have all coproducts.
- 3) Show that in the category of unital commutative rings, the coproduct of R and S is given by $R \otimes S$ with the maps $R \rightarrow R \otimes S, r \mapsto r \otimes 1$ and $S \rightarrow R \otimes S, s \mapsto 1 \otimes s$.
- 4) We say that $I \in \mathcal{C}$ is an *initial* object if for any object $X \in \mathcal{C}$, there is a unique morphism $I \rightarrow X$. Similarly, $T \in \mathcal{C}$ is *terminal* if there is a unique morphism from any $X \rightarrow T$.
 - a. Show that if such objects exist, they are unique.
 - b. Show that if an initial object exists, any coproduct can be written as a pushout. Similarly, if a terminal object exists, a product is a pullback.
 - c. Determine the initial and terminal object in the categories **Top**, **Top_{*}**, **Groups**, **$R - \text{Mod}$** .
- 5) In **Set**, show that the pullback of $f: X \rightarrow Z$ and $g: Y \rightarrow Z$ is given by the set of pairs $X \times_Z Y = \{(x, y) | x \in X, y \in Y, f(x) = g(y)\}$.

3. CW Complexes

- 1) Show that a single cell attachment is a pushout along the attachment map.
- 2) Let X, Y be CW complexes containing finitely many cells. Show that $X \times Y$ is a CW complex with d -cells given by products of k and $d - k$ cells.
- 3) Let X, Y be CW complexes, A a subcomplex of X and $f: A \rightarrow Y$ a cellular map. Show that the pushout of f along the inclusion $A \hookrightarrow X$ is a CW complex with the cells of Y and of $X - A$ as cells. You can assume for convenience that all CW complexes have finitely many cells (and therefore it is enough to consider the case of a single cell attachment)
- 4) We constructed in class a cellular structure on \mathbb{RP}^n as the quotient of a cellular structure on S^n . Describe with precise formulas for the attaching maps the cellular structure of \mathbb{RP}^2 , without using quotients at any point.

- 5) Let Γ be a finite graph with e edges, v vertices and c connected components.

Compute $H_\bullet(\Gamma; \mathbb{Z})$ in terms of e, v and c .

Hint: Use the fact that given a CW-complex X and A a contractible subcomplex $X \rightarrow X/A$ is a homotopy equivalence.

4. Chain Complexes [This exercise was supposed to appear on an earlier exercise sheet, but didn't by mistake]

- 1) Show that the kernel, image, and cokernel of a chain complex morphism $f: C \rightarrow D$ are chain complexes. Show that if $H(\ker f) = 0 = H(\operatorname{coker} f)$, then f induces an isomorphism in homology.
- 2) Show that the direct sum of chain complexes is a chain complex, and compare $H(A \oplus B)$ with $H(A) \oplus H(B)$.
- 3) Compute the homology of the complex

$$0 \rightarrow R \xrightarrow{\times 2} R \rightarrow 0$$

for $R = \mathbb{Z}$ and R a field (pay attention to the characteristic). [Important to retain from this: The ring we are working with changes a lot the result. Hatcher only works with \mathbb{Z} which does not allow us to see such differences later on.]

- 4) Show that the dual of a chain complex $C_\bullet^\vee = \{\operatorname{Hom}_R(C_{-i}, R)\}_{i \in \mathbb{Z}}$ is a chain complex with boundary map given by the dual of the original one. Prove that over \mathbb{Z} , the dual of the homology of C is not isomorphic to the homology of the dual of C .

Hint: The subexercise just above.