

# KMAS9AA1 – Algebraic Topology

## Exercise Sheet 6

### 1. Tor

- 1) Check carefully that the fundamental theorem of homological algebra implies that for any two resolutions of the  $R$ -module  $M$ ,  $F_\bullet \rightarrow M$  and  $F'_\bullet \rightarrow M$ , we have that  $H_n(F_\bullet \otimes B)$  is canonically isomorphic to  $H_n(F'_\bullet \otimes B)$ .
- 2) Show that  $\text{Tor}_i^r(A \oplus B, C) = \text{Tor}(A, C) \oplus \text{Tor}(B, C)$ .

Recall that given an exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ , there is a long exact sequence of Tor functors given by tensoring this short exact sequence with  $F_\bullet$ , a resolution of  $M$ . From now on, assume that  $R$  is a PID.

- 3) Take a free resolution  $E_\bullet \rightarrow C$  such that  $E_i = 0$  for  $i \geq 2$ . Consider the long exact sequence associated to

$$0 \rightarrow E_1 \otimes F_\bullet \rightarrow E_0 \otimes F_\bullet \rightarrow C \otimes F_\bullet \rightarrow 0$$

to conclude that  $\text{Tor}_1(C, M) = \text{Tor}_1(M, C)$ .

- 4) Assume that  $H_n(X; \mathbb{Z})$  and  $H_{n-1}(X; \mathbb{Z})$  are finitely generated. Show that for any prime  $p$ ,  $H_n(X; \mathbb{Z}/p\mathbb{Z})$  consists of:
  - i. A  $\mathbb{Z}/p\mathbb{Z}$  summand for each  $\mathbb{Z}$  summand of  $H_n(X; \mathbb{Z})$ ,
  - ii. A  $\mathbb{Z}/p\mathbb{Z}$  summand for each  $\mathbb{Z}/p^k\mathbb{Z}$  summand of  $H_n(X; \mathbb{Z})$ ,
  - iii. A  $\mathbb{Z}/p\mathbb{Z}$  summand for each  $\mathbb{Z}/p^k\mathbb{Z}$  summand of  $H_{n-1}(X; \mathbb{Z})$ ,
- 5) Use the universal coefficient theorem to show that if  $H_*(X; \mathbb{Z})$  is finitely generated, so the Euler characteristic

$$\chi(X) = \sum_n (-1)^n \text{rank} H_n(X; \mathbb{Z})$$

is defined, then for any coefficient field  $\mathbb{F}$  we have  $\chi(X) = \sum_n (-1)^n \dim H_n(X; \mathbb{F})$ .

- 2. Torsion-free** I claimed in class that while  $\mathbb{Q}$  is not free, it is torsion-free and therefore  $\text{Tor}_1^{\mathbb{Z}}(\mathbb{Q}, C) = 0, \forall C \in R - \text{Mod}$ . Let us show this.

Let  $G$  be an abelian group.

- 1) Show that any element of  $G \otimes \mathbb{Q}$  is of the form  $g \otimes \frac{1}{n}$ .
- 2) Show that if  $G$  is a torsion group, then  $\mathbb{Q} \otimes G = 0$ .
- 3) Show that if  $G$  is torsion free, then  $g \otimes \frac{1}{n} = g' \otimes \frac{1}{n'}$  is equivalent to  $gn' = ng'$ .
- 4) Take a free resolution  $F_1 \rightarrow F_0$  of  $G$ . Show that  $F_1 \otimes \mathbb{Q} \rightarrow F_0 \otimes \mathbb{Q}$  is injective and conclude that  $\text{Tor}_1^{\mathbb{Z}}(\mathbb{Q}, G) = 0$ .

### 3. Ext

- 1) Show if  $A \rightarrow B \rightarrow C \rightarrow 0$  is exact, then  $\text{Hom}(A, N) \leftarrow \text{Hom}(B, N) \leftarrow \text{Hom}(C, N) \leftarrow 0$  is exact.  
[This is what is used to conclude that  $\text{Ext}_R^0(M, N) = \text{Hom}_R(M, N)$ !]
- 2) Show that  $\text{Ext}_R^i(A \oplus B, N) = \text{Ext}_R^i(A, N) \oplus \text{Ext}_R^i(B, N)$  and  $\text{Ext}_R^i(R^7, N) = 0$ .
- 3) Show that  $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}/n\mathbb{Z}, N) = N/nN$ .
- 4) Show that  $\text{Ext}_R^1(M, -)$  is a covariant functor and that  $\text{Ext}^1(-, N)$  is a contravariant functor.
- 5) Show that  $\text{Ext}_{\mathbb{Z}/4\mathbb{Z}}^n(\mathbb{Z}/2\mathbb{Z}, \mathbb{Z}/2\mathbb{Z}) = \mathbb{Z}/2\mathbb{Z}$ .

### 4. Cup product

- 1) Show that  $H^\bullet(X \sqcup Y; R)$  and  $H^\bullet(X; R) \oplus H^\bullet(Y; R)$  are isomorphic as graded commutative  $R$ -algebras. Deduce a similar statement for the wedge product (assuming that the basepoints are deformation retracts of open neighbourhoods).
- 2) Let  $X$  be a CW complex with one 0-cell, one 5-cell, one 7-cell and one 10-cell. What is the cohomology ring structure of  $X$  with coefficients in  $\mathbb{Q}$ ?

The cohomology of the torus with coefficients in  $\mathbb{F}_2$  is spanned by **degree 0:** 1, **degree 1:**  $\alpha, \beta$  and **degree 2:**  $\gamma$ .

- 3) Use the same strategy that we used in class for  $\mathbb{RP}^2$  to show that  $\alpha \cup \beta = \gamma$ .
- 4) Show that  $\alpha \cup \alpha = 0$ .